



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2024 – 09:41 PM EDT

PDB ID : 5LIQ  
Title : The structure of C160S,C508S,C578S mutant of Nt.BspD6I nicking endonuclease at 0.185 nm resolution .  
Authors : Kachalova, G.S.; Artyukh, R.I.; Perevyazova, T.A.; Yunusova, A.K.; Popov, A.N.; Bartunik, H.D.; Zheleznaya, L.A.  
Deposited on : 2016-07-15  
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	610	
1	B	610	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	702	-	-	X	-
2	GOL	A	706	-	-	X	-
3	PO4	B	705	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nicking endonuclease N.BspD6I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	583	Total	C	N	O	S	0	13	0
			4920	3167	826	913	14			
1	B	583	Total	C	N	O	S	0	5	0
			4870	3133	822	902	13			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q8GCA3
A	-4	HIS	-	expression tag	UNP Q8GCA3
A	-3	HIS	-	expression tag	UNP Q8GCA3
A	-2	HIS	-	expression tag	UNP Q8GCA3
A	-1	HIS	-	expression tag	UNP Q8GCA3
A	0	HIS	-	expression tag	UNP Q8GCA3
A	160	SER	CYS	engineered mutation	UNP Q8GCA3
A	508	SER	CYS	engineered mutation	UNP Q8GCA3
A	578	SER	CYS	engineered mutation	UNP Q8GCA3
B	-5	HIS	-	expression tag	UNP Q8GCA3
B	-4	HIS	-	expression tag	UNP Q8GCA3
B	-3	HIS	-	expression tag	UNP Q8GCA3
B	-2	HIS	-	expression tag	UNP Q8GCA3
B	-1	HIS	-	expression tag	UNP Q8GCA3
B	0	HIS	-	expression tag	UNP Q8GCA3
B	160	SER	CYS	engineered mutation	UNP Q8GCA3
B	508	SER	CYS	engineered mutation	UNP Q8GCA3
B	578	SER	CYS	engineered mutation	UNP Q8GCA3

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	186	Total	O	0	0
			186	186		
4	B	109	Total	O	0	0
			109	109		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.20Å 92.48Å 113.81Å 90.00° 105.67° 90.00°	Depositor
Resolution (Å)	19.58 – 1.85 19.56 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (19.58-1.85) 99.6 (19.56-1.85)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.209 , 0.251 0.209 , 0.251	Depositor DCC
$R_{free}$ test set	6247 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.2	Xtrriage
Anisotropy	0.055	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10168	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 75.75 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1906e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.98	1/5041 (0.0%)	0.96	8/6782 (0.1%)
1	B	0.96	1/4973 (0.0%)	0.98	11/6692 (0.2%)
All	All	0.97	2/10014 (0.0%)	0.97	19/13474 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	461	TYR	CE1-CZ	6.94	1.47	1.38
1	B	8	TYR	CE1-CZ	-5.11	1.31	1.38

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	320	ARG	NE-CZ-NH2	-9.19	115.71	120.30
1	B	203[A]	ARG	NE-CZ-NH1	-8.55	116.02	120.30
1	B	203[B]	ARG	NE-CZ-NH1	-8.55	116.02	120.30
1	A	580	ARG	NE-CZ-NH1	8.35	124.47	120.30
1	A	203	ARG	NE-CZ-NH1	-6.92	116.84	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4920	0	4986	101	0
1	B	4870	0	4921	84	0
2	A	36	0	48	18	0
2	B	12	0	16	3	0
3	A	20	0	0	0	0
3	B	15	0	0	2	0
4	A	186	0	0	16	0
4	B	109	0	0	10	0
All	All	10168	0	9971	188	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 188 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:GLU:OE2	1:A:203:ARG:NH1	1.64	1.28
1:B:175:GLU:OE2	1:B:203[A]:ARG:NH1	1.73	1.21
1:A:531:ASN:ND2	2:A:706:GOL:H11	1.56	1.19
1:B:581:LYS:HG2	1:B:584:GLN:NE2	1.58	1.19
1:A:310:LEU:CD1	1:A:313:ARG:HH21	1.54	1.19

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	590/610 (97%)	578 (98%)	12 (2%)	0	100	100
1	B	582/610 (95%)	565 (97%)	15 (3%)	2 (0%)	41	26
All	All	1172/1220 (96%)	1143 (98%)	27 (2%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	186	ILE
1	B	225	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	545/554 (98%)	541 (99%)	4 (1%)	84 79
1	B	536/554 (97%)	534 (100%)	2 (0%)	91 89
All	All	1081/1108 (98%)	1075 (99%)	6 (1%)	86 83

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	327	GLU
1	B	67	PHE
1	B	579	GLU
1	A	200	PHE
1	A	68	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	375	GLN
1	B	478	GLN
1	A	395	HIS
1	A	375	GLN
1	B	516	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	A	708	-	4,4,4	0.98	0	6,6,6	1.01	0
2	GOL	A	702	-	5,5,5	0.44	0	5,5,5	1.31	1 (20%)
2	GOL	A	703	-	5,5,5	0.64	0	5,5,5	0.62	0
3	PO4	A	707	-	4,4,4	1.27	1 (25%)	6,6,6	1.13	0
2	GOL	B	702	-	5,5,5	0.48	0	5,5,5	0.92	0
2	GOL	A	701	-	5,5,5	0.37	0	5,5,5	0.80	0
2	GOL	A	706	-	5,5,5	0.59	0	5,5,5	0.86	0
3	PO4	A	710	-	4,4,4	0.99	0	6,6,6	0.36	0
2	GOL	B	701	-	5,5,5	0.45	0	5,5,5	1.38	1 (20%)
3	PO4	B	704	-	4,4,4	1.35	1 (25%)	6,6,6	0.93	0
3	PO4	A	709	-	4,4,4	0.82	0	6,6,6	0.89	0
3	PO4	B	703	-	4,4,4	1.33	1 (25%)	6,6,6	1.37	1 (16%)
3	PO4	B	705	-	4,4,4	0.94	0	6,6,6	1.32	0
2	GOL	A	704	-	5,5,5	0.42	0	5,5,5	0.76	0
2	GOL	A	705	-	5,5,5	0.50	0	5,5,5	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	702	-	-	4/4/4/4	-
2	GOL	A	703	-	-	0/4/4/4	-
2	GOL	B	702	-	-	3/4/4/4	-
2	GOL	A	701	-	-	2/4/4/4	-
2	GOL	A	706	-	-	1/4/4/4	-
2	GOL	B	701	-	-	2/4/4/4	-
2	GOL	A	704	-	-	3/4/4/4	-
2	GOL	A	705	-	-	4/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	704	PO4	P-O1	2.32	1.56	1.50
3	B	703	PO4	P-O1	2.27	1.56	1.50
3	A	707	PO4	P-O3	-2.13	1.48	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	703	PO4	O4-P-O1	-2.37	102.22	110.89
2	B	701	GOL	O3-C3-C2	-2.26	99.35	110.20
2	A	702	GOL	C3-C2-C1	-2.02	103.87	111.70

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	GOL	O1-C1-C2-C3
2	A	702	GOL	O1-C1-C2-C3
2	A	702	GOL	C1-C2-C3-O3
2	A	704	GOL	C1-C2-C3-O3
2	A	705	GOL	O1-C1-C2-C3

There are no ring outliers.

8 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	702	GOL	6	0
2	A	703	GOL	2	0
2	B	702	GOL	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	GOL	1	0
2	A	706	GOL	8	0
2	B	701	GOL	1	0
3	B	705	PO4	2	0
2	A	704	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	583/610 (95%)	0.30	41 (7%) 16   15	25, 41, 78, 130	7 (1%)
1	B	583/610 (95%)	0.48	56 (9%) 8   7	26, 46, 81, 151	6 (1%)
All	All	1166/1220 (95%)	0.39	97 (8%) 11   11	25, 44, 81, 151	13 (1%)

The worst 5 of 97 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	67	PHE	15.5
1	B	66	ALA	14.3
1	A	67	PHE	13.1
1	B	58	GLY	10.8
1	B	57	LEU	10.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	A	706	6/6	0.57	0.27	41,61,65,70	6
3	PO4	B	705	5/5	0.73	0.33	40,40,56,59	5
2	GOL	A	705	6/6	0.75	0.26	66,69,81,84	0
3	PO4	B	704	5/5	0.78	0.35	47,50,54,56	5
2	GOL	A	704	6/6	0.79	0.15	42,54,56,63	6
2	GOL	A	703	6/6	0.83	0.46	47,57,64,72	6
3	PO4	A	708	5/5	0.85	0.45	56,57,59,62	5
2	GOL	A	702	6/6	0.87	0.33	39,60,62,65	0
2	GOL	A	701	6/6	0.89	0.13	50,56,58,63	6
2	GOL	B	701	6/6	0.89	0.14	45,52,54,60	6
2	GOL	B	702	6/6	0.89	0.28	50,67,71,76	0
3	PO4	A	709	5/5	0.90	0.30	73,73,75,83	5
3	PO4	A	710	5/5	0.91	0.24	77,81,84,88	5
3	PO4	A	707	5/5	0.93	0.14	39,52,64,71	0
3	PO4	B	703	5/5	0.95	0.16	40,51,59,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.