



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 24, 2023 – 12:13 PM EST

PDB ID : 3LM2
Title : Crystal structure of Putative kinase. (17743352) from AGROBACTERIUM TUMEFACIENS str. C58 (Dupont) at 1.70 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2010-01-29
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	226	 88% 12%
1	B	226	 89% 9%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3911 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative kinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	225	1772	1123	306	327	2	14	0	15	0
1	B	222	1747	1106	298	329	2	12	0	14	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A9CH74
B	0	GLY	-	expression tag	UNP A9CH74

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	186	187	187	0	1
3	B	194	197	197	0	3

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative kinase

Chain A:  88% 12%



- Molecule 1: Putative kinase

Chain B:  89% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	47.91Å 49.60Å 57.33Å 97.98° 104.64° 109.97°	Depositor
Resolution (Å)	29.13 – 1.70 29.13 – 1.70	Depositor EDS
% Data completeness (in resolution range)	94.5 (29.13-1.70) 94.6 (29.13-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.5.0102, PHENIX	Depositor
R, R_{free}	0.160 , 0.186 0.207 , 0.227	Depositor DCC
R_{free} test set	2471 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtrriage
Anisotropy	0.592	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3911	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	2/1840 (0.1%)	0.80	1/2466 (0.0%)
1	B	0.68	0/1813	0.72	0/2434
All	All	0.71	2/3653 (0.1%)	0.76	1/4900 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	136	GLU	CD-OE2	5.67	1.31	1.25
1	A	136	GLU	CB-CG	5.03	1.61	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	ASP	CB-CG-OD1	8.10	125.59	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1772	0	1773	31	0
1	B	1747	0	1729	25	0
2	A	8	0	12	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	187	0	0	1	0
3	B	197	0	0	7	0
All	All	3911	0	3514	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:SER:OG	1:B:209[B]:MSE:HE2	1.34	1.25
1:A:107:SER:OG	1:A:209[B]:MSE:HE2	1.56	1.02
1:A:220[B]:ASN:HD21	1:A:223:LEU:HB2	1.35	0.92
1:A:220[B]:ASN:ND2	1:A:223:LEU:HB2	1.86	0.90
1:B:107:SER:OG	1:B:209[B]:MSE:CE	2.25	0.81
1:A:205:GLY:CA	1:A:209[B]:MSE:HE1	2.16	0.76
1:A:205:GLY:HA3	1:A:209[B]:MSE:HE1	1.71	0.73
1:A:133[B]:GLN:HG2	3:B:261[B]:HOH:O	1.88	0.72
1:B:114:LEU:CD1	1:B:135[B]:MSE:HE1	2.21	0.71
1:B:6[A]:THR:HG23	1:B:23:THR:HG23	1.74	0.69
1:B:128:VAL:HG12	1:B:129[A]:GLU:HG3	1.75	0.68
1:B:205:GLY:CA	1:B:209[B]:MSE:HE1	2.24	0.67
1:A:107:SER:OG	1:A:209[B]:MSE:CE	2.40	0.64
1:B:116[B]:LEU:HD11	1:B:122:LEU:HD11	1.83	0.60
1:B:206:ASP:H	1:B:209[B]:MSE:HE3	1.67	0.60
1:A:220[B]:ASN:ND2	1:A:223:LEU:CB	2.62	0.59
1:B:205:GLY:HA3	1:B:209[B]:MSE:HE1	1.85	0.57
1:A:133[B]:GLN:CG	3:B:261[B]:HOH:O	2.49	0.56
1:A:72[A]:ARG:HD2	1:B:140:LEU:HD22	1.87	0.56
1:A:103[B]:GLN:NE2	2:A:227:EDO:O1	2.38	0.56
1:A:220[B]:ASN:HD21	1:A:223:LEU:CB	2.12	0.55
1:B:6[B]:THR:HG22	1:B:23:THR:CG2	2.35	0.55
1:A:187:ILE:HD11	1:A:202[A]:CYS:SG	2.47	0.54
1:A:154:GLU:HG2	1:A:191:ASN:HD21	1.73	0.53
1:A:205:GLY:HA2	1:A:209[B]:MSE:HE1	1.90	0.53
1:A:140:LEU:HD22	1:B:72[A]:ARG:HD2	1.90	0.52
1:B:114:LEU:HD13	1:B:135[B]:MSE:HE1	1.90	0.52
1:B:205:GLY:HA2	1:B:209[B]:MSE:HE1	1.91	0.52
1:B:101:LEU:HD11	1:B:132:ALA:HB1	1.92	0.50
1:A:205:GLY:CA	1:A:209[B]:MSE:CE	2.88	0.50
1:A:206:ASP:O	1:A:209[B]:MSE:HG2	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:SER:CB	1:B:209[B]:MSE:HE2	2.37	0.49
1:A:128:VAL:HG12	1:A:129[A]:GLU:HG3	1.96	0.48
1:B:209[B]:MSE:HE3	3:B:340:HOH:O	2.14	0.47
1:B:130:ASN:ND2	3:B:358:HOH:O	2.47	0.47
1:A:205:GLY:HA3	1:A:209[B]:MSE:CE	2.41	0.47
1:A:154:GLU:CG	1:A:191:ASN:HD21	2.28	0.46
1:A:156:TYR:CE1	1:A:160:LYS:HG3	2.51	0.46
1:B:162[B]:ASN:OD1	1:B:166:GLN:NE2	2.48	0.46
2:A:227:EDO:C2	3:A:328:HOH:O	2.63	0.46
1:B:129[B]:GLU:HG2	3:B:277:HOH:O	2.16	0.46
1:A:206:ASP:O	1:A:209[B]:MSE:CG	2.63	0.46
1:A:224:ILE:HD12	1:A:224:ILE:N	2.31	0.46
1:B:72[A]:ARG:NH1	3:B:293:HOH:O	2.44	0.44
1:A:223:LEU:C	1:A:224:ILE:HD12	2.38	0.44
1:B:116[B]:LEU:CD1	1:B:122:LEU:HD11	2.48	0.44
1:A:154:GLU:OE2	1:A:191:ASN:ND2	2.51	0.44
1:A:103[B]:GLN:CD	2:A:227:EDO:O1	2.57	0.43
1:A:166:GLN:HE22	1:A:197:ASN:H	1.65	0.43
1:A:187:ILE:CD1	1:A:202[A]:CYS:SG	3.07	0.43
1:A:206:ASP:N	1:A:209[B]:MSE:HE3	2.34	0.41
1:B:206:ASP:O	1:B:209[B]:MSE:CG	2.69	0.41
1:A:101:LEU:HD11	1:A:132:ALA:HB1	2.01	0.41
1:B:220:ASN:HD22	1:B:223:LEU:HB2	1.86	0.40
1:B:209[B]:MSE:CE	3:B:340:HOH:O	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	238/226 (105%)	230 (97%)	8 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	234/226 (104%)	228 (97%)	6 (3%)	0	100	100
All	All	472/452 (104%)	458 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/165 (110%)	178 (98%)	4 (2%)	52	34
1	B	180/165 (109%)	180 (100%)	0	100	100
All	All	362/330 (110%)	358 (99%)	4 (1%)	78	63

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	146	LYS
1	A	202[A]	CYS
1	A	202[B]	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	170	GLN
1	A	191	ASN
1	B	76	ASN
1	B	130	ASN
1	B	220	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	226	-	3,3,3	0.52	0	2,2,2	0.33	0
2	EDO	A	227	-	3,3,3	0.31	0	2,2,2	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	226	-	-	1/1/1/1	-
2	EDO	A	227	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	226	EDO	O1-C1-C2-O2
2	A	227	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	227	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.