

wwPDB X-ray Structure Validation Summary Report (i)

Nov 19, 2023 – 04:57 PM JST

PDB ID : 6LPE

Title: Phascolosoma esculenta ferritin

Authors : Su, X.R.; Ming, T.H.

Deposited on : 2020-01-10

Resolution : 1.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	174	94%	5% •
1	В	174	93%	6% •
1	С	174	90%	9% •
1	D	174	91%	7% ••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-



ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	A	203	-	-	-	X
3	SO4	A	206	-	-	-	X
3	SO4	С	205	-	-	X	-
3	SO4	D	205	-	-	X	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6589 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ferritin.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	172	Total	С	N	О	S	0	0	0
1	A	172	1457	907	253	291	6	0	0	U
1	В	172	Total	С	N	О	S	0	12	0
1	Б	112	1488	928	259	293	8	0	12	0
1	С	172	Total	tal C N O S	0	11	0			
1		172	1479	921	256	295	7	0	11	U
1	1 D	172	Total	С	N	О	S	0	0	0
	112	1462	912	253	290	7	0	9	U	

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	4	Total Mg 4 4	0	0
2	В	3	Total Mg 3 3	0	0
2	С	2	Total Mg 2 2	0	0
2	D	3	Total Mg 3 3	0	0

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Ato	ms		ZeroOcc	AltConf	
3	A	1	Total	О	S	0	0	
3	A	1	5	4	1	0		
3	A	1	Total	О	S	0	0	
	11	1	5	4	1	0	U	
3	A	1	Total	Ο	S	0	0	
	11	1	5	4	1	Ü	Ü	
3	A	1	Total	O	S	0	0	
		_	5	4	1	Ü	0	
3	A	1	Total	O	S	0	0	
		_	5	4	1		Ū	
3	В	1	Total	O	S	0	0	
			5	4	1			
3	В	1	Total	O	S	0	0	
			5	4	1			
3	В	1	Total	O	S	0	0	
			5	4	1			
3	С	1	Total	O	S 1	0	0	
			5	4 O	S			
3	С	1	Total	4	5 1	0	0	
			5 Total	O	$\frac{1}{S}$			
3	С	1	5	4	1	0	0	
			Total	O	S			
3	С	1	5	4	1	0	0	
			Total	O	S			
3	D	1	5	4	1	0	0	
	_		Total	O	S		_	
3	D	1	5	4	1	0	0	
						1: 7		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total O S 5 4 1	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	158	Total O 158 158	0	0
4	В	158	Total O 158 158	0	0
4	С	152	Total O 152 152	0	0
4	D	148	Total O 148 148	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants	232.76Å 232.76Å 232.76Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.18 - 1.99	Depositor
Resolution (A)	41.15 - 1.99	EDS
% Data completeness	99.5 (41.18-1.99)	Depositor
(in resolution range)	99.5 (41.15-1.99)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.07 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
D D.	0.249 , 0.302	Depositor
R, R_{free}	0.258 , 0.305	DCC
R_{free} test set	3603 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 44.5	EDS
L-test for twinning ²	$ < L > = 0.44, < L^2> = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	6589	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 96.26 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.4820e-10. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.85	0/1486	0.85	1/2003 (0.0%)	
1	В	0.84	0/1517	0.85	0/2044	
1	С	0.83	0/1508	0.83	1/2032 (0.0%)	
1	D	0.84	$2/1491 \ (0.1\%)$	0.82	0/2009	
All	All	0.84	$2/6002 \ (0.0\%)$	0.84	2/8088 (0.0%)	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	D	168	GLU	CD-OE2	-5.79	1.19	1.25
1	D	120	GLU	CD-OE1	-5.01	1.20	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	87	ASP	CB-CG-OD1	-5.61	113.25	118.30
1	С	87	ASP	CB-CG-OD1	-5.08	113.73	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1457	0	1321	5	0
1	В	1488	0	1362	10	2
1	С	1479	0	1344	15	1
1	D	1462	0	1333	10	1
2	A	4	0	0	0	0
2	В	3	0	0	0	0
2	С	2	0	0	0	0
2	D	3	0	0	1	0
3	A	25	0	0	1	0
3	В	15	0	0	0	0
3	С	20	0	0	2	2
3	D	15	0	0	1	1
4	A	158	0	0	1	0
4	В	158	0	0	0	0
4	С	152	0	0	2	1
4	D	148	0	0	3	0
All	All	6589	0	5360	40	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:A:168[A]:GLU:OE2	3:A:206:SO4:O3	1.63	1.14
2:D:203:MG:MG	4:D:301:HOH:O	1.04	1.00
1:C:139[B]:GLN:O	1:C:143[B]:ILE:HG22	1.63	0.98
1:B:172:GLY:O	1:B:173:ASP:HB2	1.73	0.89
1:D:139:GLN:O	1:D:143[B]:ILE:HG22	1.72	0.89

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)
1:B:168:GLU:OE2	3:C:205:SO4:O3[38_555]	1.61	0.59
1:C:168:GLU:OE2	3:C:205:SO4:O1[2_655]	1.86	0.34
1:D:168:GLU:OE2	3:D:205:SO4:O3[17_555]	1.91	0.29
1:B:173:ASP:OD2	4:C:411:HOH:O[37_545]	2.09	0.11



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	178/174 (102%)	174 (98%)	4 (2%)	0	100	100
1	В	182/174 (105%)	180 (99%)	2 (1%)	0	100	100
1	С	181/174 (104%)	179 (99%)	2 (1%)	0	100	100
1	D	179/174 (103%)	176 (98%)	3 (2%)	0	100	100
All	All	720/696 (103%)	709 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	A	148/153 (97%)	145 (98%)	3 (2%)	55	58
1	В	152/153 (99%)	149 (98%)	3 (2%)	55	58
1	С	151/153 (99%)	150 (99%)	1 (1%)	84	88
1	D	149/153 (97%)	146 (98%)	3 (2%)	55	58
All	All	600/612 (98%)	590 (98%)	10 (2%)	69	65

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	30	TYR
1	D	138[A]	GLU
1	D	138[B]	GLU

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	В	30	TYR
1	В	138[A]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	HIS
1	В	128	GLN
1	С	134	HIS
1	D	134	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 27 ligands modelled in this entry, 12 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain R	Res L	Res Link	Bond lengths			Bond angles			
WIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	SO4	В	205	-	4,4,4	0.24	0	6,6,6	0.10	0
3	SO4	С	204	-	4,4,4	0.33	0	6,6,6	0.05	0
3	SO4	С	205	-	4,4,4	0.40	0	6,6,6	0.32	0



Mal	Trmo	Chain	Res	Link	В	Bond lengths			Bond angles		
Mol	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
3	SO4	С	203	-	4,4,4	0.20	0	6,6,6	0.22	0	
3	SO4	A	208	-	4,4,4	0.28	0	6,6,6	0.09	0	
3	SO4	D	205	-	4,4,4	0.31	0	6,6,6	0.49	0	
3	SO4	A	205	-	4,4,4	0.45	0	6,6,6	0.24	0	
3	SO4	В	204	-	4,4,4	0.18	0	6,6,6	0.17	0	
3	SO4	D	206	-	4,4,4	0.19	0	6,6,6	0.13	0	
3	SO4	A	206	-	4,4,4	0.40	0	6,6,6	0.27	0	
3	SO4	С	206	-	4,4,4	0.30	0	6,6,6	0.20	0	
3	SO4	D	204	-	4,4,4	0.19	0	6,6,6	0.17	0	
3	SO4	A	209	-	4,4,4	0.45	0	6,6,6	0.16	0	
3	SO4	A	207	-	4,4,4	0.40	0	6,6,6	0.25	0	
3	SO4	В	206	-	4,4,4	0.30	0	6,6,6	0.17	0	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	С	205	SO4	2	2
3	D	205	SO4	1	1
3	A	206	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.










































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	172/174 (98%)	-0.42	3 (1%) 70 68	32, 36, 44, 96	1 (0%)
1	В	172/174 (98%)	-0.42	1 (0%) 89 88	32, 36, 44, 96	2 (1%)
1	С	172/174 (98%)	-0.42	1 (0%) 89 88	32, 35, 44, 96	2 (1%)
1	D	172/174 (98%)	-0.43	1 (0%) 89 88	32, 36, 44, 94	2 (1%)
All	All	688/696 (98%)	-0.42	6 (0%) 84 83	32, 36, 44, 96	7 (1%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	173	ASP	3.2
1	С	173	ASP	3.1
1	В	173	ASP	3.1
1	D	173	ASP	2.8
1	A	172	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
2	MG	A	203	1/1	-0.25	0.53	75,75,75,75	1
3	SO4	A	206	5/5	0.20	0.41	76,77,103,114	5
2	MG	В	203	1/1	0.28	0.27	64,64,64,64	1
3	SO4	С	205	5/5	0.54	0.20	49,56,101,143	5
2	MG	D	203	1/1	0.65	0.25	86,86,86,86	0
2	MG	A	204	1/1	0.80	0.25	85,85,85,85	1
2	MG	D	201	1/1	0.84	0.12	47,47,47,47	1
2	MG	A	201	1/1	0.84	0.10	57,57,57,57	0
3	SO4	D	205	5/5	0.84	0.09	121,167,185,190	5
3	SO4	В	205	5/5	0.85	0.29	83,104,108,113	5
3	SO4	A	208	5/5	0.86	0.39	89,101,114,116	5
3	SO4	D	206	5/5	0.89	0.40	77,96,113,116	5
3	SO4	A	207	5/5	0.94	0.24	40,55,81,90	5
3	SO4	С	204	5/5	0.95	0.09	36,59,61,75	5
2	MG	С	201	1/1	0.96	0.14	56,56,56,56	0
3	SO4	В	206	5/5	0.97	0.12	65,67,71,76	5
2	MG	D	202	1/1	0.97	0.10	31,31,31,31	0
2	MG	В	202	1/1	0.97	0.10	31,31,31,31	0
3	SO4	С	206	5/5	0.97	0.11	64,65,70,71	5
3	SO4	A	209	5/5	0.97	0.13	61,62,65,72	5
2	MG	В	201	1/1	0.97	0.10	48,48,48,48	1
2	MG	С	202	1/1	0.98	0.11	31,31,31,31	0
3	SO4	В	204	5/5	0.98	0.07	50,54,57,62	1
2	MG	A	202	1/1	0.98	0.10	31,31,31,31	0
3	SO4	D	204	5/5	0.98	0.06	49,54,55,56	1
3	SO4	A	205	5/5	0.98	0.11	39,45,46,54	5
3	SO4	С	203	5/5	0.98	0.06	50,55,58,60	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around SO4 A 206:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around MG B 203: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around MG A 204: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around MG D 201: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c \ (\mathrm{at}\ 0.7\ \mathrm{rmsd}) \ \mathrm{in}\ \mathrm{gray}$

 ${
m mF}_o{
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around MG A 201:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)















Electron density around SO4 B 205: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around SO4 A 207:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around SO4 C 204: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around MG C 201: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 B 206: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around MG D 202: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around MG B 202: $2mF_o$ -DF $_c$ (at 0.7 rmsd) in gray mF_o -DF $_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SO4 C 206: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)








Electron density around MG B 201: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SO4 B 204:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around MG A 202: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 D 204: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

