



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 11, 2024 – 04:00 PM EDT

PDB ID : 2M5K
EMDB ID : EMD-5590
BMRB ID : 19058
Title : Atomic-resolution structure of a doublet cross-beta amyloid fibril
Authors : Fitzpatrick, A.W.P.; Debelouchina, G.T.; Bayro, M.J.; Clare, D.K.; Caporini, M.A.; Bajaj, V.S.; Jaroniec, C.P.; Wang, L.; Ladizhansky, V.; Muller, S.; MacPhee, C.E.; Waudby, C.A.; Mott, H.R.; de Simone, A.; Knowles, T.P.J.; Saibil, H.R.; Vendruscolo, M.; Orlova, E.V.; Griffin, R.G.; Dobson, C.M.
Deposited on : 2013-02-27

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **NOT EXECUTED**
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : **NOT EXECUTED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.1

1 Overall quality at a glance


The following experimental techniques were used to determine the structure:

SOLID-STATE NMR, ELECTRON MICROSCOPY

The reported resolution of this entry is 12.70 Å.

The overall completeness of chemical shifts assignment is 5%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	11	55% 36% 9%
1	B	11	55% 36% 9%
1	C	11	55% 36% 9%
1	D	11	55% 36% 9%
1	E	11	55% 36% 9%
1	F	11	55% 36% 9%
1	G	11	55% 36% 9%
1	H	11	55% 36% 9%

2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1376 atoms, of which 696 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transthyretin.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	H	N	O		
1	A	11	172	57	87	11	17	0	
1	B	11	172	57	87	11	17	0	
1	C	11	172	57	87	11	17	0	
1	D	11	172	57	87	11	17	0	
1	E	11	172	57	87	11	17	0	
1	F	11	172	57	87	11	17	0	
1	G	11	172	57	87	11	17	0	
1	H	11	172	57	87	11	17	0	

4 Residue-property plots [i](#)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Transthyretin

Chain A:  55% 36% 9%




- Molecule 1: Transthyretin

Chain B:  55% 36% 9%



- Molecule 1: Transthyretin

Chain C:  55% 36% 9%




- Molecule 1: Transthyretin

Chain D:  55% 36% 9%



- Molecule 1: Transthyretin

Chain E:  55% 36% 9%




- Molecule 1: Transthyretin

Chain F:  55% 36% 9%



- Molecule 1: Transthyretin



- Molecule 1: Transthyretin



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 1 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNSSOLVE	structure solution	
CNSSOLVE	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	65
Number of shifts mapped to atoms	65
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	5%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	85	87	87	3
1	B	85	87	87	3
1	C	85	87	87	3
1	D	85	87	87	3
1	E	85	87	87	3
1	F	85	87	87	3
1	G	85	87	87	3
1	H	85	87	87	3
All	All	680	696	696	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

5 of 12 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:C:7:LEU:HD21	1:D:6:LEU:HD23	0.54	1.79
1:G:7:LEU:HD21	1:H:6:LEU:HD23	0.53	1.79
1:A:7:LEU:HD21	1:B:6:LEU:HD23	0.53	1.80
1:E:7:LEU:HD21	1:F:6:LEU:CD2	0.53	2.34
1:A:7:LEU:HD21	1:B:6:LEU:CD2	0.52	2.34

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
1	B	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
1	C	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
1	D	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
1	E	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
1	F	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
1	G	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
1	H	9/11 (82%)	9 (100%)	0 (0%)	0 (0%)	100	100
All	All	72/88 (82%)	72 (100%)	0 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	9/9 (100%)	5 (56%)	4 (44%)	0	2
1	B	9/9 (100%)	5 (56%)	4 (44%)	0	2
1	C	9/9 (100%)	5 (56%)	4 (44%)	0	2
1	D	9/9 (100%)	5 (56%)	4 (44%)	0	2
1	E	9/9 (100%)	5 (56%)	4 (44%)	0	2
1	F	9/9 (100%)	5 (56%)	4 (44%)	0	2
1	G	9/9 (100%)	5 (56%)	4 (44%)	0	2
1	H	9/9 (100%)	5 (56%)	4 (44%)	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	72/72 (100%)	40 (56%)	32 (44%)	0 2

5 of 32 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	2	THR
1	A	6	LEU
1	A	7	LEU
1	A	11	SER
1	B	2	THR

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 5% for the well-defined parts and 5% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	65
Number of shifts mapped to atoms	65
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 5%, i.e. 62 atoms were assigned a chemical shift out of a possible 1160. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	30/424 (7%)	0/168 (0%)	22/176 (12%)	8/80 (10%)
Sidechain	22/592 (4%)	0/408 (0%)	22/184 (12%)	0/0 (—%)
Aromatic	10/144 (7%)	0/64 (0%)	10/80 (12%)	0/0 (—%)
Overall	62/1160 (5%)	0/640 (0%)	54/440 (12%)	8/80 (10%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1408
Intra-residue ($ i-j =0$)	384
Sequential ($ i-j =1$)	672
Medium range ($ i-j >1$ and $ i-j <5$)	160
Long range ($ i-j \geq 5$)	96
Inter-chain	96
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	144
Number of unmapped restraints	0
Number of restraints per residue	17.6
Number of long range restraints per residue ¹	1.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	32.0	0.13
0.2-0.5 (Medium)	None	None
>0.5 (Large)	288.0	11.44

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	28.0	1.88
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None

9 Distance violation analysis


9.1 Summary of distance violations

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	384	27.3	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	384	27.3	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	672	47.7	32	4.8	2.3	32	4.8	2.3
Backbone-Backbone	64	4.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	608	43.2	32	5.3	2.3	32	5.3	2.3
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Medium range (i-j >1 & i-j <5)	160	11.4	96	60.0	6.8	96	60.0	6.8
Backbone-Backbone	64	4.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	96	6.8	96	100.0	6.8	96	100.0	6.8
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range (i-j ≥5)	96	6.8	96	100.0	6.8	96	100.0	6.8
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	32	2.3	32	100.0	2.3	32	100.0	2.3
Sidechain-Sidechain	64	4.5	64	100.0	4.5	64	100.0	4.5
Inter-chain	96	6.8	96	100.0	6.8	96	100.0	6.8
Backbone-Backbone	80	5.7	80	100.0	5.7	80	100.0	5.7
Backbone-Sidechain	16	1.1	16	100.0	1.1	16	100.0	1.1
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1408	100.0	320	22.7	22.7	320	22.7	22.7
Backbone-Backbone	208	14.8	80	38.5	5.7	80	38.5	5.7
Backbone-Sidechain	1136	80.7	176	15.5	12.5	176	15.5	12.5
Sidechain-Sidechain	64	4.5	64	100.0	4.5	64	100.0	4.5

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis


9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	0	32	96	96	96	320	6.17	11.44	3.3	7.16

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right


9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1088(IR:384, SQ:640, MR:64, LR:0, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	32	96	96	96	320	1	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble [\(i\)](#)




9.4 Most violated distance restraints in the ensemble [\(i\)](#)

No violations found

9.5 All violated distance restraints [\(i\)](#)

9.5.1 Histogram : Distribution of distance violations [\(i\)](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1728)	1:3:H:ILE:CD1	1:9:H:PRO:CB	1	11.44
(1,1684)	1:3:F:ILE:CD1	1:9:F:PRO:CB	1	11.44
(1,1638)	1:3:D:ILE:CD1	1:9:D:PRO:CB	1	11.44
(1,1546)	1:3:H:ILE:CD1	1:9:H:PRO:CB	1	11.44
(1,1502)	1:3:F:ILE:CD1	1:9:F:PRO:CB	1	11.44
(1,1456)	1:3:D:ILE:CD1	1:9:D:PRO:CB	1	11.44
(1,1592)	1:3:B:ILE:CD1	1:9:B:PRO:CB	1	11.43
(1,1410)	1:3:B:ILE:CD1	1:9:B:PRO:CB	1	11.43
(1,1707)	1:3:G:ILE:CD1	1:9:G:PRO:CB	1	11.36
(1,1615)	1:3:C:ILE:CD1	1:9:C:PRO:CB	1	11.36

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
PSI	72	50.0	13	18.1	9.0	13	18.1	9.0
PHI	72	50.0	15	20.8	10.4	15	20.8	10.4
Total	144	100.0	28	19.4	19.4	28	19.4	19.4

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model [i](#)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations			Mean (°)	Max (°)	SD (°)	Median (°)
	PSI	PHI	Total				
1	13	15	28	1.42	1.88	0.27	1.36

10.2.1 Bar graph : Dihedral violation statistics for each model [i](#)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right


10.3 Dihedral-angle violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints			Fraction of the ensemble	
PSI	PHI	Total	Count ¹	%
13	15	28	1	100.0

¹ Number of models with violations

10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble [i](#)




10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

No violations found

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,85)	1:4:E:ALA:C	1:5:E:ALA:N	1:5:E:ALA:CA	1:5:E:ALA:C	1	1.88
(1,4)	1:4:A:ALA:N	1:4:A:ALA:CA	1:4:A:ALA:C	1:5:A:ALA:N	1	1.86
(1,76)	1:4:E:ALA:N	1:4:E:ALA:CA	1:4:E:ALA:C	1:5:E:ALA:N	1	1.85
(1,49)	1:4:C:ALA:C	1:5:C:ALA:N	1:5:C:ALA:CA	1:5:C:ALA:C	1	1.85
(1,121)	1:4:G:ALA:C	1:5:G:ALA:N	1:5:G:ALA:CA	1:5:G:ALA:C	1	1.76
(1,112)	1:4:G:ALA:N	1:4:G:ALA:CA	1:4:G:ALA:C	1:5:G:ALA:N	1	1.75
(1,13)	1:4:A:ALA:C	1:5:A:ALA:N	1:5:A:ALA:CA	1:5:A:ALA:C	1	1.74
(1,40)	1:4:C:ALA:N	1:4:C:ALA:CA	1:4:C:ALA:C	1:5:C:ALA:N	1	1.6
(1,130)	1:4:H:ALA:N	1:4:H:ALA:CA	1:4:H:ALA:C	1:5:H:ALA:N	1	1.56
(1,66)	1:3:D:ILE:C	1:4:D:ALA:N	1:4:D:ALA:CA	1:4:D:ALA:C	1	1.56