

wwPDB X-ray Structure Validation Summary Report (i)

Jun 15, 2020 – 10:19 pm BST

PDB ID : 1MCK

Title: PRINCIPLES AND PITFALLS IN DESIGNING SITE DIRECTED PEP-

TIDE LIGANDS

Authors: Edmundson, A.B.; Harris, D.L.; Fan, Z.-C.; Guddat, L.W.

Deposited on : 1993-02-25

Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	A	216	39%	44%	17%	-	
1	В	216	44%	39%	15%	•	
2	Р	5	60%	20%	20%		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called IMMUNOGLOBULIN LAMBDA DIMER MCG (LIGHT CHAIN).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	216	Total	С	N	О	S	0	0	0
1	A	210	1605	1000	266	334	5	U	0	0
1	D	216	Total	С	N	О	S	0	0	0
1	D	210	1605	1000	266	334	5	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	ILE	PHE	CONFLICT	PIR S14675
A	23	THR	SER	CONFLICT	PIR S14675
A	29	VAL	ILE	CONFLICT	PIR S14675
A	31	GLY	ASN	CONFLICT	PIR S14675
A	39	GLN	ARG	CONFLICT	PIR S14675
A	42	ALA	PRO	CONFLICT	PIR S14675
A	48	VAL	LEU	CONFLICT	PIR S14675
A	49	ILE	MET	CONFLICT	PIR S14675
A	54	ASN	THR	CONFLICT	PIR S14675
A	62	ASP	ASN	CONFLICT	PIR S14675
A	94	GLU	ALA	CONFLICT	PIR S14675
A	97	ASP	ASN	CONFLICT	PIR S14675
A	98	ASN	SER	CONFLICT	PIR S14675
A	99	PHE	LEU	CONFLICT	PIR S14675
A	100	VAL	ILE	CONFLICT	PIR S14675
A	103	THR	GLY	CONFLICT	PIR S14675
A	106	LYS	ARG	CONFLICT	PIR S14675
A	107	VAL	LEU	CONFLICT	PIR S14675
A	116	ASN	ALA	CONFLICT	PIR S14675
A	118	THR	SER	CONFLICT	PIR S14675
A	156	GLY	SER	CONFLICT	PIR S14675
A	167	LYS	THR	CONFLICT	PIR S14675
В	20	ILE	PHE	CONFLICT	PIR S14675
В	23	THR	SER	CONFLICT	PIR S14675

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	29	VAL	ILE	CONFLICT	PIR S14675
В	31	GLY	ASN	CONFLICT	PIR S14675
В	39	GLN	ARG	CONFLICT	PIR S14675
В	42	ALA	PRO	CONFLICT	PIR S14675
В	48	VAL	LEU	CONFLICT	PIR S14675
В	49	ILE	MET	CONFLICT	PIR S14675
В	54	ASN	THR	CONFLICT	PIR S14675
В	62	ASP	ASN	CONFLICT	PIR S14675
В	94	GLU	ALA	CONFLICT	PIR S14675
В	97	ASP	ASN	CONFLICT	PIR S14675
В	98	ASN	SER	CONFLICT	PIR S14675
В	99	PHE	LEU	CONFLICT	PIR S14675
В	100	VAL	ILE	CONFLICT	PIR S14675
В	103	THR	GLY	CONFLICT	PIR S14675
В	106	LYS	ARG	CONFLICT	PIR S14675
В	107	VAL	LEU	CONFLICT	PIR S14675
В	116	ASN	ALA	CONFLICT	PIR S14675
В	118	THR	SER	CONFLICT	PIR S14675
В	156	GLY	SER	CONFLICT	PIR S14675
В	167	LYS	THR	CONFLICT	PIR S14675

 \bullet Molecule 2 is a protein called PEPTIDE N-ACETYL-D-GLU-L-HIS-D-PRO-NH2.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
9	D	5	Total	С	N	О	0	0	1
	1	9	30	18	6	6	0	U	1

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0
3	В	1	Total O 1 1	0	0
3	Р	2	Total O 2 2	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: IMMUNOGLOBULIN LAMBDA DIMER MCG (LIGHT CHAIN)





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants	72.30Å 72.30Å 185.90Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 - 2.70	Depositor
% Data completeness	(Not available) (6.00-2.70)	Depositor
(in resolution range)	, , , , , , , , , , , , , , , , , , , ,	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.189 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3244	wwPDB-VP
Average B, all atoms (Å ²)	0.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2, DGL, ACE, DPR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.86	0/1644	1.42	$11/2241 \ (0.5\%)$	
1	В	0.86	0/1644	1.38	$12/2241 \ (0.5\%)$	
2	Р	0.51	0/10	1.46	0/12	
All	All	0.86	0/3298	1.40	$23/4494 \ (0.5\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	Р	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	A	56	ARG	NE-CZ-NH1	7.95	124.28	120.30
1	В	34	TYR	CA-CB-CG	7.81	128.25	113.40
1	A	202	GLU	CA-CB-CG	7.15	129.12	113.40
1	В	52	GLU	CA-CB-CG	6.76	128.26	113.40
1	A	193	ARG	NE-CZ-NH1	6.42	123.51	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	167	LYS	Peptide
2	Р	2	HIS	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1605	0	1540	87	0
1	В	1605	0	1540	95	0
2	Р	30	0	22	3	0
3	A	1	0	0	1	0
3	В	1	0	0	0	0
3	Р	2	0	0	0	0
All	All	3244	0	3102	177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance } (ext{Å}) \end{array}$	Clash overlap (Å)	
1:B:158:PRO:HB2	1:B:160:LYS:HD2	1.56	0.86	
1:B:187:GLU:HA	1:B:190:LYS:HD2	1.57	0.86	
1:B:38:TYR:HE1	1:B:91:SER:HB2	1.39	0.86	
1:B:70:GLY:O	1:B:72:THR:N	2.09	0.86	
1:A:37:TRP:HB2	1:A:50:ILE:HB	1.61	0.82	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Favoured	Allowed	Outliers	Percent	iles
1	A	214/216 (99%)	174 (81%)	26 (12%)	14 (6%)	1 2	?
1	В	214/216 (99%)	176 (82%)	28 (13%)	10 (5%)	2 4	:
2	Р	1/5 (20%)	1 (100%)	0	0	100 1	.00
All	All	429/437 (98%)	351 (82%)	54 (13%)	24 (6%)	2 3	

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	ALA
1	A	16	GLN
1	A	62	ASP
1	A	80	LEU
1	A	82	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	181/181 (100%)	140 (77%)	41 (23%)	1 2
1	В	181/181 (100%)	139 (77%)	42 (23%)	1 2
2	Р	1/1 (100%)	0	1 (100%)	0 0
All	All	363/363 (100%)	279 (77%)	84 (23%)	1 2

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	ASN
1	В	48	VAL
1	В	183	SER
1	A	183	SER
1	В	14	LEU



Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	${f Res}$	\mathbf{Type}
1	В	16	GLN
1	В	174	ASN
1	В	39	GLN
1	В	6	GLN
1	В	130	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

