



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 02:31 am BST

PDB ID : 1MLB  
Title : MONOCLONAL ANTIBODY FAB D44.1 RAISED AGAINST CHICKEN EGG-WHITE LYSOZYME  
Authors : Braden, B.C.; Souchon, H.; Eisele, J.-L.; Bentley, G.A.; Bhat, T.N.; Navaza, J.; Poljak, R.J.  
Deposited on : 1995-03-08  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	214	
2	B	218	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG1-KAPPA D44.1 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	214	1670	1029	289	345	7	0	2	0

- Molecule 2 is a protein called IGG1-KAPPA D44.1 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	218	1639	1033	269	329	8	0	2	0

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	5	GLN	VAL	CONFLICT	PIR PC4202
B	11	VAL	LEU	CONFLICT	PIR PC4202
B	26	GLY	ASP	CONFLICT	PIR PC4202
B	28	THR	ARG	CONFLICT	PIR PC4202
B	31	THR	SER	CONFLICT	PIR PC4202
B	50	GLU	ASP	CONFLICT	PIR PC4202
B	57	SER	ASN	CONFLICT	PIR PC4202
B	59	TYR	ASN	CONFLICT	PIR PC4202
B	63	LYS	ARG	CONFLICT	PIR PC4202
B	98	ARG	ILE	CONFLICT	PIR PC4202
B	99	GLY	PRO	CONFLICT	PIR PC4202
B	101	GLY	-	INSERTION	PIR PC4202
B	102	ASN	-	INSERTION	PIR PC4202
B	103	TYR	-	INSERTION	PIR PC4202
B	104	GLY	-	INSERTION	PIR PC4202
B	118	SER	LYS	CONFLICT	PIR PC4202
B	125	PHE	TYR	CONFLICT	PIR PC4202

- Molecule 3 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	71	Total 71	O 71	0	0
3	B	44	Total 44	O 44	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: IGG1-KAPPA D44.1 FAB (LIGHT CHAIN)



- Molecule 2: IGG1-KAPPA D44.1 FAB (HEAVY CHAIN)



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.70Å 136.20Å 43.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) (7.00-2.10)	Depositor
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.181 , 0.274	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3424	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.00	1/1717 (0.1%)	1.99	39/2326 (1.7%)
2	B	1.00	0/1695	1.76	27/2315 (1.2%)
All	All	1.00	1/3412 (0.0%)	1.88	66/4641 (1.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	GLU	CD-OE2	-5.43	1.19	1.25

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	ARG	NE-CZ-NH1	23.13	131.87	120.30
1	A	211	ARG	NE-CZ-NH2	-17.58	111.51	120.30
1	A	211	ARG	NE-CZ-NH1	16.38	128.49	120.30
1	A	108	ARG	NE-CZ-NH2	-12.82	113.89	120.30
1	A	188[A]	ARG	NE-CZ-NH2	-10.73	114.93	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1670	0	1590	70	0
2	B	1639	0	1582	100	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	71	0	0	0	0
3	B	44	0	0	1	0
All	All	3424	0	3172	161	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 161 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188[B]:ARG:CZ	1:A:188[B]:ARG:NE	1.82	1.42
1:A:24[B]:ARG:CG	1:A:24[B]:ARG:CD	2.02	1.38
2:B:51:ILE:HG13	2:B:58:THR:HG22	1.45	0.97
2:B:187:PRO:O	2:B:190:PRO:HD2	1.73	0.89
2:B:48:ILE:HG21	2:B:81[A]:MET:CE	2.02	0.89

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/214 (100%)	201 (94%)	10 (5%)	3 (1%)	11	6
2	B	218/218 (100%)	201 (92%)	12 (6%)	5 (2%)	6	2
All	All	432/432 (100%)	402 (93%)	22 (5%)	8 (2%)	8	3

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	SER
1	A	110	ASP

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type
2	B	190	PRO
2	B	42	GLY
1	A	77	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	196/194 (101%)	175 (89%)	21 (11%)	<b>6</b> <b>3</b>
2	B	186/184 (101%)	159 (86%)	27 (14%)	<b>3</b> <b>1</b>
All	All	382/378 (101%)	334 (87%)	48 (13%)	<b>4</b> <b>2</b>

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	3	GLN
2	B	57	SER
2	B	205	SER
2	B	4	LEU
2	B	30	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	157	ASN
1	A	161	ASN
2	B	3	GLN
1	A	156	GLN
1	A	198	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.