

# wwPDB NMR Structure Validation Summary Report (i)

#### Jun 3, 2023 – 06:20 PM EDT

PDB ID	:	2MMV
BMRB ID	:	19869
Title	:	ZapA mutant dimer from Geobacillus stearothermophilus
Authors	:	Nogueira, M.L.; Sforca, M.; Zeri, A.
Deposited on	:	2014-03-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment is 39%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	$\mathbf{NMR}$ archive $(\#\mathbf{Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	А	86	79%	5%	16%		
1	В	86	74%	9%	16%		



# 2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core	Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:10-A:81, B:10-B:81 (144)	0.68	11			

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 14, 15, 16, 18, 19, 20
2	6, 13, 17



# 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2823 atoms, of which 1434 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Cell division protein ZapA.

Mol	Chain	Residues	Atoms					Trace	
1	٨	86	Total	С	Н	Ν	0	S	0
	I A	80	1412	441	717	121	130	3	0
1	В	96	Total	С	Н	Ν	0	S	0
	D	86	1411	441	717	121	129	3	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	1	GLY	-	expression tag	UNP A0A078N0N2
А	2	SER	-	expression tag	UNP A0A078N0N2
А	3	HIS	-	expression tag	UNP A0A078N0N2
А	50	GLN	ASN	engineered mutation	UNP A0A078N0N2
A	65	GLN	ASN	engineered mutation	UNP A0A078N0N2
А	78	GLN	ASN	engineered mutation	UNP A0A078N0N2
В	1	GLY	-	expression tag	UNP A0A078N0N2
В	2	SER	-	expression tag	UNP A0A078N0N2
В	3	HIS	-	expression tag	UNP A0A078N0N2
В	50	GLN	ASN	engineered mutation	UNP A0A078N0N2
В	65	GLN	ASN	engineered mutation	UNP A0A078N0N2
В	78	GLN	ASN	engineered mutation	UNP A0A078N0N2




# 4 Residue-property plots (i)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Cell division protein ZapA



# 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 11. Colouring as in section 4.1 above.

• Molecule 1: Cell division protein ZapA

Chain A:	77%	7%	16%
G 1 82 82 84 84 86 7 88 8 88 8 88 8 82 8 82 8 82 8 82 8	L80 R81 E82 E83 K85 K85 K85		
• Molecule 1: Cell division	protein ZapA		
Chain B:	65%	19%	16%
61 83 84 85 85 86 88 88 18 83 18 18 18 18 18 18 18 18 18 18 18 18 18	844 844 844 844 844 846 844 846 846 846	E82 Q8 3 K86 K86 K86	



# 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: torsion angle dynamics, SIMULATED AN-NEALING.

Of the ? calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	2.1
CYANA	structure solution	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	898
Number of shifts mapped to atoms	898
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	39%



# 6 Model quality (i)

### 6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		E	Sond lengths	Bond angles		
	Ullalli	RMSZ	$\#Z{>}5$	RMSZ	#Z > 5	
1	А	$0.28 {\pm} 0.02$	$0{\pm}0/592~(~0.0{\pm}~0.0\%)$	$0.37 {\pm} 0.02$	$0{\pm}0/801~(~0.0{\pm}~0.0\%)$	
1	В	$0.27 {\pm} 0.00$	$0{\pm}0/592~(~0.0{\pm}~0.0\%)$	$0.37 {\pm} 0.01$	$0{\pm}0/801~(~0.0{\pm}~0.0\%)$	
All	All	0.27	1/23680 ( $0.0%$ )	0.37	1/32040 ( $0.0%$ )	

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$	Moo Worst	I
1	А	48	GLU	CD-OE2	-5.24	1.19	1.25	2	1

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$	<b>Moc</b> Worst	<b>lels</b> Total
1	А	54	ASP	CB-CG-OD2	5.19	122.97	118.30	3	1

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	581	598	596	$3\pm 2$
1	В	581	598	596	3±1
All	All	23240	23920	23840	115

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 2.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:12:VAL:HG11	1:A:32:ILE:HD12	0.59	1.74	20	4
1:B:41:ASP:HA	1:B:44:HIS:ND1	0.57	2.15	14	1
1:B:12:VAL:HG11	1:B:32:ILE:HD12	0.57	1.76	20	3
1:A:31:HIS:O	1:A:35:VAL:HG23	0.56	2.01	12	6
1:A:42:LYS:O	1:A:45:GLU:HG2	0.56	2.01	7	8

5 of 64 unique clashes are listed below, sorted by their clash magnitude.

### 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	72/86~(84%)	$68 \pm 1 (94 \pm 2\%)$	$4\pm1~(5\pm2\%)$	$1\pm1 (1\pm1\%)$	24 71
1	В	72/86~(84%)	$67 \pm 1 (94 \pm 2\%)$	$4\pm1~(5\pm2\%)$	$1\pm1 (1\pm1\%)$	20 68
All	All	2880/3440~(84%)	2702 (94%)	152 (5%)	26 (1%)	21 69

5 of 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	В	51	PRO	10
1	А	51	PRO	6
1	А	27	GLU	3
1	В	27	GLU	3
1	А	55	VAL	2

#### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	63/76~(83%)	$55\pm2$ (87 $\pm4\%$ )	$8\pm2~(13\pm4\%)$	7 48
1	В	63/76~(83%)	$54\pm2$ (85 $\pm3\%$ )	$9\pm2~(15\pm3\%)$	6 44
All	All	2520/3040~(83%)	2166~(86%)	354 (14%)	6 46

5 of 82 unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	В	80	LEU	13
1	В	23	ILE	12
1	В	42	LYS	12
1	А	17	TYR	12
1	В	17	TYR	11

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.6 Ligand geometry (i)

There are no ligands in this entry.

#### 6.7 Other polymers (i)

There are no such molecules in this entry.

#### 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 39% for the well-defined parts and 37% for the entire structure.

### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: assigned\_chem\_shift\_list\_1

#### 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	898
Number of shifts mapped to atoms	898
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

#### 7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction}\pm{\rm precision},ppm$	Suggested action
$^{13}C_{\alpha}$	79	$-0.40 \pm 0.08$	None needed ( $< 0.5$ ppm)
$^{13}C_{\beta}$	75	$0.15 \pm 0.09$	None needed ( $< 0.5$ ppm)
$^{13}C'$	0		None (insufficient data)
$^{15}N$	74	$0.13 \pm 0.17$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 39%, i.e. 806 atoms were assigned a chemical shift out of a possible 2066. 0 out of 30 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	284/712~(40%)	143/286~(50%)	72/288~(25%)	69/138~(50%)
Sidechain	489/1210~(40%)	335/790~(42%)	154/372~(41%)	0/48~(0%)

Continued on next page...



$\alpha$ $\cdot$ $\cdot$ $\cdot$	C	•	
Continued.	from	previous	page

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Aromatic	33/144~(23%)	22/68~(32%)	11/68~(16%)	0/8~(0%)
Overall	806/2066~(39%)	500/1144~(44%)	237/728~(33%)	69/194~(36%)

#### 7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	А	17	TYR	HD2	8.54	5.48 - 8.39	5.5
1	А	17	TYR	HD1	8.54	5.49 - 8.39	5.5

#### 7.1.5 Random Coil Index (RCI) plots (1)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





# 8 NMR restraints analysis (i)

### 8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1776
Intra-residue ( i-j =0)	498
Sequential ( i-j =1)	588
Medium range ( $ i-j >1$ and $ i-j <5$ )	340
Long range $( i-j  \ge 5)$	178
Inter-chain	172
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	10.3
Number of long range restraints per residue <sup>1</sup>	1.0

<sup>1</sup>Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

### 8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

#### 8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	43.0	0.2
0.2-0.5 (Medium)	76.1	0.5
>0.5 (Large)	37.4	3.78



#### 8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than  $1^\circ$  are not included in the calculation. There are no dihedral-angle violations



# 9 Distance violation analysis (i)


### 9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destroints type	Count	$\%^1$	$\mathbf{Violated}^3$			Consistently		Violated <sup>4</sup>
Restraints type	Count	70-	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue $( i-j =0)$	498	28.0	36	7.2	2.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	424	23.9	34	8.0	1.9	0	0.0	0.0
Sidechain-Sidechain	74	4.2	2	2.7	0.1	0	0.0	0.0
Sequential $( i-j =1)$	588	33.1	87	14.8	4.9	1	0.2	0.1
Backbone-Backbone	202	11.4	10	5.0	0.6	0	0.0	0.0
Backbone-Sidechain	348	19.6	61	17.5	3.4	1	0.3	0.1
Sidechain-Sidechain	38	2.1	16	42.1	0.9	0	0.0	0.0
Medium range ( $ i-j  > 1 \&  i-j  < 5$ )	340	19.1	140	41.2	7.9	1	0.3	0.1
Backbone-Backbone	120	6.8	35	29.2	2.0	1	0.8	0.1
Backbone-Sidechain	152	8.6	69	45.4	3.9	0	0.0	0.0
Sidechain-Sidechain	68	3.8	36	52.9	2.0	0	0.0	0.0
Long range $( i-j  \ge 5)$	178	10.0	97	54.5	5.5	7	3.9	0.4
Backbone-Backbone	28	1.6	22	78.6	1.2	5	17.9	0.3
Backbone-Sidechain	82	4.6	46	56.1	2.6	2	2.4	0.1
Sidechain-Sidechain	68	3.8	29	42.6	1.6	0	0.0	0.0
Inter-chain	172	9.7	66	38.4	3.7	5	2.9	0.3
Backbone-Backbone	3	0.2	1	33.3	0.1	0	0.0	0.0
Backbone-Sidechain	54	3.0	17	31.5	1.0	0	0.0	0.0
Sidechain-Sidechain	115	6.5	48	41.7	2.7	5	4.3	0.3
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1776	100.0	426	24.0	24.0	14	0.8	0.8
Backbone-Backbone	353	19.9	68	19.3	3.8	6	1.7	0.3
Backbone-Sidechain	1060	59.7	227	21.4	12.8	3	0.3	0.2
Sidechain-Sidechain	363	20.4	131	36.1	7.4	5	1.4	0.3

 $^1$  percentage calculated with respect to the total number of distance restraints,  $^2$  percentage calculated with respect to the number of restraints in a particular restraint category,  $^3$  violated in at least one model,  $^4$  violated in all the models





#### 9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

#### 9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	$SD^6$ (Å)	Median (Å)
Model ID	$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (A)	Max (A)	$SD^{*}(A)$	Median (A)
1	8	32	42	41	30	153	0.4	2.38	0.31	0.37
2	13	24	48	40	31	156	0.39	2.33	0.34	0.33
3	10	19	31	47	30	137	0.42	2.42	0.35	0.39
4	14	24	40	50	32	160	0.39	2.35	0.36	0.32
5	11	28	39	49	28	155	0.39	2.23	0.28	0.34
6	9	21	42	49	29	150	0.39	2.22	0.34	0.31
7	5	32	40	41	29	147	0.4	3.78	0.38	0.33
8	15	26	33	41	29	144	0.38	2.43	0.28	0.34
9	12	33	41	40	29	155	0.37	2.18	0.26	0.33
10	17	41	42	43	31	174	0.38	2.11	0.3	0.33
11	15	33	38	40	32	158	0.36	2.38	0.28	0.3

Continued on next page...




Madal ID	_	Nun	nber o	f viola	ations	5	Mean (Å)		$SD^{6}$ (Å)	Madian (Å)
Model ID	$\mathrm{IR}^{1}$	$SQ^2$	$MR^3$	$LR^4$	$  IC^5  $	Total	Mean (A)	Max (Å)	$SD^{*}(A)$	Median (Å)
12	11	28	27	42	31	139	0.39	2.39	0.33	0.33
13	11	31	42	51	32	167	0.39	2.66	0.34	0.31
14	16	32	37	39	36	160	0.39	2.05	0.32	0.32
15	10	26	38	50	26	150	0.39	2.89	0.31	0.36
16	10	27	49	50	44	180	0.41	3.42	0.41	0.33
17	10	32	45	48	29	164	0.37	2.45	0.33	0.29
18	11	38	40	40	33	162	0.4	2.35	0.34	0.35
19	15	30	38	44	33	160	0.38	2.4	0.31	0.33
20	11	32	49	40	25	157	0.38	2.49	0.3	0.35

Continued from previous page...

 $^1$ Intra-residue restraints,  $^2$ S<br/>equential restraints,  $^3$ Medium range restraints,<br/>  $^4$ Long range restraints,  $^5$ Inter-chain restraints,<br/>  $^6$ Standard deviation





The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right



### 9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1350(IR:462, SQ:501, MR:200, LR:81, IC:106) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	restra	aints	Fractio	n of the ensemble
$IR^1$	$SQ^2$	$MR^3$	LR <sup>4</sup>	IC <sup>5</sup>	Total	$\operatorname{Count}^6$	%
7	20	32	13	6	78	1	5.0
6	10	17	8	6	47	2	10.0
2	5	13	8	3	31	3	15.0
4	6	10	5	2	27	4	20.0
1	5	11	4	1	22	5	25.0
0	4	6	3	6	19	6	30.0
2	1	9	3	7	22	7	35.0
3	3	8	2	7	23	8	40.0
2	6	7	4	1	20	9	45.0
1	4	4	5	2	16	10	50.0
0	4	6	4	2	16	11	55.0
1	2	3	7	2	15	12	60.0
1	4	2	6	3	16	13	65.0
1	2	2	1	0	6	14	70.0
1	1	0	4	2	8	15	75.0
2	3	3	2	1	11	16	80.0
0	3	1	3	4	11	17	85.0
2	3	1	4	4	14	18	90.0
0	0	4	4	2	10	19	95.0
0	1	1	7	5	14	20	100.0


 $^{1}$ Intra-residue restraints,  $^{2}$ Sequential restraints,  $^{3}$ Medium range restraints,  $^{4}$ Long range restraints,  $^{5}$ Inter-chain restraints,  $^{6}$  Number of models with violations



Intra-residue

Sequential MediumRange

LongRange InterChain




#### 9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

### 9.4 Most violated distance restraints in the ensemble (i)

#### 9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





#### 9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	$\mathbf{Models}^1$	Mean (Å)	$SD^1$ (Å)	Median (Å)
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	20	2.26	0.6	2.18
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD22	20	2.26	0.6	2.18
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD23	20	2.26	0.6	2.18
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD11	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD12	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD13	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD11	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD12	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD13	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD11	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD12	20	1.81	0.5	1.79
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD13	20	1.81	0.5	1.79
(1,914)	1:A:80:LEU:HD21	1:B:81:ARG:HB2	20	1.53	0.41	1.42
(1,914)	1:A:80:LEU:HD22	1:B:81:ARG:HB2	20	1.53	0.41	1.42
(1,914)	1:A:80:LEU:HD23	1:B:81:ARG:HB2	20	1.53	0.41	1.42
(1,156)	1:A:19:GLN:HB2	1:B:52:MET:HG3	20	1.21	0.6	1.21

Continued on next page...



Key	Atom-1	Atom-2	$Models^1$	Mean (Å)	$SD^1$ (Å)	Median (Å)
(1,62)	1:A:12:VAL:HG11	1:A:32:ILE:H	20	0.49	0.08	0.52
(1,62)	1:A:12:VAL:HG12	1:A:32:ILE:H	20	0.49	0.08	0.52
(1,62)	1:A:12:VAL:HG13	1:A:32:ILE:H	20	0.49	0.08	0.52
(1,62)	1:A:12:VAL:HG21	1:A:32:ILE:H	20	0.49	0.08	0.52
(1,62)	1:A:12:VAL:HG22	1:A:32:ILE:H	20	0.49	0.08	0.52
(1,62)	1:A:12:VAL:HG23	1:A:32:ILE:H	20	0.49	0.08	0.52
(1,973)	1:A:84:LEU:HD11	1:A:85:LYS:H	20	0.49	0.09	0.49
(1,973)	1:A:84:LEU:HD12	1:A:85:LYS:H	20	0.49	0.09	0.49
(1,973)	1:A:84:LEU:HD13	1:A:85:LYS:H	20	0.49	0.09	0.49
(1,973)	1:A:84:LEU:HD21	1:A:85:LYS:H	20	0.49	0.09	0.49
(1,973)	1:A:84:LEU:HD22	1:A:85:LYS:H	20	0.49	0.09	0.49
(1,973)	1:A:84:LEU:HD23	1:A:85:LYS:H	20	0.49	0.09	0.49
(1,1043)	1:B:12:VAL:HG11	1:B:32:ILE:H	20	0.47	0.09	0.49
(1,1043)	1:B:12:VAL:HG12	1:B:32:ILE:H	20	0.47	0.09	0.49
(1,1043)	1:B:12:VAL:HG13	1:B:32:ILE:H	20	0.47	0.09	0.49
(1,1043)	1:B:12:VAL:HG21	1:B:32:ILE:H	20	0.47	0.09	0.49
(1,1043)	1:B:12:VAL:HG22	1:B:32:ILE:H	20	0.47	0.09	0.49
(1,1043)	1:B:12:VAL:HG23	1:B:32:ILE:H	20	0.47	0.09	0.49
(1,258)	1:A:32:ILE:HA	1:A:36:ALA:H	20	0.47	0.13	0.52
(1,113)	1:A:16:ILE:H	1:A:21:TYR:H	20	0.47	0.09	0.51
(1,1034)	1:B:12:VAL:H	1:B:25:GLY:H	20	0.44	0.13	0.51

Continued from previous page...


<sup>1</sup>Number of violated models, <sup>2</sup>Standard deviation

### 9.5 All violated distance restraints (i)

#### 9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





#### 9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	7	3.78
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD22	7	3.78
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD23	7	3.78
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	16	3.42
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD22	16	3.42
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD23	16	3.42
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	15	2.89
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD22	15	2.89
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD23	15	2.89
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	13	2.66
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD22	13	2.66
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD23	13	2.66
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD11	16	2.5
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD12	16	2.5
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD13	16	2.5
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD11	16	2.5
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD12	16 	2.5

Continued on next page...



Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD13	16	2.5
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD11	16	2.5
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD12	16	2.5
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD13	16	2.5
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD11	20	2.49
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD12	20	2.49
(1,597)	1:A:61:LEU:HD21	1:B:23:ILE:HD13	20	2.49
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD11	20	2.49
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD12	20	2.49
(1,597)	1:A:61:LEU:HD22	1:B:23:ILE:HD13	20	2.49
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD11	20	2.49
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD12	20	2.49
(1,597)	1:A:61:LEU:HD23	1:B:23:ILE:HD13	20	2.49
(1,158)	1:A:19:GLN:HG3	1:B:52:MET:HG3	16	2.47
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	17	2.45
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD22	17	2.45
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD23	17	2.45
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	8	2.43
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD22	8	2.43
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD23	8	2.43
(1,751)	1:A:71:LEU:HB2	1:B:34:LEU:HD21	3	2.42

Continued from previous page...



# 10 Dihedral-angle violation analysis (i)

No dihedral-angle restraints found

