

# Full wwPDB X-ray Structure Validation Report (i)

#### Sep 12, 2023 – 02:02 AM EDT

PDB ID : 4MMW

Title : Crystal structure of D-glucarate dehydratase from Agrobacterium tumefaciens

complexed with magnesium, L-Xylarohydroxamate and L-Lyxarohydroxamate

Authors: Fedorov, A.A.; Fedorov, E.V.; Sakai, A.; Gerlt, J.A.; Almo, S.C.

Deposited on : 2013-09-09

Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$ 

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4: 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\# \text{Entries, resolution range}(\mathring{A}))$		
$R_{free}$	130704	3122 (1.66-1.62)		
Clashscore	141614	3268 (1.66-1.62)		
Ramachandran outliers	138981	3215 (1.66-1.62)		
Sidechain outliers	138945	3215 (1.66-1.62)		
RSRZ outliers	127900	3079 (1.66-1.62)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	405	91%	•	6%			
1	В	405	89%	5%	6%			

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	XYH	В	404[A]	-	X	-	-
6	PGE	A	411	-	-	X	-



## 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 6920 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Isomerase/lactonizing enzyme.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	A	381	Total 3055	C 1923	N 552	O 563	S 17	0	5	0
1	В	381	Total 3046	C 1918	N 548	O 563	S 17	0	4	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	expression tag	UNP Q7CSI0
A	-21	GLY	-	expression tag	UNP Q7CSI0
A	-20	SER	-	expression tag	UNP Q7CSI0
A	-19	SER	-	expression tag	UNP Q7CSI0
A	-18	HIS	-	expression tag	UNP Q7CSI0
A	-17	HIS	-	expression tag	UNP Q7CSI0
A	-16	HIS	-	expression tag	UNP Q7CSI0
A	-15	HIS	-	expression tag	UNP Q7CSI0
A	-14	HIS	-	expression tag	UNP Q7CSI0
A	-13	HIS	-	expression tag	UNP Q7CSI0
A	-12	SER	-	expression tag	UNP Q7CSI0
A	-11	SER	-	expression tag	UNP Q7CSI0
A	-10	GLY	-	expression tag	UNP Q7CSI0
A	-9	LEU	-	expression tag	UNP Q7CSI0
A	-8	VAL	-	expression tag	UNP Q7CSI0
A	-7	PRO	-	expression tag	UNP Q7CSI0
A	-6	ARG	-	expression tag	UNP Q7CSI0
A	-5	GLY	-	expression tag	UNP Q7CSI0
A	-4	SER	-	expression tag	UNP Q7CSI0
A	-3	HIS	-	expression tag	UNP Q7CSI0
A	-2	MET	-	expression tag	UNP Q7CSI0
A	-1	ILE	-	expression tag	UNP Q7CSI0
A	0	ILE	-	expression tag	UNP Q7CSI0
В	-22	MET	-	expression tag	UNP Q7CSI0
В	-21	GLY	-	expression tag	UNP Q7CSI0



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	-20	SER	-	expression tag	UNP Q7CSI0
В	-19	SER	-	expression tag	UNP Q7CSI0
В	-18	HIS	-	expression tag	UNP Q7CSI0
В	-17	HIS	-	expression tag	UNP Q7CSI0
В	-16	HIS	-	expression tag	UNP Q7CSI0
В	-15	HIS	-	expression tag	UNP Q7CSI0
В	-14	HIS	-	expression tag	UNP Q7CSI0
В	-13	HIS	-	expression tag	UNP Q7CSI0
В	-12	SER	-	expression tag	UNP Q7CSI0
В	-11	SER	-	expression tag	UNP Q7CSI0
В	-10	GLY	-	expression tag	UNP Q7CSI0
В	-9	LEU	-	expression tag	UNP Q7CSI0
В	-8	VAL	-	expression tag	UNP Q7CSI0
В	-7	PRO	-	expression tag	UNP Q7CSI0
В	-6	ARG	-	expression tag	UNP Q7CSI0
В	-5	GLY	-	expression tag	UNP Q7CSI0
В	-4	SER	-	expression tag	UNP Q7CSI0
В	-3	HIS	-	expression tag	UNP Q7CSI0
В	-2	MET	-	expression tag	UNP Q7CSI0
В	-1	ILE	-	expression tag	UNP Q7CSI0
В	0	ILE	-	expression tag	UNP Q7CSI0

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Mg 3 3	0	0
2	В	3	Total Mg 3 3	0	0


• Molecule 3 is XYLAROHYDROXAMATE (three-letter code: XYH) (formula:  $C_5H_8NO_7$ ).





Mol	Chain	Residues	Ato	ms		ZeroOcc	AltConf
3	A	1	Total C 13 5			0	1
3	В	1	Total C	N 1	O 7	0	1

• Molecule 4 is (2R,3S,4R)-2,3,4-TRIHYDROXY-5-(HYDROXYAMINO)-5-OXOPENTANO IC ACID (three-letter code: LLH) (formula:  $C_5H_9NO_7$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total 13	C 5	N 1	O 7	0	1




Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	В	1	Total 13	C 5	_	O 7	0	1

 $\bullet$  Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	5	Total Cl 5 5	0	0
5	В	5	Total Cl 5 5	0	0

 $\bullet$  Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $\mathrm{C_6H_{14}O_4}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 10 6 4	0	0

• Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	375	Total O 375 375	0	0
7	В	366	Total O 366 366	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Isomerase/lactonizing enzyme





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.10Å 118.10Å 133.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	$\begin{array}{r} 30.00 & 30.00 & 30.00 \\ \hline 49.10 & - & 1.65 \\ 49.10 & - & 1.65 \end{array}$	Depositor EDS
% Data completeness	99.5 (49.10-1.65)	Depositor
(in resolution range)	99.5 (49.10-1.65)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.09 (at 1.64Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
D.D.	0.170 , 0.194	Depositor
$R, R_{free}$	0.172 , $0.195$	DCC
$R_{free}$ test set	5473 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.0	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37, 32.0	EDS
L-test for twinning <sup>2</sup>	$< L >=0.52, < L^2>=0.36$	Xtriage
Estimated twinning fraction	0.478 for -h,k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6920	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: XYH, LLH, MG, PGE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.34	0/3133	0.53	0/4249	
1	В	0.34	0/3124	0.52	0/4238	
All	All	0.34	0/6257	0.52	0/8487	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3055	0	2963	8	0
1	В	3046	0	2950	14	0
2	A	3	0	0	0	0
2	В	3	0	0	0	0
3	A	13	0	8	0	0
3	В	13	0	8	0	0
4	A	13	0	7	0	0
4	В	13	0	7	0	0
5	A	5	0	0	1	0
5	В	5	0	0	1	0
6	A	10	0	14	7	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	375	0	0	1	0
7	В	366	0	0	1	0
All	All	6920	0	5957	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:A:261:ALA:H	6:A:411:PGE:H1	1.59	0.68
6:A:411:PGE:C5	1:B:261:ALA:H	2.12	0.62
1:B:221[B]:ARG:NH1	7:B:840:HOH:O	2.28	0.60
6:A:411:PGE:H6	1:B:238:GLN:HG2	1.84	0.58
1:A:261:ALA:H	6:A:411:PGE:C1	2.18	0.56
6:A:411:PGE:H52	1:B:261:ALA:H	1.74	0.53
6:A:411:PGE:H5	1:B:261:ALA:H	1.73	0.53
1:B:202:ARG:NE	1:B:229[B]:ASP:OD2	2.45	0.49
1:A:257:PRO:HB3	1:A:270:TRP:CZ2	2.49	0.48
1:B:257:PRO:HB3	1:B:270:TRP:CZ2	2.50	0.47
1:B:263:LYS:HD2	1:B:265:TRP:HZ3	1.80	0.47
1:B:46:GLN:HG2	1:B:115:SER:OG	2.15	0.46
1:B:190:CYS:SG	1:B:205:ILE:HD12	2.56	0.46
1:A:158:GLU:OE2	7:A:769:HOH:O	2.20	0.46
1:A:171:HIS:CD2	1:A:206:ASP:HB3	2.51	0.46
1:B:1:MET:HG3	5:B:410:CL:CL	2.54	0.44
1:A:166:LYS:NZ	1:A:356:ASP:O	2.49	0.44
6:A:411:PGE:H5	1:B:261:ALA:HB3	2.00	0.43
1:B:211:TYR:HB2	1:B:234:PRO:O	2.19	0.42
1:A:1:MET:HG3	5:A:410:CL:CL	2.57	0.41
1:B:27:PRO:HG3	1:B:334:HIS:CE1	2.57	0.40
1:A:347:LYS:HE3	1:A:372:ASP:HA	2.04	0.40

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	384/405~(95%)	374 (97%)	10 (3%)	0	100	100
1	В	383/405~(95%)	374 (98%)	9 (2%)	0	100	100
All	All	767/810 (95%)	748 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric   Outliers		Percentiles		
1	A	319/335~(95%)	317 (99%)	2 (1%)	86 75		
1	В	318/335 (95%)	316 (99%)	2 (1%)	86 75		
All	All	$637/670 \ (95\%)$	633 (99%)	4 (1%)	86 75		

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	HIS
1	A	230	TRP
1	В	31	HIS
1	В	230	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 21 ligands modelled in this entry, 16 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Res Link Bond lengths		Bond angles				
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	LLH	В	405[B]	2	12,12,12	0.81	0	14,16,16	0.99	0
6	PGE	A	411	-	9,9,9	0.61	0	8,8,8	1.81	2 (25%)
3	XYH	A	404[A]	2	12,12,12	2.68	4 (33%)	14,16,16	4.22	7 (50%)
4	LLH	A	405[B]	2	12,12,12	0.80	0	14,16,16	0.91	0
3	XYH	В	404[A]	2	12,12,12	2.70	4 (33%)	14,16,16	4.23	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LLH	В	405[B]	2	-	7/18/18/18	-
6	PGE	A	411	-	-	3/7/7/7	-
3	XYH	A	404[A]	2	-	7/18/18/18	-
4	LLH	A	405[B]	2	-	6/18/18/18	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	XYH	В	404[A]	2	-	8/18/18/18	-

#### All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\textup{\AA})$	$Ideal(\AA)$
3	В	404[A]	XYH	OH5-C5	8.08	1.39	1.23
3	A	404[A]	XYH	OH5-C5	8.02	1.39	1.23
3	В	404[A]	XYH	C5-N6	-2.57	1.30	1.33
3	A	404[A]	XYH	C5-N6	-2.46	1.30	1.33
3	A	404[A]	XYH	O1B-C1	2.29	1.29	1.22
3	В	404[A]	XYH	O1B-C1	2.28	1.29	1.22
3	A	404[A]	XYH	O1A-C1	-2.21	1.23	1.30
3	В	404[A]	XYH	O1A-C1	-2.20	1.23	1.30

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	404[A]	XYH	OH4-C4-C5	-13.31	81.95	110.63
3	В	404[A]	XYH	OH4-C4-C5	-13.19	82.21	110.63
3	В	404[A]	XYH	OH4-C4-C3	-4.06	101.84	110.45
3	A	404[A]	XYH	OH4-C4-C3	-3.77	102.47	110.45
3	В	404[A]	XYH	C3-C4-C5	3.46	118.32	109.91
3	A	404[A]	XYH	C3-C4-C5	3.46	118.32	109.91
3	В	404[A]	XYH	OH5-C5-N6	-3.40	118.87	123.27
3	A	404[A]	XYH	OH5-C5-N6	-3.25	119.06	123.27
3	В	404[A]	XYH	OH3-C3-C2	3.09	114.83	109.21
3	В	404[A]	XYH	OH3-C3-C4	3.05	114.77	109.21
3	A	404[A]	XYH	OH3-C3-C2	3.03	114.73	109.21
3	A	404[A]	XYH	OH3-C3-C4	3.02	114.72	109.21
6	A	411	PGE	O3-C5-C6	2.81	122.42	110.07
3	A	404[A]	XYH	C3-C2-C1	2.38	112.31	109.32
3	В	404[A]	XYH	C3-C2-C1	2.35	112.28	109.32
6	A	411	PGE	O2-C2-C1	2.07	119.14	110.07

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	404[A]	XYH	OH3-C3-C4-C5
3	A	404[A]	XYH	OH4-C4-C5-OH5
3	A	404[A]	XYH	OH4-C4-C5-N6



Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	В	404[A]	XYH	OH3-C3-C4-C5
3	В	404[A]	XYH	OH4-C4-C5-OH5
3	В	404[A]	XYH	OH4-C4-C5-N6
4	A	405[B]	LLH	O1-C1-C2-C3
4	A	405[B]	LLH	N-C1-C2-C3
4	A	405[B]	LLH	O1-C1-C2-O2
4	A	405[B]	LLH	N-C1-C2-O2
4	В	405[B]	LLH	C1-C2-C3-O3
4	В	405[B]	LLH	O1-C1-C2-C3
4	В	405[B]	LLH	N-C1-C2-C3
4	В	405[B]	LLH	O1-C1-C2-O2
4	В	405[B]	LLH	N-C1-C2-O2
6	A	411	PGE	O2-C3-C4-O3
3	A	404[A]	XYH	C3-C4-C5-OH5
3	A	404[A]	XYH	C3-C4-C5-N6
3	В	404[A]	XYH	C3-C4-C5-OH5
3	В	404[A]	XYH	C3-C4-C5-N6
3	A	404[A]	XYH	OH2-C2-C3-OH3
3	В	404[A]	XYH	OH2-C2-C3-OH3
6	A	411	PGE	O3-C5-C6-O4
4	A	405[B]	LLH	C1-C2-C3-O3
3	A	404[A]	XYH	C2-C3-C4-C5
3	В	404[A]	XYH	C2-C3-C4-C5
4	В	405[B]	LLH	O2-C2-C3-C4
4	A	405[B]	LLH	O2-C2-C3-C4
3	В	404[A]	XYH	C1-C2-C3-OH3
6	A	411	PGE	C3-C4-O3-C5
4	В	405[B]	LLH	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	$\operatorname{Res}$	Type	Clashes	Symm-Clashes	
6	A	411	PGE	7	0	

# 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		Z>2	$OWAB(A^2)$	Q<0.9
1	A	381/405 (94%)	-0.68	0	100	100	13, 18, 31, 42	0
1	В	381/405 (94%)	-0.69	0	100	100	13, 18, 30, 41	0
All	All	762/810 (94%)	-0.68	0	100	100	13, 18, 31, 42	0

There are no RSRZ outliers to report.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathrm{\AA}^2)$	Q<0.9
6	PGE	A	411	10/10	0.84	0.14	30,40,45,47	0
5	CL	В	410	1/1	0.92	0.08	42,42,42,42	0
2	MG	В	401	1/1	0.93	0.08	22,22,22,22	1
5	CL	A	410	1/1	0.93	0.07	41,41,41,41	0
2	MG	A	401	1/1	0.95	0.16	35,35,35,35	0
3	XYH	A	404[A]	13/13	0.95	0.10	12,21,26,28	13
4	LLH	A	405[B]	13/13	0.95	0.10	13,22,25,26	13



 $Continued\ from\ previous\ page...$ 

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
4	LLH	В	405[B]	13/13	0.96	0.09	13,23,27,29	13
3	XYH	В	404[A]	13/13	0.96	0.10	13,22,27,30	13
5	CL	A	409	1/1	0.98	0.03	31,31,31,31	0
5	CL	A	406	1/1	0.98	0.05	22,22,22,22	0
5	CL	A	407	1/1	0.98	0.06	28,28,28,28	0
5	CL	A	408	1/1	0.98	0.07	31,31,31,31	0
5	CL	В	406	1/1	0.99	0.04	21,21,21,21	0
5	CL	В	408	1/1	0.99	0.05	29,29,29,29	0
5	CL	В	409	1/1	0.99	0.04	30,30,30,30	0
2	MG	В	403	1/1	0.99	0.09	14,14,14,14	0
2	MG	A	402	1/1	0.99	0.06	15,15,15,15	0
2	MG	A	403	1/1	1.00	0.07	15,15,15,15	0
5	CL	В	407	1/1	1.00	0.06	24,24,24,24	0
2	MG	В	402	1/1	1.00	0.04	14,14,14,14	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

