

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 04:59 AM EDT

PDB ID : 6MNI

Title : Structure of the tandem CACHE domain of PscC Authors : Johnston, J.M.; Gerth, M.L.; Ehrhardt, M.K.G.

Deposited on : 2018-10-01

Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Methyl-accepting chemotaxis protein.

\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	258	Total	С	N	О	S	0	3	0
1	Λ	250	1995	1261	337	388	9	U		
1	B	251	Total	С	N	О	S	0	4	0
1	D	201	1929	1222	324	374	9		4	

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	MET	-	initiating methionine	UNP A0A2V0QX35
A	8	GLY	_	expression tag	UNP A0A2V0QX35
A	9	SER	-	expression tag	UNP A0A2V0QX35
A	10	SER	-	expression tag	UNP A0A2V0QX35
A	11	HIS	-	expression tag	UNP A0A2V0QX35
A	12	HIS	-	expression tag	UNP A0A2V0QX35
A	13	HIS	-	expression tag	UNP A0A2V0QX35
A	14	HIS	-	expression tag	UNP A0A2V0QX35
A	15	HIS	-	expression tag	UNP A0A2V0QX35
A	16	HIS	-	expression tag	UNP A0A2V0QX35
A	17	SER	-	expression tag	UNP A0A2V0QX35
A	18	SER	-	expression tag	UNP A0A2V0QX35
A	19	GLY	-	expression tag	UNP A0A2V0QX35
A	20	LEU	-	expression tag	UNP A0A2V0QX35
A	21	VAL	-	expression tag	UNP A0A2V0QX35
A	22	PRO	-	expression tag	UNP A0A2V0QX35
A	23	ARG	-	expression tag	UNP A0A2V0QX35
A	24	GLY	-	expression tag	UNP A0A2V0QX35
A	25	SER	-	expression tag	UNP A0A2V0QX35
A	26	HIS	-	expression tag	UNP A0A2V0QX35
A	27	MET	_	expression tag	UNP A0A2V0QX35
В	7	MET	-	initiating methionine	UNP A0A2V0QX35
В	8	GLY	-	expression tag	UNP A0A2V0QX35
В	9	SER	-	expression tag	UNP A0A2V0QX35
В	10	SER	-	expression tag	UNP A0A2V0QX35


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	11	HIS	-	expression tag	UNP A0A2V0QX35
В	12	HIS	-	expression tag	UNP A0A2V0QX35
В	13	HIS	-	expression tag	UNP A0A2V0QX35
В	14	HIS	-	expression tag	UNP A0A2V0QX35
В	15	HIS	-	expression tag	UNP A0A2V0QX35
В	16	HIS	-	expression tag	UNP A0A2V0QX35
В	17	SER	-	expression tag	UNP A0A2V0QX35
В	18	SER	-	expression tag	UNP A0A2V0QX35
В	19	GLY	-	expression tag	UNP A0A2V0QX35
В	20	LEU	-	expression tag	UNP A0A2V0QX35
В	21	VAL	-	expression tag	UNP A0A2V0QX35
В	22	PRO	-	expression tag	UNP A0A2V0QX35
В	23	ARG	-	expression tag	UNP A0A2V0QX35
В	24	GLY	-	expression tag	UNP A0A2V0QX35
В	25	SER	-	expression tag	UNP A0A2V0QX35
В	26	HIS	-	expression tag	UNP A0A2V0QX35
В	27	MET	-	expression tag	UNP A0A2V0QX35


• Molecule 2 is PROLINE (three-letter code: PRO) (formula: $C_5H_9NO_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 8		N 1		0	0
2	В	1	Total 8	C 5	N 1	O 2	0	0



• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 6	C 3	O 3	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total Cl 1 1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	188	Total O 188 188	0	0
5	В	194	Total O 194 194	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants	34.24Å 53.92Å 73.31Å	Depositor
a, b, c, α , β , γ	104.79° 91.20° 92.71°	Depositor
Resolution (Å)	48.27 - 1.70	Depositor
% Data completeness	96.6 (48.27-1.70)	Depositor
(in resolution range)	, , ,	•
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.37 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.162 , 0.200	Depositor
Wilson B-factor (\mathring{A}^2)	22.4	Xtriage
Anisotropy	0.215	Xtriage
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4329	wwPDB-VP
Average B, all atoms $(Å^2)$	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 10.34% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Res	Link	В	ond leng	gths	Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PRO	В	301	-	8,8,8	0.96	1 (12%)	10,10,10	0.95	0
2	PRO	A	301	-	8,8,8	0.78	0	10,10,10	1.16	1 (10%)
3	GOL	A	302	-	5,5,5	0.99	0	5,5,5	1.34	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PRO	В	301	_	-	0/4/11/11	0/1/1/1
2	PRO	A	301	-	-	0/4/11/11	0/1/1/1
3	GOL	A	302	_	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
2	В	301	PRO	OXT-C	-2.25	1.23	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	A	302	GOL	C3-C2-C1	-2.30	102.77	111.70
2	A	301	PRO	OXT-C-CA	2.10	120.37	113.40

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	GOL	O1-C1-C2-C3
3	A	302	GOL	O1-C1-C2-O2


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In



addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

