



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2020 – 10:52 AM BST

PDB ID : 3MQE
Title : Structure of SC-75416 bound at the COX-2 active site
Authors : Wang, J.L.; Limburg, D.; Graneto, M.J.; Springer, J.; Rogier, J.; Kiefer, J.R.
Deposited on : 2010-04-28
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION







The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	 78% 15% • 6%
1	B	587	 80% 12% • 6%
1	C	587	 82% 11% • 6%
1	D	587	 81% 11% • 6%
2	E	3	 100%
2	F	3	 33% 67%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	3	 100%
2	H	3	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	G	3	-	-	-	X
2	NAG	H	3	-	-	-	X
4	NAG	A	661	X	-	-	-
4	NAG	B	661	X	-	-	-
4	NAG	C	661	X	-	-	-
4	NAG	D	661	X	-	-	-

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 18573 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostaglandin G/H synthase 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	552	4474	2885	750	814	25	0	0	0
1	B	552	4474	2885	750	814	25	0	0	0
1	C	552	4474	2885	750	814	25	0	0	0
1	D	552	4474	2885	750	814	25	0	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	3	42	24	3	15	0	0	0
2	F	3	42	24	3	15	0	0	0
2	G	3	42	24	3	15	0	0	0
2	H	3	42	24	3	15	0	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
3	A	1	43	34	1	4	4	0	0
3	B	1	43	34	1	4	4	0	0
3	C	1	43	34	1	4	4	0	0
3	D	1	43	34	1	4	4	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula: C₁₄H₂₈O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			20	14	6		
5	C	1	Total	C	O	0	0
			20	14	6		

- Molecule 6 is (2S)-7-tert-butyl-6-chloro-2-(trifluoromethyl)-2H-chromene-3-carboxylic acid (three-letter code: 416) (formula: C₁₅H₁₄ClF₃O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	F	O		
6	A	1	22	15	1	3	3	0	0
6	B	1	22	15	1	3	3	0	0
6	C	1	22	15	1	3	3	0	0
6	D	1	22	15	1	3	3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	31	Total	O	0	0
			31	31		
7	B	24	Total	O	0	0
			24	24		
7	C	23	Total	O	0	0
			23	23		
7	D	19	Total	O	0	0
			19	19		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Prostaglandin G/H synthase 2




- Molecule 1: Prostaglandin G/H synthase 2





- Molecule 1: Prostaglandin G/H synthase 2





- Molecule 1: Prostaglandin G/H synthase 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  33% 67%

MAG1
MAG2
MAG3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	183.70Å 140.38Å 129.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 34.28 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (20.00-2.80) 99.1 (34.28-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.236 , 0.290 0.236 , 0.292	Depositor DCC
R_{free} test set	8283 reflections (10.05%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtrriage
Anisotropy	0.261	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	18573	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8586e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, 416, NAG, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/4601	0.62	1/6239 (0.0%)
1	B	0.50	0/4601	0.60	0/6239
1	C	0.47	0/4601	0.58	0/6239
1	D	0.48	0/4601	0.59	0/6239
All	All	0.49	0/18404	0.60	1/24956 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	215	ASP	CB-CG-OD1	5.34	123.11	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4474	0	4373	50	0
1	B	4474	0	4373	46	0
1	C	4474	0	4373	42	0
1	D	4474	0	4373	40	0
2	E	42	0	37	0	0
2	F	42	0	37	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	G	42	0	37	0	0
2	H	42	0	37	1	0
3	A	43	0	30	2	0
3	B	43	0	30	1	0
3	C	43	0	30	1	0
3	D	43	0	30	0	0
4	A	28	0	26	0	0
4	B	28	0	26	0	0
4	C	28	0	26	0	0
4	D	28	0	26	0	0
5	A	20	0	28	0	0
5	C	20	0	28	0	0
6	A	22	0	13	2	0
6	B	22	0	13	2	0
6	C	22	0	13	1	0
6	D	22	0	13	3	0
7	A	31	0	0	0	0
7	B	24	0	0	1	0
7	C	23	0	0	0	0
7	D	19	0	0	0	0
All	All	18573	0	17972	185	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 185 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:328:LYS:NZ	1:D:546:ASN:O	2.13	0.81
1:C:513:ALA:HB3	1:C:514:PRO:HD3	1.63	0.80
1:B:128:PHE:O	1:B:362:ARG:NH2	2.18	0.76
1:C:179:GLY:O	1:C:568:VAL:HG23	1.85	0.76
1:A:184:PHE:CZ	1:A:338:LEU:HD11	2.22	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/587 (94%)	519 (94%)	29 (5%)	2 (0%)	34	66
1	B	550/587 (94%)	519 (94%)	30 (6%)	1 (0%)	47	78
1	C	550/587 (94%)	528 (96%)	21 (4%)	1 (0%)	47	78
1	D	550/587 (94%)	512 (93%)	38 (7%)	0	100	100
All	All	2200/2348 (94%)	2078 (94%)	118 (5%)	4 (0%)	47	78

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	HIS
1	B	212	HIS
1	A	384	GLU
1	C	568	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/525 (94%)	466 (94%)	27 (6%)	21	52
1	B	493/525 (94%)	466 (94%)	27 (6%)	21	52
1	C	493/525 (94%)	465 (94%)	28 (6%)	20	50
1	D	493/525 (94%)	464 (94%)	29 (6%)	19	49
All	All	1972/2100 (94%)	1861 (94%)	111 (6%)	21	51

5 of 111 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	478	LYS
1	C	171	ARG
1	D	405	LEU
1	B	494	LEU
1	C	86	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	313	GLN
1	B	360	GLN
1	C	374	HIS
1	B	264	HIS
1	C	190	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	E	1	1,2	14,14,15	0.47	0	17,19,21	2.44	3 (17%)
2	NAG	E	2	2	14,14,15	0.61	0	17,19,21	2.06	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	E	3	2	14,14,15	0.72	1 (7%)	17,19,21	2.00	5 (29%)
2	NAG	F	1	1,2	14,14,15	0.72	0	17,19,21	1.05	0
2	NAG	F	2	2	14,14,15	0.47	0	17,19,21	1.68	4 (23%)
2	NAG	F	3	2	14,14,15	0.91	1 (7%)	17,19,21	1.86	3 (17%)
2	NAG	G	1	1,2	14,14,15	0.65	0	17,19,21	1.74	2 (11%)
2	NAG	G	2	2	14,14,15	0.45	0	17,19,21	1.70	5 (29%)
2	NAG	G	3	2	14,14,15	0.98	2 (14%)	17,19,21	2.57	7 (41%)
2	NAG	H	1	1,2	14,14,15	0.64	0	17,19,21	1.43	2 (11%)
2	NAG	H	2	2	14,14,15	0.69	0	17,19,21	1.38	2 (11%)
2	NAG	H	3	2	14,14,15	0.89	1 (7%)	17,19,21	2.45	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	3/6/23/26	0/1/1/1
2	NAG	E	3	2	-	2/6/23/26	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	1/6/23/26	0/1/1/1
2	NAG	F	3	2	-	2/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	3/6/23/26	0/1/1/1
2	NAG	G	3	2	-	2/6/23/26	0/1/1/1
2	NAG	H	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	1/6/23/26	0/1/1/1
2	NAG	H	3	2	-	1/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	3	NAG	C1-C2	2.49	1.56	1.52
2	F	3	NAG	C1-C2	2.32	1.55	1.52
2	G	3	NAG	C1-C2	2.30	1.55	1.52
2	G	3	NAG	C2-N2	2.18	1.50	1.46
2	E	3	NAG	C1-C2	2.08	1.55	1.52

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	NAG	C1-O5-C5	8.62	123.88	112.19
2	H	3	NAG	C1-O5-C5	6.40	120.86	112.19
2	G	3	NAG	C2-N2-C7	5.99	131.43	122.90
2	E	3	NAG	C1-O5-C5	5.79	120.04	112.19
2	G	1	NAG	C1-O5-C5	5.50	119.65	112.19

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	H	3	NAG	C3-C2-N2-C7
2	F	2	NAG	C3-C2-N2-C7
2	G	2	NAG	O5-C5-C6-O6
2	E	2	NAG	O5-C5-C6-O6
2	G	2	NAG	C4-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	2	NAG	1	0
2	H	3	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	HEM	D	605	1	27,50,50	2.29	6 (22%)	17,82,82	1.60	3 (17%)
4	NAG	D	681	1	14,14,15	0.48	0	17,19,21	0.94	1 (5%)
3	HEM	B	605	1	27,50,50	2.18	7 (25%)	17,82,82	1.48	2 (11%)
4	NAG	B	661	1	14,14,15	0.59	0	17,19,21	1.78	5 (29%)
6	416	A	701	-	19,23,23	1.15	1 (5%)	29,37,37	1.26	2 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	605	1	27,50,50	2.30	7 (25%)	17,82,82	1.83	4 (23%)
4	NAG	B	681	1	14,14,15	0.58	0	17,19,21	0.93	0
6	416	B	701	-	19,23,23	0.98	0	29,37,37	1.50	3 (10%)
4	NAG	A	681	1	14,14,15	0.57	0	17,19,21	1.09	1 (5%)
4	NAG	C	681	1	14,14,15	0.72	0	17,19,21	1.21	1 (5%)
5	BOG	A	704	-	20,20,20	0.54	0	25,25,25	0.61	0
6	416	C	701	-	19,23,23	1.03	1 (5%)	29,37,37	1.31	3 (10%)
4	NAG	A	661	1	14,14,15	0.82	1 (7%)	17,19,21	2.16	6 (35%)
6	416	D	701	-	19,23,23	1.14	2 (10%)	29,37,37	1.74	4 (13%)
4	NAG	C	661	1	14,14,15	0.48	0	17,19,21	1.20	1 (5%)
5	BOG	C	703	-	20,20,20	0.53	0	25,25,25	0.95	1 (4%)
4	NAG	D	661	1	14,14,15	0.75	1 (7%)	17,19,21	2.69	6 (35%)
3	HEM	C	605	1	27,50,50	2.21	6 (22%)	17,82,82	1.50	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	D	605	1	-	2/6/54/54	-
4	NAG	D	681	1	-	0/6/23/26	0/1/1/1
3	HEM	B	605	1	-	0/6/54/54	-
4	NAG	B	661	1	1/1/5/7	1/6/23/26	0/1/1/1
6	416	A	701	-	-	0/12/28/28	0/2/2/2
3	HEM	A	605	1	-	2/6/54/54	-
4	NAG	B	681	1	-	0/6/23/26	0/1/1/1
6	416	B	701	-	-	0/12/28/28	0/2/2/2
4	NAG	C	661	1	1/1/5/7	1/6/23/26	0/1/1/1
4	NAG	C	681	1	-	2/6/23/26	0/1/1/1
5	BOG	A	704	-	-	5/11/31/31	0/1/1/1
6	416	C	701	-	-	0/12/28/28	0/2/2/2
4	NAG	A	661	1	1/1/5/7	3/6/23/26	0/1/1/1
6	416	D	701	-	-	0/12/28/28	0/2/2/2
4	NAG	A	681	1	-	0/6/23/26	0/1/1/1
5	BOG	C	703	-	-	7/11/31/31	0/1/1/1
4	NAG	D	661	1	1/1/5/7	2/6/23/26	0/1/1/1
3	HEM	C	605	1	-	0/6/54/54	-

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	605	HEM	C3D-C2D	5.78	1.54	1.37
3	D	605	HEM	C3D-C2D	5.33	1.53	1.37
3	C	605	HEM	C3D-C2D	5.26	1.53	1.37
3	A	605	HEM	C3C-C2C	-5.16	1.33	1.40
3	D	605	HEM	C3C-C2C	-5.15	1.33	1.40

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	661	NAG	C1-O5-C5	6.27	120.68	112.19
6	D	701	416	C3-C7-C12	-5.78	115.09	122.54
4	D	661	NAG	C1-C2-N2	5.50	119.88	110.49
6	D	701	416	C14-C12-C7	4.86	124.48	118.43
6	B	701	416	C3-C7-C12	-4.78	116.37	122.54

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	661	NAG	C1
4	A	661	NAG	C1
4	C	661	NAG	C1
4	D	661	NAG	C1

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	605	HEM	C1A-C2A-CAA-CBA
3	D	605	HEM	C3A-C2A-CAA-CBA
4	B	661	NAG	C3-C2-N2-C7
3	A	605	HEM	C3A-C2A-CAA-CBA
5	C	703	BOG	C4-C5-C6-O6

There are no ring outliers.

7 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	605	HEM	1	0
6	A	701	416	2	0
3	A	605	HEM	2	0
6	B	701	416	2	0
6	C	701	416	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	701	416	3	0
3	C	605	HEM	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	552/587 (94%)	-0.40	1 (0%) 95 94	29, 47, 64, 79	0
1	B	552/587 (94%)	-0.41	3 (0%) 91 88	26, 46, 64, 76	0
1	C	552/587 (94%)	-0.29	6 (1%) 80 75	35, 54, 74, 88	0
1	D	552/587 (94%)	-0.33	4 (0%) 87 84	35, 53, 71, 82	0
All	All	2208/2348 (94%)	-0.36	14 (0%) 89 86	26, 50, 69, 88	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	18	ALA	4.6
1	B	59	PHE	4.2
1	D	59	PHE	3.4
1	D	354	ASN	3.3
1	D	63	ILE	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	G	3	14/15	0.66	0.49	60,62,63,64	0
2	NAG	H	3	14/15	0.67	0.45	49,52,53,54	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	E	3	14/15	0.81	0.39	51,53,53,54	0
2	NAG	F	3	14/15	0.82	0.45	52,53,56,57	0
2	NAG	H	2	14/15	0.89	0.31	44,46,47,48	0
2	NAG	E	1	14/15	0.90	0.25	33,36,38,40	0
2	NAG	F	2	14/15	0.90	0.32	42,45,46,49	0
2	NAG	E	2	14/15	0.91	0.30	34,40,43,47	0
2	NAG	G	1	14/15	0.91	0.23	37,39,42,46	0
2	NAG	G	2	14/15	0.92	0.34	49,51,53,57	0
2	NAG	H	1	14/15	0.93	0.17	32,34,37,40	0
2	NAG	F	1	14/15	0.94	0.18	27,34,36,40	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	681	14/15	0.72	0.38	54,57,58,58	0
4	NAG	A	681	14/15	0.81	0.35	49,52,53,54	0
4	NAG	A	661	14/15	0.82	0.34	31,36,39,40	0
4	NAG	D	661	14/15	0.83	0.36	34,37,38,38	0
4	NAG	B	661	14/15	0.84	0.43	36,39,41,41	0
4	NAG	C	661	14/15	0.88	0.38	35,39,42,42	0
4	NAG	D	681	14/15	0.89	0.37	43,44,46,46	0
4	NAG	B	681	14/15	0.91	0.43	44,46,47,48	0
3	HEM	A	605	43/43	0.93	0.17	24,32,39,41	0
6	416	D	701	22/22	0.94	0.18	52,55,56,58	0
6	416	A	701	22/22	0.94	0.18	49,52,53,53	0
6	416	C	701	22/22	0.94	0.16	55,59,63,63	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	HEM	D	605	43/43	0.94	0.17	32,35,44,47	0
3	HEM	C	605	43/43	0.94	0.16	29,33,38,40	0
6	416	B	701	22/22	0.95	0.17	49,52,55,57	0
3	HEM	B	605	43/43	0.95	0.15	31,32,38,41	0
5	BOG	C	703	20/20	0.95	0.14	55,56,58,58	0
5	BOG	A	704	20/20	0.96	0.13	39,41,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 416 D 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 416 A 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 416 C 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM D 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 416 B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM B 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BOG C 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BOG A 704:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.