



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 01:01 pm BST

PDB ID : 5MQ4  
Title : Crystal Structure of the leucine zipper of human PRKCBP1  
Authors : Krojer, T.; Savitsky, P.; Picaud, S.; Newman, J.; Tallant, C.; Heroven, C.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Filippakopoulos, P.  
Deposited on : 2016-12-20  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	127	 2% 79% 9% • 11%
1	B	127	 69% 17% • 13%
1	C	127	 7% 76% 17% 8%
1	D	127	 2% 69% 19% 13%
1	E	127	 4% 76% 9% • 13%
1	F	127	 2% 72% 13% • 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
2	SO4	B	1201	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein kinase C-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	113	918	573	166	168	11	0	1	0
1	B	111	911	567	161	172	11	0	0	0
1	C	117	970	601	178	180	11	0	1	0
1	D	111	929	578	166	174	11	0	0	0
1	E	111	863	536	154	163	10	0	0	0
1	F	111	929	578	166	174	11	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	997	SER	-	expression tag	UNP Q9ULU4
A	998	MET	-	expression tag	UNP Q9ULU4
B	997	SER	-	expression tag	UNP Q9ULU4
B	998	MET	-	expression tag	UNP Q9ULU4
C	-1	SER	-	expression tag	UNP Q9ULU4
C	0	MET	-	expression tag	UNP Q9ULU4
D	997	SER	-	expression tag	UNP Q9ULU4
D	998	MET	-	expression tag	UNP Q9ULU4
E	997	SER	-	expression tag	UNP Q9ULU4
E	998	MET	-	expression tag	UNP Q9ULU4
F	997	SER	-	expression tag	UNP Q9ULU4
F	998	MET	-	expression tag	UNP Q9ULU4

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	2	Total	Zn	0	0
			2	2		
3	E	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		
3	C	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		
3	F	2	Total	Zn	0	0
			2	2		


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	B	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Protein kinase C-binding protein 1




- Molecule 1: Protein kinase C-binding protein 1




- Molecule 1: Protein kinase C-binding protein 1




- Molecule 1: Protein kinase C-binding protein 1



- Molecule 1: Protein kinase C-binding protein 1



- Molecule 1: Protein kinase C-binding protein 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.64Å 237.05Å 210.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.62 – 2.70 78.62 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (78.62-2.70) 100.0 (78.62-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.208 , 0.236 0.208 , 0.236	Depositor DCC
$R_{free}$ test set	2345 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.9	Xtrriage
Anisotropy	0.152	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5558	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.47	0/934	0.60	0/1257
1	B	0.48	0/928	0.63	0/1249
1	C	0.47	0/987	0.64	0/1326
1	D	0.45	0/946	0.61	0/1269
1	E	0.51	1/876 (0.1%)	0.64	1/1180 (0.1%)
1	F	0.50	0/946	0.62	0/1269
All	All	0.48	1/5617 (0.0%)	0.62	1/7550 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	1112	CYS	CB-SG	5.49	1.91	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	1035	LEU	CA-CB-CG	6.17	129.49	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	918	0	886	7	0
1	B	911	0	867	19	0
1	C	970	0	942	16	1
1	D	929	0	904	19	0
1	E	863	0	805	11	1
1	F	929	0	904	17	0
2	A	5	0	0	0	0
2	B	5	0	0	3	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
4	B	1	0	0	0	0
All	All	5558	0	5308	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1069:VAL:HG13	1:E:1072:THR:HG21	1.63	0.79
1:D:1042:MET:HG3	1:F:1042:MET:HG3	1.66	0.75
1:C:1002:THR:HG23	1:C:1004:GLY:H	1.52	0.74
1:D:1060:GLN:HG2	1:F:1060:GLN:HG2	1.72	0.71
1:B:1049:GLU:OE2	1:B:1052:ARG:NH1	2.23	0.71
1:B:1007:ILE:HG13	1:D:1006:THR:HG22	1.72	0.70
1:B:1011:ARG:NH2	2:B:1201:SO4:S	2.75	0.58
1:C:1060:GLN:HE21	1:C:1064:GLU:HG3	1.69	0.57
1:B:1098:TYR:CG	1:B:1099:PRO:HD3	2.40	0.56
1:D:1046:LEU:HD23	1:F:1046:LEU:HD23	1.88	0.56
1:A:1007:ILE:HG23	1:F:1009:GLU:HG3	1.87	0.56
1:A:1035:LEU:HD21	1:F:1038:THR:HG22	1.89	0.55
1:D:1060:GLN:CG	1:F:1060:GLN:HG2	2.37	0.55
1:A:1077:TRP:CE3	1:D:1073:LYS:HE2	2.42	0.55
1:B:1014:ARG:NH1	1:D:1009:GLU:OE2	2.40	0.54
1:B:1073:LYS:HE2	1:F:1077:TRP:CE3	2.43	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1095:TYR:CE2	1:D:1101:GLN:HG3	2.43	0.53
1:E:1050:ARG:HG3	1:E:1051:ASP:N	2.22	0.53
1:C:1073:LYS:HE2	1:E:1077:TRP:CE3	2.45	0.52
1:C:1097:ASP:HB3	1:C:1099:PRO:HD2	1.92	0.52
1:E:1095:TYR:CE2	1:E:1101:GLN:HG3	2.44	0.52
1:B:1054:ILE:HD13	1:F:1054:ILE:HD13	1.90	0.51
1:A:1017:ILE:HD11	1:D:1017:ILE:HD13	1.92	0.51
1:C:1002:THR:HG23	1:C:1004:GLY:N	2.22	0.51
1:C:1058:LYS:HG2	1:E:1061:LEU:HD11	1.92	0.50
1:C:1077:TRP:CE3	1:E:1073:LYS:HE2	2.46	0.50
1:D:1056:GLU:O	1:D:1060:GLN:HG3	2.12	0.49
1:C:1112:CYS:SG	1:C:1113:THR:N	2.86	0.49
1:A:1019:LYS:HA	1:A:1022:TRP:CE3	2.49	0.47
1:D:1031:MET:HB2	1:F:1031:MET:HE2	1.98	0.46
1:B:1098:TYR:CD1	1:B:1099:PRO:HD3	2.51	0.46
1:F:1112:CYS:SG	1:F:1113:THR:N	2.88	0.46
1:C:1090:CYS:HB3	1:C:1113:THR:O	2.16	0.46
1:E:1088:TYR:HA	1:E:1094:SER:HA	1.98	0.46
1:D:1044:GLN:O	1:D:1048:GLN:HG3	2.17	0.45
1:C:1017:ILE:HG21	1:E:1017:ILE:HD11	1.99	0.45
1:F:1106:PRO:HA	1:F:1109:MET:HE2	1.99	0.45
1:C:1073:LYS:HE2	1:E:1077:TRP:CZ3	2.52	0.44
1:D:1112:CYS:SG	1:D:1113:THR:N	2.90	0.44
1:F:1082:LYS:HA	1:F:1082:LYS:HD3	1.82	0.43
1:B:1014:ARG:NH2	2:B:1201:SO4:O1	2.52	0.43
1:B:1077:TRP:CE3	1:F:1073:LYS:HE2	2.54	0.42
1:D:1027:GLU:O	1:D:1031:MET:HG3	2.20	0.42
1:F:1088:TYR:HA	1:F:1094:SER:HA	2.00	0.42
1:D:1093:THR:HG21	1:D:1108:HIS:CD2	2.55	0.42
1:F:1113:THR:HG22	1:F:1114:GLN:N	2.35	0.42
1:B:1112:CYS:SG	1:B:1113:THR:N	2.92	0.42
1:B:1082:LYS:HD3	1:B:1082:LYS:HA	1.62	0.42
1:B:1020:LEU:HD23	1:B:1020:LEU:HA	1.66	0.41
1:C:1037:LEU:HD23	1:C:1037:LEU:HA	1.81	0.41
1:B:1045:SER:OG	1:F:1043:ARG:NH2	2.53	0.41
1:D:1109:MET:HB3	1:D:1109:MET:HE3	1.95	0.41
1:B:1088:TYR:HA	1:B:1094:SER:HA	2.01	0.41
1:B:1109:MET:HB3	1:B:1109:MET:HE3	1.91	0.41
1:C:1077:TRP:CZ3	1:E:1073:LYS:HE2	2.55	0.41
1:C:1093:THR:HG21	1:C:1108:HIS:CE1	2.54	0.41
1:B:1077:TRP:CD1	1:F:1073:LYS:HG2	2.55	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:1061:LEU:HA	1:E:1061:LEU:HD23	1.81	0.41
1:A:1049:GLU:OE2	1:A:1052:ARG:NH2	2.44	0.40
1:A:1093:THR:HG21	1:A:1108:HIS:CE1	2.55	0.40
1:B:1035:LEU:HD22	1:D:1034:ASN:HB3	2.03	0.40
1:D:1104:HIS:CE1	1:D:1108:HIS:HB2	2.57	0.40
1:B:1011:ARG:NH2	2:B:1201:SO4:O3	2.47	0.40
1:C:1061:LEU:HA	1:C:1061:LEU:HD23	1.93	0.40
1:D:1105:TRP:CZ2	1:D:1109:MET:HG3	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1049:GLU:OE2	1:E:1050:ARG:NH2[3_655]	1.97	0.23

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/127 (88%)	111 (99%)	1 (1%)	0	100	100
1	B	109/127 (86%)	107 (98%)	2 (2%)	0	100	100
1	C	116/127 (91%)	115 (99%)	1 (1%)	0	100	100
1	D	109/127 (86%)	108 (99%)	1 (1%)	0	100	100
1	E	109/127 (86%)	109 (100%)	0	0	100	100
1	F	109/127 (86%)	108 (99%)	1 (1%)	0	100	100
All	All	664/762 (87%)	658 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/116 (84%)	93 (96%)	4 (4%)	30	59
1	B	98/116 (84%)	91 (93%)	7 (7%)	14	34
1	C	105/116 (90%)	103 (98%)	2 (2%)	57	82
1	D	102/116 (88%)	100 (98%)	2 (2%)	55	81
1	E	87/116 (75%)	84 (97%)	3 (3%)	37	66
1	F	102/116 (88%)	99 (97%)	3 (3%)	42	71
All	All	591/696 (85%)	570 (96%)	21 (4%)	35	64

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1002	THR
1	A	1012	ARG
1	A	1031	MET
1	A	1052	ARG
1	B	1005	SER
1	B	1007	ILE
1	B	1023	LEU
1	B	1037	LEU
1	B	1052	ARG
1	B	1061	LEU
1	B	1107	GLU
1	C	999	SER
1	C	1006	THR
1	D	1014	ARG
1	D	1039	MET
1	E	1023	LEU
1	E	1035	LEU
1	E	1050	ARG
1	F	1026	GLN
1	F	1043	ARG
1	F	1060	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1108	HIS
1	C	1060	GLN
1	D	1108	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 12 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	E	1201	-	4,4,4	0.26	0	6,6,6	0.56	0
2	SO4	B	1201	-	4,4,4	0.14	0	6,6,6	0.10	0
2	SO4	D	1201	-	4,4,4	0.17	0	6,6,6	0.46	0
2	SO4	A	1201	-	4,4,4	0.19	0	6,6,6	0.20	0
2	SO4	F	1201	-	4,4,4	0.31	0	6,6,6	0.79	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1201	SO4	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	113/127 (88%)	0.58	2 (1%) 68 70	44, 75, 132, 158	0
1	B	111/127 (87%)	0.36	0 100 100	34, 73, 108, 131	0
1	C	117/127 (92%)	0.62	9 (7%) 13 11	40, 73, 116, 124	0
1	D	111/127 (87%)	0.53	3 (2%) 54 55	41, 68, 94, 119	0
1	E	111/127 (87%)	0.57	5 (4%) 33 31	35, 77, 158, 179	0
1	F	111/127 (87%)	0.52	2 (1%) 68 70	39, 63, 88, 116	0
All	All	674/762 (88%)	0.53	21 (3%) 49 49	34, 71, 122, 179	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1022	TRP	4.3
1	E	1029	SER	4.0
1	D	1004	GLY	3.4
1	C	1031	MET	3.2
1	E	1028	LEU	3.1
1	C	1035	LEU	2.7
1	F	1017	ILE	2.6
1	C	1037	LEU	2.6
1	E	1037	LEU	2.5
1	C	1024	HIS	2.4
1	C	1046	LEU	2.4
1	C	1003	THR	2.2
1	F	1015	ILE	2.2
1	E	1040	ALA	2.2
1	D	1035	LEU	2.1
1	E	1009	GLU	2.1
1	D	1042	MET	2.1
1	A	1011	ARG	2.1
1	C	1000	LYS	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	1053	LEU	2.0
1	C	1042	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	1201	5/5	0.92	0.13	149,149,151,151	0
2	SO4	A	1201	5/5	0.93	0.17	108,111,121,126	0
2	SO4	D	1201	5/5	0.94	0.39	60,89,104,108	0
2	SO4	F	1201	5/5	0.94	0.33	50,88,91,107	0
2	SO4	E	1201	5/5	0.95	0.28	51,57,104,112	0
3	ZN	A	1202	1/1	0.98	0.22	62,62,62,62	0
3	ZN	A	1203	1/1	0.99	0.24	55,55,55,55	0
3	ZN	D	1203	1/1	0.99	0.20	68,68,68,68	0
3	ZN	F	1203	1/1	0.99	0.24	60,60,60,60	0
3	ZN	D	1202	1/1	0.99	0.20	59,59,59,59	0
3	ZN	F	1202	1/1	0.99	0.23	45,45,45,45	0
3	ZN	E	1202	1/1	0.99	0.21	45,45,45,45	0
3	ZN	C	1201	1/1	0.99	0.21	53,53,53,53	0
3	ZN	B	1203	1/1	0.99	0.22	60,60,60,60	0
3	ZN	B	1202	1/1	1.00	0.23	48,48,48,48	0
3	ZN	C	1202	1/1	1.00	0.20	72,72,72,72	0
3	ZN	E	1203	1/1	1.00	0.20	41,41,41,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.