

# wwPDB X-ray Structure Validation Summary Report (i)

Jan 7, 2024 – 11:53 pm GMT

PDB ID : 5MQJ

Title: Crystal structure of dCK mutant C3S

Authors: Saez-Ayala, M.; Rebuffet, E.; Hammam, K.; Gros, L.; Lopez, S.; Hajem, B.;

Humbert, M.; Baudelet, E.; Audebert, S.; Betzi, S.; Lugari, A.; Combes, S.; Pez, D.; Letard, S.; Mansfield, C.; Moussy, A.; de Sepulveda, P.; Morelli, X.;

Dubreuil, P.

Deposited on : 2016-12-20

Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$ 

Mogul : 1.8.4, CSD as541be (2020)

 $Xtriage\ (Phenix) \quad : \quad 1.13$ 

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36




# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	1049 (3.88-3.52)
Clashscore	141614	1027 (3.86-3.54)
Ramachandran outliers	138981	1069 (3.88-3.52)
Sidechain outliers	138945	1065 (3.88-3.52)
RSRZ outliers	127900	1578 (3.90-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	281	60%	22%	•	17%			
1	В	281	58%	22%	·	19%			
1	С	281	62%	19%		18%			
1	D	281	58%	22%	·	19%			



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Deoxycytidine kinase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	232	Total	С	N	О	S	0	0	0
1	A	232	1850	1184	305	354	7	0	U	0
1	В	228	Total	С	N	О	S	0	0	0
1	Ъ	220	1793	1155	291	340	7	U	U	0
1	С	230	Total	С	N	О	S	0	1	0
1		230	1811	1163	301	340	7	0	1	
1	D	229	Total	С	N	О	S	0	0	0
1	ש	429	1809	1164	296	342	7	0	U	

There are 100 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP P27707
A	-19	SER	-	expression tag	UNP P27707
A	-18	TYR	-	expression tag	UNP P27707
A	-17	TYR	-	expression tag	UNP P27707
A	-16	HIS	-	expression tag	UNP P27707
A	-15	HIS	-	expression tag	UNP P27707
A	-14	HIS	-	expression tag	UNP P27707
A	-13	HIS	-	expression tag	UNP P27707
A	-12	HIS	-	expression tag	UNP P27707
A	-11	HIS	-	expression tag	UNP P27707
A	-10	LEU	-	expression tag	UNP P27707
A	-9	GLU	-	expression tag	UNP P27707
A	-8	SER	-	expression tag	UNP P27707
A	-7	THR	-	expression tag	UNP P27707
A	-6	SER	-	expression tag	UNP P27707
A	-5	LEU	-	expression tag	UNP P27707
A	-4	TYR		expression tag	UNP P27707
A	-3	LYS	-	expression tag	UNP P27707
A	-2	LYS	-	expression tag	UNP P27707
A	-1	ALA		expression tag	UNP P27707
A	0	GLY	-	expression tag	UNP P27707

Continued on next page...



Continued from previous page...

Chain	$oxed{\mathbf{Residue}}$	Modelled	Actual	Comment	Reference
A	1	LEU	_	expression tag	UNP P27707
A	9	SER	CYS	engineered mutation	UNP P27707
A	45	SER	CYS	engineered mutation	UNP P27707
A	59	SER	CYS	engineered mutation	UNP P27707
В	-20	MET	_	initiating methionine	UNP P27707
В	-19	SER	-	expression tag	UNP P27707
В	-18	TYR	-	expression tag	UNP P27707
В	-17	TYR	-	expression tag	UNP P27707
В	-16	HIS	-	expression tag	UNP P27707
В	-15	HIS	-	expression tag	UNP P27707
В	-14	HIS	-	expression tag	UNP P27707
В	-13	HIS	-	expression tag	UNP P27707
В	-12	HIS	-	expression tag	UNP P27707
В	-11	HIS	-	expression tag	UNP P27707
В	-10	LEU	-	expression tag	UNP P27707
В	-9	GLU	-	expression tag	UNP P27707
В	-8	SER	-	expression tag	UNP P27707
В	-7	THR	-	expression tag	UNP P27707
В	-6	SER	-	expression tag	UNP P27707
В	-5	LEU	-	expression tag	UNP P27707
В	-4	TYR	-	expression tag	UNP P27707
В	-3	LYS	-	expression tag	UNP P27707
В	-2	LYS	-	expression tag	UNP P27707
В	-1	ALA	-	expression tag	UNP P27707
В	0	GLY	-	expression tag	UNP P27707
В	1	LEU	-	expression tag	UNP P27707
В	9	SER	CYS	engineered mutation	UNP P27707
В	45	SER	CYS	engineered mutation	UNP P27707
В	59	SER	CYS	engineered mutation	UNP P27707
С	-20	MET	-	initiating methionine	UNP P27707
С	-19	SER	-	expression tag	UNP P27707
С	-18	TYR	-	expression tag	UNP P27707
С	-17	TYR	-	expression tag	UNP P27707
С	-16	HIS	-	expression tag	UNP P27707
С	-15	HIS	-	expression tag	UNP P27707
С	-14	HIS	-	expression tag	UNP P27707
С	-13	HIS	-	expression tag	UNP P27707
C	-12	HIS	-	expression tag	UNP P27707
С	-11	HIS	-	expression tag	UNP P27707
C	-10	LEU	-	expression tag	UNP P27707
C	-9	GLU	-	expression tag	UNP P27707
С	-8	SER	-	expression tag	UNP P27707

Continued on next page...




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
С	-7	THR	-	expression tag	UNP P27707
С	-6	SER	-	expression tag	UNP P27707
С	-5	LEU	-	expression tag	UNP P27707
С	-4	TYR	-	expression tag	UNP P27707
С	-3	LYS	-	expression tag	UNP P27707
С	-2	LYS	-	expression tag	UNP P27707
С	-1	ALA	-	expression tag	UNP P27707
С	0	GLY	-	expression tag	UNP P27707
С	1	LEU	-	expression tag	UNP P27707
С	9	SER	CYS	engineered mutation	UNP P27707
С	45	SER	CYS	engineered mutation	UNP P27707
С	59	SER	CYS	engineered mutation	UNP P27707
D	-20	MET	_	initiating methionine	UNP P27707
D	-19	SER	-	expression tag	UNP P27707
D	-18	TYR	_	expression tag	UNP P27707
D	-17	TYR	-	expression tag	UNP P27707
D	-16	HIS	_	expression tag	UNP P27707
D	-15	HIS	-	expression tag	UNP P27707
D	-14	HIS	-	expression tag	UNP P27707
D	-13	HIS	-	expression tag	UNP P27707
D	-12	HIS	-	expression tag	UNP P27707
D	-11	HIS	-	expression tag	UNP P27707
D	-10	LEU	-	expression tag	UNP P27707
D	-9	GLU	-	expression tag	UNP P27707
D	-8	SER	-	expression tag	UNP P27707
D	-7	THR	-	expression tag	UNP P27707
D	-6	SER	-	expression tag	UNP P27707
D	-5	LEU	-	expression tag	UNP P27707
D	-4	TYR	-	expression tag	UNP P27707
D	-3	LYS	-	expression tag	UNP P27707
D	-2	LYS	-	expression tag	UNP P27707
D	-1	ALA	-	expression tag	UNP P27707
D	0	GLY	-	expression tag	UNP P27707
D	1	LEU	-	expression tag	UNP P27707
D	9	SER	CYS	engineered mutation	UNP P27707
D	45	SER	CYS	engineered mutation	UNP P27707
D	59	SER	CYS	engineered mutation	UNP P27707


 $\bullet \ \ \mathrm{Molecule} \ 2 \ \mathrm{is} \ \mathrm{URIDINE} - 5\text{'} - \mathrm{DIPHOSPHATE} \ (\mathrm{three-letter} \ \mathrm{code} \colon \ \mathrm{UDP}) \ (\mathrm{formula:} \ \ \mathrm{C}_9\mathrm{H}_{14}\mathrm{N}_2\mathrm{O}_{12}\mathrm{P}_2).$ 





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	Λ	1	Total	С	N	О	Р	0	0	
2	Λ	1	25	9	2	12	2		U	
2	B	1	Total	С	N	О	Р	0		
2	Б	1	25	9	2	12	2	0	U	
2	С	1	Total	С	N	О	Р	0	0	
2		1	25	9	2	12	2	0	U	
2	D	1	Total	С	N	О	Р	0	0	
	ש	1	25	9	2	12	2		U	

• Molecule 3 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula:  $C_9H_{13}N_2O_8P$ ).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	Λ	1	Total	С	N	О	Р	0	0
)	А	1	20	9	2	8	1	0	U
2	D	1	Total	С	N	О	Р	0	0
)	Б	1	20	9	2	8	1	0	U
3	C	1	Total	С	N	О	Р	0	0
)	C	1	20	9	2	8	1	0	U
3	D	1	Total	С	N	О	Р	0	0
)	D	1	20	9	2	8	1	0	0

 $\bullet$  Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	В	1	Total Mg 1 1	0	0
4	С	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Deoxycytidine kinase







• Molecule 1: Deoxycytidine kinase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	92.69Å 92.69Å 338.08Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.06 - 3.70	Depositor
resolution (A)	46.35 - 3.70	EDS
% Data completeness	99.9 (47.06-3.70)	Depositor
(in resolution range)	99.9 (46.35-3.70)	EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.73  (at  3.66Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
P.P.	0.181 , $0.251$	Depositor
$R, R_{free}$	0.191 , $0.254$	DCC
$R_{free}$ test set	830 reflections $(5.00\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	108.0	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.32 , 143.3	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.44, < L^2> = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7447	wwPDB-VP
Average B, all atoms $(Å^2)$	131.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, MG, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.53	0/1896	0.72	0/2584	
1	В	0.53	0/1839	0.78	0/2512	
1	С	0.52	0/1856	0.76	0/2533	
1	D	0.56	0/1855	0.75	0/2532	
All	All	0.54	0/7446	0.75	0/10161	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1850	0	1695	78	0
1	В	1793	0	1628	76	0
1	С	1811	0	1651	64	0
1	D	1809	0	1652	48	0
2	A	25	0	11	0	0
2	В	25	0	11	1	0
2	С	25	0	11	3	0
2	D	25	0	11	6	0
3	A	20	0	11	0	0

Continued on next page...



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	20	0	11	2	0
3	С	20	0	11	0	0
3	D	20	0	11	1	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
All	All	7447	0	6714	242	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:158:TRP:CZ2	1:A:162:MET:HE1	1.44	1.49
1:C:38:VAL:CG2	1:C:50:VAL:HG11	1.62	1.28
1:A:158:TRP:CE2	1:A:162:MET:CE	2.27	1.18
1:A:158:TRP:CZ2	1:A:162:MET:CE	2.28	1.15
1:C:186:LEU:HA	1:C:189:ILE:HD12	1.29	1.13

There are no symmetry-related clashes.

# 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	228/281 (81%)	209 (92%)	18 (8%)	1 (0%)	34	69
1	В	224/281 (80%)	194 (87%)	29 (13%)	1 (0%)	34	69
1	С	227/281 (81%)	203 (89%)	22 (10%)	2 (1%)	17	54
1	D	225/281 (80%)	202 (90%)	22 (10%)	1 (0%)	34	69

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured Allower		Outliers	Percentiles	
All	All	904/1124 (80%)	808 (89%)	91 (10%)	5 (1%)	25 62	

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	170	LEU
1	С	115	LYS
1	С	170	LEU
1	D	120	GLU
1	A	238	VAL

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	191/258~(74%)	185 (97%)	6 (3%)	40 65
1	В	$181/258 \ (70\%)$	174 (96%)	7 (4%)	32 60
1	С	$181/258 \ (70\%)$	169 (93%)	12 (7%)	16 48
1	D	$184/258 \ (71\%)$	167 (91%)	17 (9%)	9 36
All	All	$737/1032 \ (71\%)$	695 (94%)	42 (6%)	20 52

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	54	PRO
1	D	160	ASP
1	D	83	GLN
1	D	112	LEU
1	D	200	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:



Mol	Chain	Res	Type
1	С	108	GLN
1	D	229	GLN
1	D	97	GLN
1	В	165	GLN
1	С	97	GLN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	UDP	В	301	-	24,26,26	0.60	0	37,40,40	0.79	1 (2%)
2	UDP	A	301	-	24,26,26	0.57	1 (4%)	37,40,40	0.62	0
3	UMP	В	302	4	21,21,21	1.20	1 (4%)	31,31,31	1.68	8 (25%)
2	UDP	С	301	-	24,26,26	0.47	0	37,40,40	0.69	1 (2%)
3	UMP	D	302	4	21,21,21	1.39	3 (14%)	31,31,31	1.66	8 (25%)
3	UMP	С	302	4	21,21,21	1.19	1 (4%)	31,31,31	1.76	8 (25%)
2	UDP	D	301	4	24,26,26	0.64	1 (4%)	37,40,40	0.57	0
3	UMP	A	302	4	21,21,21	1.53	7 (33%)	31,31,31	1.60	6 (19%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	В	301	-	-	2/16/32/32	0/2/2/2
2	UDP	A	301	-	-	0/16/32/32	0/2/2/2
3	UMP	В	302	4	-	0/10/22/22	0/2/2/2
2	UDP	С	301	_	-	1/16/32/32	0/2/2/2
3	UMP	D	302	4	-	3/10/22/22	0/2/2/2
3	UMP	С	302	4	-	4/10/22/22	0/2/2/2
2	UDP	D	301	4	-	7/16/32/32	0/2/2/2
3	UMP	A	302	4	-	5/10/22/22	0/2/2/2

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$Ideal(\AA)$
2	D	301	UDP	PB-O2B	-2.88	1.43	1.54
3	D	302	UMP	O4'-C1'	2.85	1.48	1.42
3	A	302	UMP	C6-N1	2.71	1.44	1.38
3	A	302	UMP	O4'-C1'	2.64	1.48	1.42
2	A	301	UDP	PB-O2B	-2.42	1.45	1.54

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	С	302	UMP	C4-N3-C2	-4.49	120.66	126.58
3	С	302	UMP	C5-C4-N3	4.15	121.05	114.84
3	В	302	UMP	C4-N3-C2	-3.95	121.37	126.58
3	A	302	UMP	O4-C4-C5	-3.84	118.40	125.16
3	D	302	UMP	C5-C4-N3	3.83	120.57	114.84

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	301	UDP	O4'-C4'-C5'-O5'
2	D	301	UDP	PA-O3A-PB-O2B
3	A	302	UMP	C5'-O5'-P-OP1
3	A	302	UMP	C5'-O5'-P-OP2
3	A	302	UMP	C5'-O5'-P-OP3




There are no ring outliers.


5 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	301	UDP	1	0
3	В	302	UMP	2	0
2	С	301	UDP	3	0
3	D	302	UMP	1	0
2	D	301	UDP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






























# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	232/281 (82%)	-0.36	0 100 100	88, 120, 187, 234	0
1	В	228/281 (81%)	-0.31	1 (0%) 92 88	91, 127, 163, 193	0
1	С	230/281 (81%)	0.08	6 (2%) 56 43	87, 135, 194, 222	0
1	D	229/281 (81%)	-0.24	1 (0%) 92 88	89, 126, 180, 208	0
All	All	919/1124 (81%)	-0.21	8 (0%) 84 76	87, 127, 184, 234	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	119	ALA	3.8
1	С	64	THR	2.9
1	С	125	PHE	2.8
1	В	166	PHE	2.6
1	С	124	LEU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
2	UDP	В	301	25/25	0.89	0.22	119,134,144,147	0
2	UDP	D	301	25/25	0.93	0.19	123,147,160,167	0
2	UDP	С	301	25/25	0.94	0.13	128,150,166,173	0
2	UDP	A	301	25/25	0.94	0.20	115,131,151,157	0
3	UMP	В	302	20/20	0.95	0.20	124,134,148,149	0
4	MG	В	303	1/1	0.97	0.11	55,55,55,55	0
4	MG	D	303	1/1	0.97	0.13	66,66,66,66	0
3	UMP	D	302	20/20	0.98	0.20	92,110,125,128	0
4	MG	A	303	1/1	0.98	0.22	64,64,64,64	0
3	UMP	A	302	20/20	0.98	0.22	104,113,124,125	0
4	MG	С	303	1/1	0.98	0.09	91,91,91,91	0
3	UMP	С	302	20/20	0.98	0.21	113,122,144,146	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.















# Electron density around UDP A 301: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)



















# 6.5 Other polymers (i)

There are no such residues in this entry.

