

wwPDB NMR Structure Validation Summary Report (i)

Jun 3, 2023 – 06:11 PM EDT

PDB ID	:	2MUR
BMRB ID	:	25230
Title	:	Solution Structure of the Human FAAP20 UBZ-Ubiquitin Complex
Authors	:	Wang, S.; Wojtaszek, J.L.; Zhou, P.
Deposited on	:	2014-09-16

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 92%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f NMR} { m archive} \ (\#{ m Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of cha	in	
1	А	44	61%	23% •	14%
2	В	78	62%	29%	5% •



2 Ensemble composition and analysis (i)

This entry contains 10 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid						
1	A:7-A:44, B:99-B:173 (113)	0.53	7			

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 6 single-model clusters were found.

Cluster number	Models
1	6, 10
2	2, 7
Single-model clusters	1; 3; 4; 5; 8; 9



3 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1909 atoms, of which 956 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Fanconi anemia-associated protein of 20 kDa.

Mol	Chain	Residues	Atoms				Trace		
1	٨	4.4	Total	С	Н	Ν	0	S	0
	A	44	652	205	317	59	66	5	U

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q6NZ36
А	2	HIS	-	expression tag	UNP Q6NZ36
А	3	MET	-	expression tag	UNP Q6NZ36

• Molecule 2 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms				Trace		
2	В	78	Total 1256	C 387	Н 639	N 109	0 120	S 1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	99	SER	-	expression tag	UNP P62987
В	100	HIS	-	expression tag	UNP P62987

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms
9	Δ	1	Total Zn
J	A	1	1 1




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Fanconi anemia-associated protein of 20 kDa



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

 \bullet Molecule 1: Fanconi anemia-associated protein of 20 kDa





5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: torsion angle dynamics.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1504
Number of shifts mapped to atoms	1504
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	92%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	297	283	283	7 ± 1
2	В	598	620	619	23 ± 3
All	All	8960	9030	9020	251

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

5 of 101 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Mod	dels
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
2:B:123:ILE:HG23	2:B:143:LEU:HD12	0.90	1.39	7	7
1:A:8:LEU:HD11	1:A:31:LEU:HD11	0.88	1.42	1	7
1:A:25:LEU:HD12	2:B:166:THR:HG21	0.83	1.48	3	1
2:B:136:ILE:HD13	2:B:171:LEU:HD11	0.80	1.53	2	6
2:B:105:VAL:HG21	2:B:130:ILE:HD11	0.80	1.53	9	6



6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	37/44~(84%)	$33 \pm 1 (90 \pm 2\%)$	$4\pm1~(10\pm2\%)$	0±0 (0±0%)	100	100
2	В	74/78~(95%)	$69 \pm 1 (94 \pm 2\%)$	$4\pm1~(6\pm2\%)$	0±0 (1±1%)	32	76
All	All	1110/1220~(91%)	1025~(92%)	81 (7%)	4 (0%)	38	78

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
2	В	121	ASP	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perc	entiles
1	А	34/37~(92%)	27 ± 1 (79 $\pm 4\%$)	$7\pm1~(21\pm4\%)$	3	32
2	В	69/70~(99%)	60 ± 1 (87 $\pm2\%$)	$9\pm1~(13\pm2\%)$	7	48
All	All	1030/1070~(96%)	869 (84%)	161 (16%)	5	43

 $5~{\rm of}~43$ unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	8	LEU	10
1	А	22	LEU	10
2	В	125	ASN	10
2	В	129	LYS	10
2	В	136	ILE	10



6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 92% for the well-defined parts and 91% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1504
Number of shifts mapped to atoms	1504
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction}\pm{\rm precision},ppm$	Suggested action
$^{13}C_{\alpha}$	118	0.21 ± 0.08	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	112	0.52 ± 0.16	Should be applied
$^{13}C'$	109	-0.14 ± 0.13	None needed (< 0.5 ppm)
¹⁵ N	110	0.43 ± 0.16	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 92%, i.e. 1439 atoms were assigned a chemical shift out of a possible 1560. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	536/559~(96%)	220/225~(98%)	213/226~(94%)	103/108~(95%)
Sidechain	845/929~(91%)	575/602~(96%)	259/293~(88%)	11/34~(32%)

Continued on next page...



	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Aromatic	58/72~(81%)	29/37~(78%)	28/31~(90%)	1/4~(25%)
Overall	1439/1560~(92%)	824/864 (95%)	500/550~(91%)	115/146~(79%)

Continued from previous page...


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)


The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:







8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	3650
Intra-residue (i-j =0)	366
Sequential (i-j =1)	813
Medium range ($ i-j >1$ and $ i-j <5$)	786
Long range $(i-j \ge 5)$	1451
Inter-chain	144
Hydrogen bond restraints	90
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	29.9
Number of long range restraints per residue ¹	12.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	2.2	0.2
0.2-0.5 (Medium)	0.2	0.21
>0.5 (Large)	1.0	0.98



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destruction to the second	Count	ount $\%^1$	Vio	lated	3	Consis	tently	$\sqrt{Violated^4}$
Restraints type	Count	701	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	366	10.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	301	8.2	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	65	1.8	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	813	22.3	4	0.5	0.1	0	0.0	0.0
Backbone-Backbone	184	5.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	519	14.2	4	0.8	0.1	0	0.0	0.0
Sidechain-Sidechain	110	3.0	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	786	21.5	4	0.5	0.1	0	0.0	0.0
Backbone-Backbone	179	4.9	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	385	10.5	1	0.3	0.0	0	0.0	0.0
Sidechain-Sidechain	222	6.1	3	1.4	0.1	0	0.0	0.0
Long range $(i-j \ge 5)$	1451	39.8	2	0.1	0.1	0	0.0	0.0
Backbone-Backbone	138	3.8	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	659	18.1	1	0.2	0.0	0	0.0	0.0
Sidechain-Sidechain	654	17.9	1	0.2	0.0	0	0.0	0.0
Inter-chain	144	3.9	3	2.1	0.1	0	0.0	0.0
Backbone-Backbone	5	0.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	65	1.8	1	1.5	0.0	0	0.0	0.0
Sidechain-Sidechain	74	2.0	2	2.7	0.1	0	0.0	0.0
Hydrogen bond	90	2.5	1	1.1	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	3650	100.0	14	0.4	0.4	0	0.0	0.0
Backbone-Backbone	592	16.2	1	0.2	0.0	0	0.0	0.0
Backbone-Sidechain	1933	53.0	7	0.4	0.2	0	0.0	0.0
Sidechain-Sidechain	1125	30.8	6	0.5	0.2	0	0.0	0.0

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)


The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	5	Mean (Å)	Max (Å)	SD^6 (Å)	Median (Å)
Model ID	IR^1	SQ^2	MR^3	LR ⁴	IC ⁵	Total	Mean (A)	Max (A)	$\left \mathbf{SD} \right (\mathbf{A})$	Median (A)
1	0	1	1	0	2	4	0.27	0.64	0.22	0.16
2	0	1	0	0	1	2	0.14	0.15	0.01	0.14
3	0	2	0	1	1	4	0.33	0.85	0.3	0.18
4	0	1	1	1	0	3	0.18	0.21	0.04	0.2
5	0	0	0	0	2	2	0.9	0.98	0.08	0.9
6	0	1	3	0	1	5	0.26	0.74	0.24	0.15
7	0	1	0	0	3	4	0.41	0.76	0.28	0.38
8	0	1	1	0	0	2	0.16	0.16	0.01	0.16
9	0	0	1	0	3	4	0.39	0.64	0.23	0.4
10	0	0	1	1	2	4	0.29	0.8	0.29	0.12

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,



⁵Inter-chain restraints, ⁶Standard deviation



9.2.1 Bar graph : Distance Violation statistics for each model (i)

The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 3547(IR:366, SQ:809, MR:782, LR:1449, IC:141) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	Fraction	n of the ensemble		
IR^1	SQ^2	MR^3	LR^4	IC ⁵	Total	Count^6	%
0	1	2	1	0	4	1	10.0
0	2	2	1	0	5	2	20.0
0	1	0	0	0	1	3	30.0
0	0	0	0	2	2	4	40.0


Continued on next page...



	Continueu from preotous page								
Nu	Number of violated restraints						n of the ensemble		
IR^1	SQ^2	MR^3	LR ⁴	IC ⁵	Total	Count^6	%		
0	0	0	0	0	0	5	50.0		
0	0	0	0	0	0	6	60.0		
0	0	0	0	1	1	7	70.0		
0	0	0	0	0	0	8	80.0		
0	0	0	0	0	0	9	90.0		
0	0	0	0	0	0	10	100.0		

Continued from previous page...

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations




9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH1	7	0.7	0.24	0.76
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH2	7	0.7	0.24	0.76
(2,899)	1:A:41:ASP:OD2	2:B:172:ARG:NH1	4	0.68	0.09	0.64
(1,817)	1:A:44:TRP:HE3	2:B:148:LYS:HA	4	0.14	0.04	0.12
(1,370)	1:A:21:ARG:HD2	1:A:22:LEU:H	3	0.14	0.01	0.14
(1,370)	1:A:21:ARG:HD3	1:A:22:LEU:H	3	0.14	0.01	0.14
(1,1798)	2:B:128:ALA:H	2:B:129:LYS:HG2	2	0.18	0.02	0.18
(1,1798)	2:B:128:ALA:H	2:B:129:LYS:HG3	2	0.18	0.02	0.18
(2,145)	1:A:29:SER:H	1:A:31:LEU:HG	2	0.18	0.02	0.18
(1,1409)	2:B:116:GLU:H	2:B:129:LYS:HG2	2	0.16	0.04	0.16
(1,1409)	2:B:116:GLU:H	2:B:129:LYS:HG3	2	0.16	0.04	0.16
(1,231)	1:A:14:CYS:HA	1:A:15:GLN:HG2	2	0.15	0.0	0.15
(1,231)	1:A:14:CYS:HA	1:A:15:GLN:HG3	2	0.15	0.0	0.15
(3,39)	2:B:123:ILE:O	2:B:127:LYS:H	2	0.12	0.01	0.12
(1,399)	1:A:22:LEU:HG	1:A:26:ASP:HB2	2	0.12	0.01	0.12




¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH1	5	0.98
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH2	5	0.98
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH1	3	0.85
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH2	3	0.85
(2,899)	1:A:41:ASP:OD2	2:B:172:ARG:NH1	5	0.83
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH1	10	0.8
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH2	10	0.8
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH1	7	0.76

Continued on next page...



Key	Atom-1	Atom-2	Model ID	Violation (Å)
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH2	7	0.76
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH1	6	0.74
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH2	6	0.74
(2,899)	1:A:41:ASP:OD2	2:B:172:ARG:NH1	1	0.64
(2,899)	1:A:41:ASP:OD2	2:B:172:ARG:NH1	9	0.64
(2,899)	1:A:41:ASP:OD2	2:B:172:ARG:NH1	7	0.62
(3,91)	1:A:41:ASP:OD1	2:B:142:ARG:NH1	9	0.59

Continued from previous page...



10 Dihedral-angle violation analysis (i)

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value

