



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 23, 2024 – 01:53 AM EDT

PDB ID : 5MV3  
Title : ACC1 Fab fragment in complex with CII583-591 (CG10)  
Authors : Dobritsch, D.; Holmdahl, R.; Ge, C.  
Deposited on : 2017-01-15  
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	89% 9% ..
1	C	218	90% 7% .
1	F	218	91% 8% .
1	I	218	91% 8% .
1	O	218	93% 6% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	R	218	 6% 91% 7%
1	U	218	 6% 88% 8%
2	B	218	 92% 7%
2	D	218	 92% 8%
2	G	218	 92% 8%
2	J	218	 91% 9%
2	M	218	 91% 9%
2	P	218	 94% 5%
2	S	218	 89% 11%
2	V	218	 91% 9%
3	E	30	 37% 7% 57%
3	H	30	 37% 60%
3	K	30	 40% 57%
3	N	30	 3% 40% 57%
3	Q	30	 30% 7% 60%
3	T	30	 37% 60%
3	W	30	 33% 7% 60%
3	X	30	 33% 13% 53%
4	L	220	 94% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	HYP	Q	12	-	-	-	X
3	HYP	T	12	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 26921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called heavy chain of ACC1 Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	216	Total 1629	C 1028	N 271	O 322	S 8	0	0	0
1	C	213	Total 1614	C 1020	N 268	O 318	S 8	0	0	0
1	F	216	Total 1629	C 1028	N 271	O 322	S 8	0	0	0
1	I	215	Total 1625	C 1026	N 270	O 321	S 8	0	0	0
1	O	215	Total 1625	C 1026	N 270	O 321	S 8	0	0	0
1	R	214	Total 1613	C 1019	N 269	O 317	S 8	0	0	0
1	U	211	Total 1597	C 1008	N 266	O 315	S 8	0	0	0

- Molecule 2 is a protein called light chain of ACC1 Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	218	Total 1666	C 1038	N 282	O 339	S 7	0	0	0
2	D	217	Total 1660	C 1035	N 281	O 338	S 6	0	0	0
2	G	217	Total 1660	C 1035	N 281	O 338	S 6	0	0	0
2	J	218	Total 1666	C 1038	N 282	O 339	S 7	0	0	0
2	M	218	Total 1666	C 1038	N 282	O 339	S 7	0	0	0
2	P	215	Total 1643	C 1026	N 278	O 333	S 6	0	0	0
2	S	218	Total 1666	C 1038	N 282	O 339	S 7	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	V	218	1666	1038	282	339	7	0	0	0

- Molecule 3 is a protein called synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	X	14	87	54	17	16	0	0	0
3	E	13	84	52	16	16	0	0	0
3	H	12	76	47	15	14	0	0	0
3	K	13	83	52	16	15	0	0	0
3	N	13	83	52	16	15	0	0	0
3	Q	12	76	47	15	14	0	0	0
3	T	12	76	47	15	14	0	0	0
3	W	12	76	47	15	14	0	0	0

- Molecule 4 is a protein called heavy chain of ACC1 Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	L	220	1655	1044	275	328	8	0	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: heavy chain of ACC1 Fab fragment

Chain A: 




- Molecule 1: heavy chain of ACC1 Fab fragment

Chain C: 




- Molecule 1: heavy chain of ACC1 Fab fragment

Chain F: 




- Molecule 1: heavy chain of ACC1 Fab fragment

Chain I: 




- Molecule 1: heavy chain of ACC1 Fab fragment

Chain O: 





- Molecule 1: heavy chain of ACC1 Fab fragment


Chain R: 



- Molecule 1: heavy chain of ACC1 Fab fragment



- Molecule 2: light chain of ACC1 Fab fragment




- Molecule 2: light chain of ACC1 Fab fragment



- Molecule 2: light chain of ACC1 Fab fragment



- Molecule 2: light chain of ACC1 Fab fragment



- Molecule 2: light chain of ACC1 Fab fragment





- Molecule 2: light chain of ACC1 Fab fragment

Chain P:  94% 5%





- Molecule 2: light chain of ACC1 Fab fragment

Chain S:  2% 89% 11%




- Molecule 2: light chain of ACC1 Fab fragment

Chain V:  5% 91% 9%





- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,

Chain X:  33% 13% 53%




- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,

Chain E:  37% 7% 57%



- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,


Chain H:  37% 7% 60%



- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,




Chain K: 



- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,

Chain N: 




- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,

Chain Q: 




- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,

Chain T: 



- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II, Collagen alpha-1(II) chain, synthetic peptide containing the CII583-591 epitope of collagen type II,

Chain W: 



- Molecule 4: heavy chain of ACC1 Fab fragment

Chain L: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.35Å 92.84Å 121.71Å 92.50° 106.53° 104.31°	Depositor
Resolution (Å)	50.01 – 2.95 48.85 – 2.95	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.01-2.95) 99.0 (48.85-2.95)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 2.96Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.202 , 0.261 0.204 , 0.262	Depositor DCC
$R_{free}$ test set	3639 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.4	Xtrriage
Anisotropy	0.301	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25 , 20.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	26921	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/1669	0.70	1/2277 (0.0%)
1	C	0.51	0/1654	0.73	0/2257
1	F	0.53	0/1669	0.71	1/2277 (0.0%)
1	I	0.53	0/1665	0.74	0/2272
1	O	0.50	0/1665	0.69	0/2272
1	R	0.56	1/1653 (0.1%)	0.71	1/2255 (0.0%)
1	U	0.48	0/1635	0.69	0/2227
2	B	0.52	0/1703	0.73	1/2310 (0.0%)
2	D	0.57	1/1697 (0.1%)	0.73	0/2302
2	G	0.56	2/1697 (0.1%)	0.72	0/2302
2	J	0.49	0/1703	0.70	0/2310
2	M	0.52	0/1703	0.72	0/2310
2	P	0.50	0/1680	0.71	0/2279
2	S	0.53	0/1703	0.71	0/2310
2	V	0.48	0/1703	0.69	0/2310
3	E	0.46	0/69	0.81	0/92
3	H	0.57	0/69	0.73	0/92
3	K	0.54	0/76	0.83	0/100
3	N	0.63	0/76	0.76	0/100
3	Q	0.65	0/69	0.79	0/92
3	T	0.54	0/69	0.73	0/92
3	W	0.51	0/69	0.87	0/92
3	X	0.59	0/80	0.65	0/106
4	L	0.52	0/1696	0.74	0/2316
All	All	0.52	4/27472 (0.0%)	0.71	4/37352 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	I	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	127	GLU	CG-CD	9.28	1.65	1.51
2	G	127	GLU	CD-OE2	8.58	1.35	1.25
1	R	218	PRO	C-O	5.73	1.34	1.23
2	G	127	GLU	CG-CD	5.67	1.60	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	27	GLU	OE1-CD-OE2	-7.65	114.12	123.30
1	A	69	ARG	NE-CZ-NH1	5.33	122.96	120.30
1	F	69	ARG	NE-CZ-NH2	5.12	122.86	120.30
1	R	210	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	I	152	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1629	0	1589	11	1
1	C	1614	0	1577	8	0
1	F	1629	0	1590	10	0
1	I	1625	0	1586	7	0
1	O	1625	0	1587	6	0
1	R	1613	0	1573	7	0
1	U	1597	0	1555	13	0
2	B	1666	0	1597	5	0
2	D	1660	0	1593	8	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	G	1660	0	1593	9	0
2	J	1666	0	1597	9	0
2	M	1666	0	1597	8	0
2	P	1643	0	1581	4	0
2	S	1666	0	1597	13	0
2	V	1666	0	1597	10	1
3	E	84	0	86	0	0
3	H	76	0	79	0	0
3	K	83	0	84	0	0
3	N	83	0	85	0	0
3	Q	76	0	79	2	0
3	T	76	0	79	0	0
3	W	76	0	78	1	0
3	X	87	0	87	3	0
4	L	1655	0	1618	8	0
All	All	26921	0	26084	127	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 127 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:U:174:LEU:HD13	2:V:164:LEU:HD11	1.48	0.96
1:F:12:VAL:HG11	1:F:88:LEU:HD13	1.73	0.69
2:S:13:VAL:HG21	2:S:19:ALA:HB2	1.77	0.67
2:V:11:LEU:HD12	2:V:13:VAL:HG23	1.76	0.67
2:G:46:GLN:HG3	2:G:47:PRO:HD2	1.77	0.66

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:THR:OG1	2:V:10:SER:OG[1_665]	2.14	0.06

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/218 (97%)	202 (95%)	9 (4%)	1 (0%)	29	64
1	C	209/218 (96%)	205 (98%)	3 (1%)	1 (0%)	29	64
1	F	212/218 (97%)	205 (97%)	7 (3%)	0	100	100
1	I	211/218 (97%)	202 (96%)	8 (4%)	1 (0%)	29	64
1	O	211/218 (97%)	203 (96%)	8 (4%)	0	100	100
1	R	210/218 (96%)	203 (97%)	7 (3%)	0	100	100
1	U	205/218 (94%)	194 (95%)	10 (5%)	1 (0%)	29	64
2	B	216/218 (99%)	208 (96%)	8 (4%)	0	100	100
2	D	215/218 (99%)	209 (97%)	6 (3%)	0	100	100
2	G	215/218 (99%)	206 (96%)	9 (4%)	0	100	100
2	J	216/218 (99%)	206 (95%)	10 (5%)	0	100	100
2	M	216/218 (99%)	212 (98%)	4 (2%)	0	100	100
2	P	213/218 (98%)	206 (97%)	7 (3%)	0	100	100
2	S	216/218 (99%)	205 (95%)	10 (5%)	1 (0%)	29	64
2	V	216/218 (99%)	209 (97%)	7 (3%)	0	100	100
3	E	11/30 (37%)	10 (91%)	1 (9%)	0	100	100
3	H	10/30 (33%)	9 (90%)	1 (10%)	0	100	100
3	K	10/30 (33%)	10 (100%)	0	0	100	100
3	N	10/30 (33%)	10 (100%)	0	0	100	100
3	Q	10/30 (33%)	10 (100%)	0	0	100	100
3	T	10/30 (33%)	10 (100%)	0	0	100	100
3	W	10/30 (33%)	10 (100%)	0	0	100	100
3	X	11/30 (37%)	10 (91%)	1 (9%)	0	100	100
4	L	218/220 (99%)	212 (97%)	5 (2%)	1 (0%)	29	64
All	All	3493/3730 (94%)	3366 (96%)	121 (4%)	6 (0%)	47	79

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	L	102	THR
1	A	102	THR
1	I	102	THR
1	U	102	THR
2	S	124	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/185 (100%)	181 (98%)	3 (2%)	62	84
1	C	183/185 (99%)	179 (98%)	4 (2%)	52	79
1	F	184/185 (100%)	181 (98%)	3 (2%)	62	84
1	I	184/185 (100%)	181 (98%)	3 (2%)	62	84
1	O	184/185 (100%)	182 (99%)	2 (1%)	73	89
1	R	182/185 (98%)	180 (99%)	2 (1%)	73	89
1	U	180/185 (97%)	178 (99%)	2 (1%)	73	89
2	B	188/188 (100%)	181 (96%)	7 (4%)	34	66
2	D	187/188 (100%)	184 (98%)	3 (2%)	62	84
2	G	187/188 (100%)	183 (98%)	4 (2%)	53	80
2	J	188/188 (100%)	183 (97%)	5 (3%)	44	74
2	M	188/188 (100%)	183 (97%)	5 (3%)	44	74
2	P	185/188 (98%)	181 (98%)	4 (2%)	52	79
2	S	188/188 (100%)	185 (98%)	3 (2%)	62	84
2	V	188/188 (100%)	180 (96%)	8 (4%)	29	62
3	E	6/12 (50%)	6 (100%)	0	100	100
3	H	6/12 (50%)	6 (100%)	0	100	100
3	K	7/12 (58%)	7 (100%)	0	100	100
3	N	7/12 (58%)	7 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	Q	6/12 (50%)	6 (100%)	0	100	100
3	T	6/12 (50%)	6 (100%)	0	100	100
3	W	6/12 (50%)	6 (100%)	0	100	100
3	X	7/12 (58%)	7 (100%)	0	100	100
4	L	187/187 (100%)	185 (99%)	2 (1%)	73	89
All	All	3018/3082 (98%)	2958 (98%)	60 (2%)	55	80

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	94	GLN
2	V	111	LYS
2	M	50	PHE
2	V	95	SER
2	V	189	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
2	V	161	ASN
2	S	128	GLN
2	G	31	ASN
2	D	149	ASN
4	L	167	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HYP	H	12	3	6,8,9	0.74	0	5,10,12	1.92	2 (40%)
3	HYP	T	12	3	6,8,9	0.70	0	5,10,12	2.06	2 (40%)
3	HYP	K	12	3	6,8,9	0.64	0	5,10,12	1.93	2 (40%)
3	HYP	Q	12	3	6,8,9	0.71	0	5,10,12	2.05	2 (40%)
3	HYP	E	24	3	6,8,9	0.44	0	5,10,12	1.73	2 (40%)
3	HYP	E	12	3	6,8,9	0.76	0	5,10,12	2.03	2 (40%)
3	HYP	X	12	3	6,8,9	1.16	1 (16%)	5,10,12	1.42	1 (20%)
3	HYP	N	12	3	6,8,9	0.56	0	5,10,12	1.97	2 (40%)
3	HYP	W	12	3	6,8,9	0.57	0	5,10,12	1.84	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HYP	H	12	3	-	0/0/11/13	0/1/1/1
3	HYP	T	12	3	-	0/0/11/13	0/1/1/1
3	HYP	K	12	3	-	0/0/11/13	0/1/1/1
3	HYP	Q	12	3	-	0/0/11/13	0/1/1/1
3	HYP	E	24	3	-	0/0/11/13	0/1/1/1
3	HYP	E	12	3	-	0/0/11/13	0/1/1/1
3	HYP	X	12	3	-	0/0/11/13	0/1/1/1
3	HYP	N	12	3	-	0/0/11/13	0/1/1/1
3	HYP	W	12	3	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	12	HYP	CD-N	2.13	1.54	1.47

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	K	12	HYP	CB-CG-CD	3.73	107.84	103.27
3	W	12	HYP	CB-CG-CD	3.46	107.51	103.27
3	Q	12	HYP	CB-CG-CD	3.05	107.00	103.27

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	12	HYP	CB-CG-CD	3.03	106.98	103.27
3	E	12	HYP	CB-CG-CD	2.97	106.91	103.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Q	12	HYP	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/218 (99%)	-0.37	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	39, 70, 107, 141	0
1	C	213/218 (97%)	-0.17	2 (0%) <span style="border: 1px solid blue; padding: 2px;">84</span> <span style="border: 1px solid blue; padding: 2px;">71</span>	39, 80, 117, 140	0
1	F	216/218 (99%)	-0.39	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	48, 71, 106, 121	0
1	I	215/218 (98%)	-0.50	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	46, 64, 87, 106	0
1	O	215/218 (98%)	-0.28	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	46, 79, 110, 132	0
1	R	214/218 (98%)	0.34	12 (5%) <span style="border: 1px solid red; padding: 2px;">24</span> <span style="border: 1px solid red; padding: 2px;">15</span>	50, 103, 136, 162	0
1	U	211/218 (96%)	0.37	14 (6%) <span style="border: 1px solid red; padding: 2px;">18</span> <span style="border: 1px solid red; padding: 2px;">10</span>	65, 116, 166, 190	0
2	B	218/218 (100%)	-0.28	1 (0%) <span style="border: 1px solid blue; padding: 2px;">91</span> <span style="border: 1px solid blue; padding: 2px;">81</span>	33, 72, 124, 144	0
2	D	217/218 (99%)	-0.37	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	35, 61, 108, 123	0
2	G	217/218 (99%)	-0.38	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	44, 66, 106, 125	0
2	J	218/218 (100%)	-0.34	2 (0%) <span style="border: 1px solid blue; padding: 2px;">84</span> <span style="border: 1px solid blue; padding: 2px;">71</span>	41, 66, 98, 117	0
2	M	218/218 (100%)	-0.35	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	44, 65, 107, 125	0
2	P	215/218 (98%)	-0.35	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	46, 74, 102, 114	0
2	S	218/218 (100%)	-0.19	4 (1%) <span style="border: 1px solid blue; padding: 2px;">68</span> <span style="border: 1px solid blue; padding: 2px;">51</span>	38, 75, 130, 145	0
2	V	218/218 (100%)	0.05	10 (4%) <span style="border: 1px solid red; padding: 2px;">32</span> <span style="border: 1px solid red; padding: 2px;">20</span>	45, 86, 153, 174	0
3	E	11/30 (36%)	-0.32	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	46, 60, 100, 112	0
3	H	11/30 (36%)	-0.43	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	66, 73, 101, 113	0
3	K	12/30 (40%)	-0.28	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	67, 71, 108, 115	0
3	N	12/30 (40%)	0.39	1 (8%) <span style="border: 1px solid red; padding: 2px;">11</span> <span style="border: 1px solid red; padding: 2px;">6</span>	52, 65, 104, 106	0
3	Q	11/30 (36%)	-0.27	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	56, 64, 89, 94	0
3	T	11/30 (36%)	-0.17	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	60, 73, 106, 125	0
3	W	11/30 (36%)	-0.06	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	74, 84, 116, 127	0
3	X	13/30 (43%)	-0.10	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	59, 71, 94, 102	0
4	L	220/220 (100%)	-0.38	1 (0%) <span style="border: 1px solid blue; padding: 2px;">91</span> <span style="border: 1px solid blue; padding: 2px;">81</span>	45, 69, 98, 117	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
All	All	3551/3730 (95%)	-0.22	47 (1%) 77 61	33, 73, 128, 190	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	N	23	PRO	4.9
1	R	200	VAL	4.8
1	U	17	SER	4.2
1	R	144	LEU	4.1
1	U	125	TYR	4.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	HYP	Q	12	8/9	0.70	0.45	102,107,111,111	0
3	HYP	T	12	8/9	0.73	0.54	97,105,114,115	0
3	HYP	E	24	8/9	0.85	0.19	95,100,104,107	0
3	HYP	W	12	8/9	0.86	0.40	104,107,111,111	0
3	HYP	K	12	8/9	0.87	0.17	92,96,102,102	0
3	HYP	H	12	8/9	0.87	0.27	89,96,100,102	0
3	HYP	X	12	8/9	0.89	0.16	72,76,82,84	0
3	HYP	N	12	8/9	0.90	0.23	89,95,99,99	0
3	HYP	E	12	8/9	0.90	0.21	88,96,101,109	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.