

# wwPDB X-ray Structure Validation Summary Report (i)

May 15, 2020 – 06:48 pm BST

PDB ID : 1MY2

Title: crystal titration experiment (AMPA complex control)

Authors : Jin, R.; Gouaux, E.

Deposited on : 2002-10-03

Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{resolution range}( ext{Å}))$
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	A	263	83%	13%	• • •
1	В	263	79%	17%	
1	С	263	80%	15%	



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GLUTAMATE RECEPTOR 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	1 1 000	258	Total	С	N	О	S	0	0	0
1	A	250	2018	1286	336	382	14	U		U
1	В	258	Total	С	N	О	S	0	0	0
1		250	2018	1286	336	382	14			
1	1 C	258	Total	С	N	О	S	0	0	0
1			2013	1282	335	382	14	U	0	

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	=	CLONING ARTIFACT	UNP P19491
A	2	ALA	-	CLONING ARTIFACT	UNP P19491
A	118	GLY	-	LINKER	UNP P19491
A	119	THR	_	LINKER	UNP P19491
В	1	GLY	-	CLONING ARTIFACT	UNP P19491
В	2	ALA	-	CLONING ARTIFACT	UNP P19491
В	118	GLY	-	LINKER	UNP P19491
В	119	THR	-	LINKER	UNP P19491
С	1	GLY	_	CLONING ARTIFACT	UNP P19491
С	2	ALA	-	CLONING ARTIFACT	UNP P19491
С	118	GLY	-	LINKER	UNP P19491
С	119	THR	-	LINKER	UNP P19491

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	2	$\begin{array}{c c} Total & Zn \\ 2 & 2 \end{array}$	0	0
2	A	1	Total Zn 1 1	0	0
2	С	2	Total Zn 2 2	0	0



• Molecule 3 is (S)-ALPHA-AMINO-3-HYDROXY-5-METHYL-4-ISOXAZOLEPROPIONIC ACID (three-letter code: AMQ) (formula:  $C_7H_{10}N_2O_4$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 13 7 2 4	0	0
3	В	1	Total C N O 13 7 2 4	0	0
3	С	1	Total C N O 13 7 2 4	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	138	Total O 138 138	0	0
4	В	164	Total O 164 164	0	0
4	С	118	Total O 118 118	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: GLUTAMATE RECEPTOR 2





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	114.54Å 164.10Å 47.40Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.73 - 1.80	Depositor
% Data completeness	94.2 (29.73-1.80)	Depositor
(in resolution range)	34.2 (23.13 1.00)	Берозгот
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
$R, R_{free}$	0.225 , $0.244$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6513	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP



## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: AMQ, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z  > 5
1	A	0.33	0/2054	0.57	0/2762
1	В	0.32	0/2054	0.58	0/2762
1	С	0.33	0/2049	0.57	0/2756
All	All	0.33	0/6157	0.57	0/8280

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2018	0	2050	30	0
1	В	2018	0	2050	39	0
1	С	2013	0	2038	49	0
2	A	1	0	0	0	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
3	A	13	0	8	0	0
3	В	13	0	8	1	0
3	С	13	0	8	0	0
4	A	138	0	0	1	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	H(added)	Clashes	Symm-Clashes
4	В	164	0	0	4	0
4	С	118	0	0	3	0
All	All	6513	0	6162	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:C:130:GLN:HE22	1:C:133:ILE:H	1.00	0.94
1:B:157:LYS:HE3	1:B:157:LYS:HA	1.55	0.88
1:C:116:LYS:O	1:C:119:THR:HG23	1.80	0.81
1:C:130:GLN:NE2	1:C:133:ILE:H	1.77	0.81
1:A:151:LYS:HA	1:A:151:LYS:HE2	1.65	0.76

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	$256/263 \ (97\%)$	254 (99%)	2 (1%)	0	100	100
1	В	256/263 (97%)	250 (98%)	5 (2%)	1 (0%)	34	21
1	С	256/263 (97%)	249 (97%)	6 (2%)	1 (0%)	34	21
All	All	768/789 (97%)	753 (98%)	13 (2%)	2 (0%)	41	27

#### All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	259	GLY

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	C	153	ALA

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	217/219 (99%)	209 (96%)	8 (4%)	34 19		
1	В	217/219 (99%)	208 (96%)	9 (4%)	30 16		
1	С	216/219 (99%)	206 (95%)	10 (5%)	27 13		
All	All	650/657 (99%)	623 (96%)	27 (4%)	30 15		

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	${f Res}$	$\mathbf{Type}$
1	В	119	THR
1	В	172	ARG
1	С	151	LYS
1	В	139	ASP
1	A	151	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	GLN
1	С	130	GLN
1	С	252	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	B	ond leng	$_{ m gths}$	E	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	AMQ	С	601	-	5,13,13	0.86	0	3,18,18	1.42	1 (33%)
3	AMQ	A	602	-	5,13,13	0.99	0	3,18,18	1.39	1 (33%)
3	AMQ	В	603	-	5,13,13	0.85	0	3,18,18	1.44	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMQ	С	601	_	-	0/2/8/8	0/1/1/1
3	AMQ	A	602	_	-	0/2/8/8	0/1/1/1
3	AMQ	В	603	-	-	0/2/8/8	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
3	В	603	AMQ	CE2-CD2-CG	2.29	134.29	126.51
3	С	601	AMQ	CE2-CD2-CG	2.29	134.28	126.51
3	A	602	AMQ	CE2-CD2-CG	2.23	134.08	126.51



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	603	AMQ	1	0

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

#### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

