



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 12, 2023 – 02:54 PM EDT

PDB ID : 4MZ9
Title : Revised structure of E. coli SSB
Authors : Oakley, A.J.
Deposited on : 2013-09-29
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	
1	B	178	
1	C	178	
1	D	178	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3691 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Single-stranded DNA-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	114	894	555	161	173	5	0	2	0
1	B	114	884	548	161	171	4	0	0	0
1	C	114	884	548	161	171	4	0	0	0
1	D	114	884	548	161	171	4	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	30	Total	O	0	0
			30	30		
2	B	45	Total	O	0	0
			45	45		
2	C	42	Total	O	0	0
			42	42		
2	D	28	Total	O	0	0
			28	28		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Single-stranded DNA-binding protein



- Molecule 1: Single-stranded DNA-binding protein




- Molecule 1: Single-stranded DNA-binding protein




ASP
PHE
ASP
ASP
ASP
ILE
PRO
PHE

● Molecule 1: Single-stranded DNA-binding protein

Chain D: 


 MET A1 S2 R3 G4 L14 V20 M23 V29 A30 N31 I32 T33 L34 A35 T36 S37 E38 S39 W40 R41 D42 K43 A44 T45 G46 E47 M48 K49 T52 E53 W54 H55 V58 L59 K62 L63 S68 L71 R72 Q76 V77 Y78 I79 E80 R84 D90 Q91


 S92 G93 Q94 D95 P99 V103 M104 V105 T108 M111 L112 G113 G114 ARG GLN GLY GLY GLY ALA ALA PRO ALA GLY GLY ASN ILE GLY GLY GLN PRO GLN GLY GLY TRP GLY GLN PRO GLN PRO GLN GLY GLY ASN PHE SER GLY GLY ALA ALA GLN SER ARG PRO GLN

SER
ALA
PRO
ALA
ALA
PRO
SER
ASN
GLU
PRO
PRO
MET
ASP
PHE
ASP
ASP
ASP
ILE
PRO
PHE

4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.37Å 62.92Å 97.79Å 90.00° 112.57° 90.00°	Depositor
Resolution (Å)	51.96 – 2.20 51.96 – 2.20	Depositor EDS
% Data completeness (in resolution range)	82.7 (51.96-2.20) 82.7 (51.96-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.210 , 0.262 0.208 , 0.257	Depositor DCC
R_{free} test set	1263 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	46.3	Xtriage
Anisotropy	0.169	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.024 for $1/2^*h+3/2^*k, 1/2^*h-1/2^*k, -1/2^*h-1/2^*k-l$ 0.020 for $1/2^*h-3/2^*k, -1/2^*h-1/2^*k, -1/2^*h+1/2^*k-l$	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3691	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	0/913	1.00	3/1232 (0.2%)
1	B	1.02	3/897 (0.3%)	1.02	0/1210
1	C	1.02	1/897 (0.1%)	1.01	3/1210 (0.2%)
1	D	0.84	1/897 (0.1%)	0.96	3/1210 (0.2%)
All	All	0.96	5/3604 (0.1%)	1.00	9/4862 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	92	SER	C-N	11.48	1.53	1.33
1	B	92	SER	C-N	8.96	1.49	1.33
1	B	92	SER	C-O	7.13	1.36	1.23
1	B	80	GLU	CB-CG	-6.31	1.40	1.52
1	D	99	THR	CB-CG2	-5.13	1.35	1.52

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	32	ILE	CB-CA-C	-7.31	96.98	111.60
1	C	84	ARG	NE-CZ-NH2	-6.33	117.14	120.30
1	A	96	ARG	NE-CZ-NH2	-6.23	117.19	120.30
1	C	41	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	A	86	ARG	NE-CZ-NH2	-5.33	117.64	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1	ALA	Peptide
1	C	42	ASP	Peptide
1	C	92	SER	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	894	0	893	40	0
1	B	884	0	880	37	0
1	C	884	0	880	36	0
1	D	884	0	880	43	0
2	A	30	0	0	7	0
2	B	45	0	0	1	0
2	C	42	0	0	1	0
2	D	28	0	0	3	0
All	All	3691	0	3533	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:104:ASN:ND2	1:B:105:VAL:HG22	1.28	1.44
1:B:105:VAL:HG23	1:B:106:GLY:H	1.15	1.09
1:A:20:VAL:HG21	1:A:68:SER:HB2	1.41	1.03
1:B:104:ASN:ND2	1:B:105:VAL:CG2	2.22	1.02
1:D:20:VAL:HG21	1:D:68:SER:HB2	1.41	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/178 (64%)	103 (90%)	5 (4%)	6 (5%)	2	0
1	B	112/178 (63%)	103 (92%)	6 (5%)	3 (3%)	5	2
1	C	112/178 (63%)	100 (89%)	5 (4%)	7 (6%)	1	0
1	D	112/178 (63%)	101 (90%)	9 (8%)	2 (2%)	8	5
All	All	450/712 (63%)	407 (90%)	25 (6%)	18 (4%)	3	1

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	TRP
1	A	41	ARG
1	C	2	SER
1	C	42	ASP
1	C	113	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/136 (70%)	79 (83%)	16 (17%)	2	1
1	B	93/136 (68%)	76 (82%)	17 (18%)	1	1
1	C	93/136 (68%)	79 (85%)	14 (15%)	3	2
1	D	93/136 (68%)	74 (80%)	19 (20%)	1	1
All	All	374/544 (69%)	308 (82%)	66 (18%)	2	1

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	77	VAL
1	D	92	SER
1	D	108	THR
1	B	43	LYS
1	B	42	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	94	GLN
1	D	76	GLN
1	B	104	ASN
1	D	82	GLN
1	D	25	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	114/178 (64%)	1.05	21 (18%) 1 1	30, 60, 143, 196	0
1	B	114/178 (64%)	0.49	12 (10%) 6 5	34, 59, 142, 176	0
1	C	114/178 (64%)	0.57	11 (9%) 8 6	32, 59, 130, 161	0
1	D	114/178 (64%)	0.64	12 (10%) 6 5	33, 63, 148, 189	0
All	All	456/712 (64%)	0.69	56 (12%) 4 3	30, 61, 143, 196	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	45	THR	10.9
1	A	114	GLY	10.9
1	A	46	GLY	9.3
1	A	41	ARG	8.6
1	B	44	ALA	7.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.