

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 12, 2024 – 11:49 AM EDT

PDB ID	:	6N80
Title	:	S. aureus ClpP bound to anti-4a
Authors	:	Lee, R.E.; Griffith, E.C.
Deposited on	:	2018-11-28
Resolution	:	1.96  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	Δ	203	2%		
	A	205	86%	•	12%
1	В	203	84%	•	13%
1	С	<u> </u>	4%		
1	U	203	88%	•	9%
1	D	203	88%	•	10%
1	Е	203	4%		100/
	Ľ	200	85%	•	10%



Mol	Chain	Length	Quality of chain	
1	F	203	85% •	12%
1	G	203	3% 83% 5%	12%
1	Ι	203	4% 87% •	11%
1	K	203	2% 84% ·	12%
1	L	203	3% 87%	9%
1	М	203	4% 85% 5%	10%
1	N	203	2%	12%
1	S	203	3% 	12%
1	Т	203	87%	12%



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 21063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	179	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
	11	115	1354	853	231	264	6	0	I	0
1	В	177	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
	D	111	1326	836	225	259	6	0	0	0
1	С	184	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
-		101	1389	875	237	271	6	Ŭ	-	Ŭ
1	D	183	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
-		100	1385	873	235	271	6	Ŭ	Ŭ	Ŭ
1	Е	182	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
-		102	1378	867	233	272	6	Ŭ	Ŭ	
1	F	179	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
-	1	110	1354	851	232	265	6	Ŭ	1	
1	G	179	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
		110	1356	854	232	264	6	Ŭ	-	
1	Ι	180	Total	С	Ν	Ο	S	0	0	0
-	1	100	1358	852	230	270	6	0	0	
1	К	178	Total	С	Ν	Ο	$\mathbf{S}$	0	0 0	0
		110	1337	841	229	261	6	0	0	0
1	L	184	Total	С	Ν	Ο	S	0	1	0
		101	1398	881	236	275	6	0	1	0
1	М	183	Total	С	Ν	Ο	$\mathbf{S}$	0	2	0
-		100	1413	890	238	279	6	Ŭ	-	0
1	Ν	179	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
	11	110	1358	855	231	266	6	0	0	0
1	S	178	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
		110	1343	845	231	261	6		1	
1	Т	179	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
	1	115	1343	845	231	261	6		1	U

• Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference				
А	196	LEU	-	expression tag	UNP Q2G036				



Chain	Residue	Modelled	Actual	Comment	Reference
А	197	GLU	-	expression tag	UNP Q2G036
А	198	HIS	-	expression tag	UNP Q2G036
А	199	HIS	-	expression tag	UNP Q2G036
А	200	HIS	-	expression tag	UNP Q2G036
А	201	HIS	-	expression tag	UNP Q2G036
А	202	HIS	-	expression tag	UNP Q2G036
А	203	HIS	-	expression tag	UNP Q2G036
В	196	LEU	-	expression tag	UNP Q2G036
В	197	GLU	-	expression tag	UNP Q2G036
В	198	HIS	-	expression tag	UNP Q2G036
В	199	HIS	-	expression tag	UNP Q2G036
В	200	HIS	-	expression tag	UNP Q2G036
В	201	HIS	-	expression tag	UNP Q2G036
В	202	HIS	-	expression tag	UNP Q2G036
В	203	HIS	-	expression tag	UNP Q2G036
С	196	LEU	-	expression tag	UNP Q2G036
С	197	GLU	-	expression tag	UNP Q2G036
С	198	HIS	-	expression tag	UNP Q2G036
С	199	HIS	-	expression tag	UNP Q2G036
С	200	HIS	-	expression tag	UNP Q2G036
С	201	HIS	-	expression tag	UNP Q2G036
С	202	HIS	-	expression tag	UNP Q2G036
С	203	HIS	_	expression tag	UNP Q2G036
D	196	LEU	-	expression tag	UNP Q2G036
D	197	GLU	_	expression tag	UNP Q2G036
D	198	HIS	_	expression tag	UNP Q2G036
D	199	HIS	_	expression tag	UNP Q2G036
D	200	HIS	-	expression tag	UNP Q2G036
D	201	HIS	-	expression tag	
D	202	HIS	_	expression tag	UNP Q2G036
D	203	HIS	_	expression tag	UNP Q2G036
Е	196	LEU	_	expression tag	UNP Q2G036
Е	197	GLU	_	expression tag	UNP Q2G036
Е	198	HIS	_	expression tag	UNP Q2G036
Е	199	HIS	-	expression tag	UNP Q2G036
Е	200	HIS	-	expression tag	UNP Q2G036
Е	201	HIS	-	expression tag	UNP Q2G036
	202	HIS	_	expression tag	UNP Q2G036
$\mathbf{E}$					-
E E	203	HIS	-	expression tag	$\cup$ NP Q2G030
		HIS LEU	-	expression tag	UNP Q2G036 UNP Q2G036
Е	203 196 197			expression tag expression tag	UNP Q2G036 UNP Q2G036 UNP Q2G036



Chain	Residue	vious page Modelled	Actual	Comment	Reference
F	199	HIS	_	expression tag	UNP Q2G036
F	200	HIS	_	expression tag	UNP Q2G036
F	201	HIS	-	expression tag	UNP Q2G036
F	202	HIS	-	expression tag	UNP Q2G036
F	203	HIS	-	expression tag	UNP Q2G036
G	196	LEU	-	expression tag	UNP Q2G036
G	197	GLU	-	expression tag	UNP Q2G036
G	198	HIS	-	expression tag	UNP Q2G036
G	199	HIS	-	expression tag	UNP Q2G036
G	200	HIS	-	expression tag	UNP Q2G036
G	201	HIS	-	expression tag	UNP Q2G036
G	202	HIS	-	expression tag	UNP Q2G036
G	203	HIS	-	expression tag	UNP Q2G036
Ι	196	LEU	-	expression tag	UNP Q2G036
Ι	197	GLU	-	expression tag	UNP Q2G036
Ι	198	HIS	-	expression tag	UNP Q2G036
Ι	199	HIS	-	expression tag	UNP Q2G036
Ι	200	HIS	-	expression tag	UNP Q2G036
Ι	201	HIS	-	expression tag	UNP Q2G036
Ι	202	HIS	-	expression tag	UNP Q2G036
Ι	203	HIS	-	expression tag	UNP Q2G036
К	196	LEU	-	expression tag	UNP Q2G036
Κ	197	GLU	-	expression tag	UNP Q2G036
Κ	198	HIS	-	expression tag	UNP Q2G036
Κ	199	HIS	-	expression tag	UNP Q2G036
Κ	200	HIS	-	expression tag	UNP Q2G036
Κ	201	HIS	-	expression tag	UNP Q2G036
Κ	202	HIS	-	expression tag	UNP Q2G036
Κ	203	HIS	-	expression tag	UNP Q2G036
L	196	LEU	-	expression tag	UNP $Q2G036$
L	197	GLU	-	expression tag	UNP Q2G036
L	198	HIS	-	expression tag	UNP Q2G036
L	199	HIS	-	expression tag	UNP Q2G036
L	200	HIS	-	expression tag	UNP $Q2G036$
L	201	HIS	-	expression tag	UNP Q2G036
L	202	HIS	-	expression tag	UNP Q2G036
L	203	HIS	-	expression tag	UNP Q2G036
М	196	LEU	-	expression tag	UNP Q2G036
М	197	GLU	-	expression tag	UNP Q2G036
М	198	HIS	-	expression tag	UNP Q2G036
М	199	HIS	-	expression tag	UNP Q2G036
М	200	HIS	-	expression tag	UNP Q2G036


Continued from previous page...



Chain	Residue	vious page Modelled	Actual	Comment	Reference
М	201	HIS	-	expression tag	UNP Q2G036
М	202	HIS	_	expression tag	UNP Q2G036
М	203	HIS	-	expression tag	UNP $Q2G036$
N	196	LEU	-	expression tag	UNP Q2G036
N	197	GLU	-	expression tag	UNP Q2G036
N	198	HIS	-	expression tag	UNP Q2G036
N	199	HIS	-	expression tag	UNP Q2G036
N	200	HIS	-	expression tag	UNP Q2G036
N	201	HIS	-	expression tag	UNP Q2G036
N	202	HIS	-	expression tag	UNP Q2G036
N	203	HIS	-	expression tag	UNP Q2G036
S	196	LEU	-	expression tag	UNP Q2G036
S	197	GLU	-	expression tag	UNP Q2G036
S	198	HIS	-	expression tag	UNP Q2G036
S	199	HIS	-	expression tag	UNP Q2G036
S	200	HIS	-	expression tag	UNP Q2G036
S	201	HIS	-	expression tag	UNP Q2G036
S	202	HIS	-	expression tag	UNP Q2G036
S	203	HIS	-	expression tag	UNP Q2G036
Т	196	LEU	-	expression tag	UNP Q2G036
Т	197	GLU	-	expression tag	UNP Q2G036
Т	198	HIS	-	expression tag	UNP $Q2G036$
Т	199	HIS	-	expression tag	UNP Q2G036
Т	200	HIS	-	expression tag	UNP Q2G036
Т	201	HIS	-	expression tag	UNP Q2G036
Т	202	HIS	-	expression tag	UNP Q2G036
Т	203	HIS	-	expression tag	UNP Q2G036

• Molecule 2 is N-[(1R)-1-borono-3-methylbutyl]-N 2 -(2-chloro-4-methoxybenzene-1-carbonyl )-L-leucinamide (three-letter code: JT7) (formula:  $C_{19}H_{30}BClN_2O_5$ ).





Mol	Chain	Residues		1	Aton	ns			ZeroOcc	AltConf	
0		1	Total	В	С	Cl	Ν	Ο	0	0	
2	А	1	28	1	19	1	2	5	0	0	
2	В	1	Total	В	С	Cl	Ν	0	0	0	
	D	1	28	1	19	1	2	5	0	0	
2	С	1	Total	В	С	Cl	Ν	0	0	0	
2	U	T	28	1	19	1	2	5	0	0	
2	D	1	Total	В	С	Cl	Ν	0	0	0	
	D	1	28	1	19	1	2	5	0	0	
2	Е	1	Total	В	$\mathbf{C}$	Cl	Ν	Ο	0	0	
		T	28	1	19	1	2	5	0	0	
2	F	1	Total	В	$\mathbf{C}$	$\operatorname{Cl}$	Ν	Ο	0	0	
	1	1	28	1	19	1	2	5	0	0	
2	G	1	Total	В	$\mathbf{C}$	$\operatorname{Cl}$	Ν	Ο	0	0	
	<u> </u>	1	28	1	19	1	2	5		0	
2	Ι	1	Total	В	$\mathbf{C}$	Cl	Ν	Ο	0	0	
_	-	-	28	1	19	1	2	5	Ŭ		
2	K	1	Total	В	С	Cl	Ν	Ο	0	0	
		-	28	1	19	1	2	5	Ŭ	0	
2	L	1	Total	В	$\mathbf{C}$	Cl	Ν	Ο	0	0	
		-	28	1	19	1	2	5	Ŭ		
2	М	1	Total	В	С	Cl	Ν	Ο	0	0	
		-	28	1	19	1	2	5	Ŭ		
2	Ν	1	Total	В	С	Cl	Ν	Ο	0	0	
		*	28	1	19	1	2	5	Ŭ		
2	S	1	Total	В	С	Cl	Ν	Ο	0	0	
	~	*	28	1	19	1	2	5	Ŭ		
2	Т	1	Total	В	С	Cl	Ν	Ο	0	0	
		*	28	1	19	1	2	5		ý	



• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	107	Total O 107 107	0	0
3	В	95	Total         O           95         95	0	0
3	С	112	Total         O           112         112	0	0
3	D	127	Total         O           127         127	0	0
3	Е	124	Total O 124 124	0	0
3	F	109	Total O 109 109	0	0
3	G	108	Total O 108 108	0	0
3	Ι	94	Total O 94 94	0	0
3	K	107	Total         O           107         107	0	0
3	L	137	Total         O           137         137	0	0
3	М	141	Total O 141 141	0	0
3	Ν	121	Total         O           121         121	0	0
3	S	97	Total         O           97         97	0	0
3	Т	100	Total         O           100         100	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-dependent Clp protease proteolytic subunit




• Molecule 1: ATP-dependent Clp protease proteolytic subunit






• Molecule 1: ATP-dependent Clp protease proteolytic subunit



• Molecule 1: ATP-dependent Clp protease proteolytic subunit





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	94.33Å 126.11Å 145.46Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $93.80^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	43.24 - 1.96	Depositor
Resolution (A)	43.24 - 1.96	EDS
% Data completeness	98.7 (43.24-1.96)	Depositor
(in resolution range)	99.3 (43.24-1.96)	EDS
R <sub>merge</sub>	0.16	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.10 (at 1.95 \text{\AA})$	Xtriage
Refinement program	REFMAC	Depositor
D D.	0.171 , $0.201$	Depositor
$R, R_{free}$	0.175 , $0.205$	DCC
$R_{free}$ test set	12064  reflections  (4.96%)	wwPDB-VP
Wilson B-factor $(Å^2)$	28.2	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , $40.0$	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	21063	wwPDB-VP
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section:  $\rm JT7$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.37	0/1375	0.58	2/1862~(0.1%)	
1	В	0.37	0/1343	0.63	4/1817~(0.2%)	
1	С	0.42	0/1410	0.61	2/1909~(0.1%)	
1	D	0.46	0/1403	0.62	2/1899~(0.1%)	
1	Е	0.45	0/1396	0.59	1/1890~(0.1%)	
1	F	0.43	0/1374	0.58	1/1859~(0.1%)	
1	G	0.38	0/1377	0.58	2/1864~(0.1%)	
1	Ι	0.37	0/1375	0.59	3/1861~(0.2%)	
1	Κ	0.43	0/1354	0.60	3/1833~(0.2%)	
1	L	0.48	0/1419	0.62	3/1921~(0.2%)	
1	М	0.45	0/1434	0.62	2/1938~(0.1%)	
1	N	0.42	0/1375	0.60	2/1859~(0.1%)	
1	S	0.38	0/1363	0.56	3/1844~(0.2%)	
1	Т	0.37	0/1363	0.57	2/1846~(0.1%)	
All	All	0.41	0/19361	0.60	32/26202~(0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1

There are no bond length outliers.

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	98	SER	O-C-N	-9.52	107.46	122.70
1	Т	97	ALA	O-C-N	8.33	136.03	122.70



Mol	Chain	$\mathbf{Res}$	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$		
1	В	97	ALA	O-C-N	8.33	136.03	122.70		
1	С	98	SER	O-C-N	-8.20	109.58	122.70		
1	А	97	ALA	O-C-N	7.75	135.09	122.70		

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	98	SER	Mainchain

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1354	0	1333	3	0
1	В	1326	0	1305	3	0
1	С	1389	0	1368	3	0
1	D	1385	0	1371	1	0
1	Е	1378	0	1352	5	0
1	F	1354	0	1339	5	0
1	G	1356	0	1343	6	0
1	Ι	1358	0	1330	1	0
1	K	1337	0	1318	3	0
1	L	1398	0	1378	4	0
1	М	1413	0	1405	7	0
1	N	1358	0	1343	5	0
1	S	1343	0	1328	4	0
1	Т	1343	0	1322	1	0
2	А	28	0	0	0	0
2	В	28	0	0	0	0
2	С	28	0	0	0	0
2	D	28	0	0	0	0
2	Е	28	0	0	0	0
2	F	28	0	0	0	0
2	G	28	0	0	0	0
2	Ι	28	0	0	0	0
2	K	28	0	0	0	0



	Continued from previous page									
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes				
2	L	28	0	0	0	0				
2	М	28	0	0	0	0				
2	Ν	28	0	0	0	0				
2	S	28	0	0	0	0				
2	Т	28	0	0	0	0				
3	А	107	0	0	0	0				
3	В	95	0	0	1	0				
3	С	112	0	0	1	0				
3	D	127	0	0	1	0				
3	Е	124	0	0	2	0				
3	F	109	0	0	1	0				
3	G	108	0	0	1	0				
3	Ι	94	0	0	1	0				
3	Κ	107	0	0	1	0				
3	L	137	0	0	1	0				
3	М	141	0	0	0	0				
3	Ν	121	0	0	0	0				
3	S	97	0	0	0	0				
3	Т	100	0	0	0	0				
All	All	21063	0	18835	41	0				

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:33:GLY:HA3	1:F:42[B]:ASN:OD1	2.03	0.57
1:L:33:GLY:HA3	1:M:42[B]:ASN:OD1	2.07	0.54
1:A:42[B]:ASN:ND2	1:G:33:GLY:HA3	2.24	0.53
1:E:142:HIS:HD2	3:E:517:HOH:O	1.93	0.51
1:E:75:PHE:CE2	1:E:149:LYS:HE3	2.46	0.50

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	176/203~(87%)	173~(98%)	3~(2%)	0	100	100
1	В	173/203~(85%)	170 (98%)	3(2%)	0	100	100
1	С	181/203~(89%)	178 (98%)	3 (2%)	0	100	100
1	D	179/203~(88%)	176 (98%)	3 (2%)	0	100	100
1	Е	178/203~(88%)	175 (98%)	3 (2%)	0	100	100
1	F	176/203~(87%)	173 (98%)	3 (2%)	0	100	100
1	G	176/203~(87%)	173 (98%)	3 (2%)	0	100	100
1	Ι	176/203~(87%)	173 (98%)	3 (2%)	0	100	100
1	Κ	174/203~(86%)	171 (98%)	3 (2%)	0	100	100
1	L	181/203~(89%)	178 (98%)	3 (2%)	0	100	100
1	М	181/203~(89%)	178 (98%)	3 (2%)	0	100	100
1	Ν	175/203~(86%)	172 (98%)	3 (2%)	0	100	100
1	S	175/203~(86%)	172 (98%)	3 (2%)	0	100	100
1	Т	176/203~(87%)	173 (98%)	3 (2%)	0	100	100
All	All	2477/2842 (87%)	2435 (98%)	42 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	P	erce	ntiles
1	А	141/171~(82%)	140 (99%)	1 (1%)		81	81
1	В	137/171~(80%)	136~(99%)	1 (1%)		81	81
1	С	144/171~(84%)	143~(99%)	1 (1%)		81	81
1	D	145/171~(85%)	144 (99%)	1 (1%)		81	81
1	Е	144/171 (84%)	143 (99%)	1 (1%)		81	81
1	F	142/171~(83%)	141 (99%)	1 (1%)		81	81
1	G	142/171~(83%)	141 (99%)	1 (1%)		81	81
1	Ι	142/171~(83%)	141 (99%)	1 (1%)		81	81
1	K	139/171 (81%)	138 (99%)	1 (1%)		81	81
1	L	146/171~(85%)	145 (99%)	1 (1%)		81	81
1	М	150/171~(88%)	149 (99%)	1 (1%)		81	81
1	Ν	143/171 (84%)	142 (99%)	1 (1%)		81	81
1	S	140/171 (82%)	139 (99%)	1 (1%)		81	81
1	Т	139/171 (81%)	138 (99%)	1 (1%)		81	81
All	All	1994/2394~(83%)	1980 (99%)	14 (1%)		81	81

5 of 14 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	Ι	123	HIS
1	Κ	123	HIS
1	Т	123	HIS
1	Ν	123	HIS
1	S	123	HIS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such side chains are listed below:

Mol	Chain	Res	Type
1	Κ	142	HIS
1	L	142	HIS
1	Т	83	HIS
1	S	142	HIS
1	Е	142	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	Bond lengths		B	ond ang	les
WIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	$\rm JT7$	Ν	301	1	24,28,28	1.84	4 (16%)	33,38,38	1.24	<mark>6 (18%)</mark>
2	$\rm JT7$	S	301	1	24,28,28	1.80	4 (16%)	33,38,38	1.13	3 (9%)
2	JT7	С	301	1	24,28,28	1.84	4 (16%)	33,38,38	1.21	4 (12%)
2	JT7	М	301	1	24,28,28	1.84	4 (16%)	33,38,38	1.14	4 (12%)
2	JT7	Т	301	1	24,28,28	1.84	4 (16%)	33,38,38	1.30	5 (15%)
2	JT7	Е	301	1	24,28,28	1.82	5 (20%)	33,38,38	1.27	6 (18%)
2	JT7	Ι	301	1	24,28,28	1.84	5 (20%)	33,38,38	1.21	5 (15%)
2	JT7	К	301	1	24,28,28	1.86	5 (20%)	33,38,38	1.18	3 (9%)
2	JT7	В	301	1	24,28,28	1.82	4 (16%)	33,38,38	1.21	4 (12%)
2	JT7	G	301	1	24,28,28	1.86	5 (20%)	33,38,38	1.21	4 (12%)
2	JT7	F	301	1	24,28,28	1.83	5 (20%)	33,38,38	1.25	6 (18%)
2	JT7	L	301	1	24,28,28	1.85	5 (20%)	33,38,38	1.23	6 (18%)
2	JT7	А	301	1	24,28,28	1.79	4 (16%)	33,38,38	1.23	5 (15%)
2	JT7	D	301	1	24,28,28	1.81	5 (20%)	33,38,38	1.28	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JT7	Ν	301	1	-	0/25/30/30	0/1/1/1
2	JT7	S	301	1	-	1/25/30/30	0/1/1/1
2	JT7	С	301	1	-	1/25/30/30	0/1/1/1
2	JT7	М	301	1	-	1/25/30/30	0/1/1/1
2	JT7	Т	301	1	-	1/25/30/30	0/1/1/1
2	JT7	Ε	301	1	-	0/25/30/30	0/1/1/1
2	$\rm JT7$	Ι	301	1	-	0/25/30/30	0/1/1/1
2	JT7	Κ	301	1	-	1/25/30/30	0/1/1/1
2	JT7	В	301	1	-	0/25/30/30	0/1/1/1
2	JT7	G	301	1	-	0/25/30/30	0/1/1/1
2	JT7	F	301	1	-	0/25/30/30	0/1/1/1
2	JT7	L	301	1	-	0/25/30/30	0/1/1/1
2	JT7	А	301	1	-	1/25/30/30	0/1/1/1
2	JT7	D	301	1	-	0/25/30/30	0/1/1/1

The worst 5 of 63 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Е	301	$\rm JT7$	C10-N09	5.54	1.45	1.34
2	F	301	JT7	C10-N09	5.54	1.45	1.34
2	Κ	301	JT7	C10-N09	5.46	1.45	1.34
2	М	301	JT7	C10-N09	5.46	1.45	1.34
2	G	301	JT7	C10-N09	5.44	1.45	1.34

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	Т	301	$\rm JT7$	C13-C12-C10	-3.33	102.71	110.59
2	Е	301	JT7	O19-C18-C20	3.08	126.69	121.03
2	В	301	JT7	C13-C12-C10	-3.03	103.40	110.59
2	N	301	JT7	C13-C12-C10	-2.90	103.72	110.59
2	K	301	JT7	O19-C18-C20	2.89	126.33	121.03

There are no chirality outliers.

5 of 6 torsion outliers are listed below
--


Mol	Chain	Res	Type	Atoms
2	А	301	JT7	C05-C04-N09-C10
2	С	301	JT7	C05-C04-N09-C10
2	Κ	301	JT7	C05-C04-N09-C10
2	М	301	JT7	C05-C04-N09-C10
2	S	301	JT7	C05-C04-N09-C10




There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




























Torsions





Rings













































### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.





## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ	>2	$OWAB(A^2)$	Q<0.9
1	А	179/203~(88%)	0.23	5 (2%) 55	61	24, 33, 49, 60	1 (0%)
1	В	177/203~(87%)	0.17	7 (3%) 43	49	28, 33, 47, 72	0
1	С	184/203~(90%)	0.08	8 (4%) 40	47	21, 28, 48, 75	1 (0%)
1	D	183/203~(90%)	-0.18	6 (3%) 49	56	19, 25, 39, 71	0
1	Е	182/203~(89%)	-0.13	8 (4%) 39	46	19, 25, 43, 71	0
1	F	179/203~(88%)	-0.09	6 (3%) 48	55	19, 27, 47, 66	1 (0%)
1	G	179/203~(88%)	0.16	6 (3%) 48	55	21, 33, 49, 61	1 (0%)
1	Ι	180/203~(88%)	0.23	8 (4%) 39	46	26, 33, 50, 77	0
1	Κ	178/203~(87%)	-0.12	4 (2%) 62	68	20, 26, 42, 60	0
1	L	184/203~(90%)	-0.17	6 (3%) 49	56	17, 24, 44, 76	1 (0%)
1	М	183/203~(90%)	-0.23	8 (4%) 39	46	16, 24, 45, 68	2 (1%)
1	Ν	179/203~(88%)	-0.12	5 (2%) 55	61	20, 27, 44, 76	0
1	S	178/203~(87%)	0.24	6 (3%) 48	55	20, 33, 50, 65	1 (0%)
1	Т	179/203~(88%)	0.29	6 (3%) 48	55	22, 33, 50, 59	1 (0%)
All	All	2524/2842 (88%)	0.02	89 (3%) 47	54	16, 30, 48, 77	9 (0%)

The worst 5 of 89 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	6	THR	5.7
1	N	4	ILE	5.6
1	F	18	TYR	5.5
1	Κ	6	THR	4.9
1	F	7	VAL	4.7



#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\operatorname{\AA}^2)$	Q < 0.9
2	$\rm JT7$	В	301	28/28	0.77	0.17	35,40,47,100	0
2	$\rm JT7$	Ι	301	28/28	0.78	0.16	32,38,46,86	0
2	$\rm JT7$	G	301	28/28	0.81	0.15	32,38,44,87	0
2	$\rm JT7$	А	301	28/28	0.81	0.14	31,40,45,86	0
2	$\rm JT7$	Т	301	28/28	0.81	0.15	34,40,49,93	0
2	$\rm JT7$	F	301	28/28	0.82	0.15	27,34,43,88	0
2	$\rm JT7$	S	301	28/28	0.82	0.14	30,37,43,80	0
2	$\rm JT7$	С	301	28/28	0.82	0.15	30,35,44,90	0
2	$\rm JT7$	N	301	28/28	0.83	0.14	26,34,43,80	0
2	$\rm JT7$	L	301	28/28	0.83	0.14	25,31,41,87	0
2	$\rm JT7$	М	301	28/28	0.83	0.15	24,33,39,89	0
2	$\rm JT7$	D	301	28/28	0.84	0.14	25,32,40,84	0
2	$\rm JT7$	Е	301	28/28	0.84	0.14	25,32,40,88	0
2	$\rm JT7$	K	301	28/28	0.85	0.12	27,33,41,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







































## 6.5 Other polymers (i)

There are no such residues in this entry.

