

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 2, 2023 – 04:25 AM EDT

:	6N97
:	Methylmalonyl-CoA decarboxylase in complex with 2-sulfonate-propionyl-am
	ino(dethia)-CoA
:	Stunkard, L.M.; Dixon, A.D.; Huth, T.J.; Lohman, J.R.
:	2018-11-30
:	1.75  Å(reported)
	:

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	FAILED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.75 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 15628 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	260	Total	С	Ν	0	$\mathbf{S}$	0	18	0
	А	200	2142	1373	360	396	13	0	10	
1	В	260	Total	С	Ν	0	S	0	23	0
	D	200	2187	1406	372	395	14	0	20	0
1	1 C	260	Total	С	Ν	Ο	S	0	18	0
	U		2139	1375	366	386	12			
1	D	260	Total	С	Ν	0	S	0	23	0
	D	200	2170	1395	365	397	13	0	20	
1	Е	260	Total	С	Ν	Ο	S	0	17	0
		200	2133	1365	364	392	12	0	11	0
1	Б	F 260	Total	С	Ν	Ο	S	0	22	0
	1 F		2169	1397	368	391	13	0		U


• Molecule 1 is a protein called Methylmalonyl-CoA decarboxylase.

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	2	ALA	SER	engineered mutation	UNP P52045
В	2	ALA	SER	engineered mutation	UNP P52045
С	2	ALA	SER	engineered mutation	UNP P52045
D	2	ALA	SER	engineered mutation	UNP P52045
Е	2	ALA	SER	engineered mutation	UNP P52045
F	2	ALA	SER	engineered mutation	UNP P52045

• Molecule 2 is (2S)-sulfonate propionyl-amino(dethia)-CoA (three-letter code: YZS) (formula:  $\rm C_{24}H_{41}N_8O_{20}P_3S).$ 






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total         C         N         O         P         S           56         24         8         20         3         1	0	1
2	В	1	Total         C         N         O         P         S           56         24         8         20         3         1	0	1
2	С	1	Total         C         N         O         P         S           56         24         8         20         3         1	0	1
2	С	1	Total C N O P 31 10 5 13 3	0	1
2	D	1	Total         C         N         O         P         S           56         24         8         20         3         1	0	1
2	Е	1	Total         C         N         O         P         S           56         24         8         20         3         1	0	1
2	Е	1	Total         C         N         O         P           31         10         5         13         3	0	1
2	F	1	Total         C         N         O         P         S           56         24         8         20         3         1	0	1
2	F	1	Total         C         N         O         P           31         10         5         13         3	0	1

• Molecule 3 is (2R)-sulfonate propionyl-amino(dethia)-CoA (three-letter code: KGP) (formula:  $\rm C_{24}H_{41}N_8O_{20}P_3S).$ 






Mol	Chain	Residues		A	ton	ıs	ZeroOcc	AltConf				
3	А	1	Total	С	Ν	Ο	Р	S	0	1		
5	Л	1	56	24	8	20	3	1	0	1		
3	В	1	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	1		
5	D	1	56	24	8	20	3	1	0	T		
3	C	С	С	1	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	1
5	U	T	56	24	8	20	3	1	0	T		
3	D	1	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	1		
5	D	1	56	24	8	20	3	1	0	T		
3	Е	1	Total	$\mathbf{C}$	Ν	Ο	Р	$\mathbf{S}$	0	1		
0	5 E	1	56	24	8	20	3	1	0	T		
3	F	1	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	1		
	5 F	1	56	24	8	20	3	1	0	1		

• Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total Ni 1 1	0	0
4	Е	1	Total Ni 1 1	0	0

• Molecule 5 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total 5	${ m C} { m 3}$	N 2	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	322	Total O 341 341	0	23
6	В	286	Total         O           296         296	0	12
6	С	295	Total         O           307         307	0	16
6	D	324	Total         O           342         342	0	21
6	Е	291	Total O 304 304	0	18
6	F	316	Total O 326 326	0	15

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	86.96Å 114.61Å 192.34Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 - 1.75	Depositor
% Data completeness	97.0 (30.00-1.75)	Depositor
(in resolution range)		-
R <sub>merge</sub>	0.09	Depositor
R <sub>sym</sub>	0.09	Depositor
$< I/\sigma(I) > 1$	2.58 (at 1.75Å)	Xtriage
Refinement program	REFMAC refmac_5.8.0238	Depositor
$R, R_{free}$	0.147 , $0.179$	Depositor
Wilson B-factor $(Å^2)$	22.7	Xtriage
Anisotropy	0.122	Xtriage
L-test for twinning <sup>2</sup>	$ < L >=0.49, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	15628	wwPDB-VP
Average B, all atoms $(Å^2)$	27.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



N/L-1	<b>—</b>	Class	Der	T 1.	Bo	ond leng	ths	B	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	KGP	С	302[A]	-	47,58,58	1.28	3 (6%)	$60,\!88,\!88$	1.33	6 (10%)
5	IMD	D	303	-	$3,\!5,\!5$	0.22	0	$4,\!5,\!5$	0.72	0
3	KGP	$\mathbf{F}$	302[A]	-	$47,\!58,\!58$	1.23	3 (6%)	$60,\!88,\!88$	1.84	13 (21%)
2	YZS	С	301[B]	-	47,58,58	1.27	3 (6%)	$60,\!88,\!88$	1.29	8 (13%)
3	KGP	D	302[A]	-	47,58,58	1.24	3 (6%)	60,88,88	1.44	7 (11%)
2	YZS	F	301[B]	-	47,58,58	1.23	3 (6%)	60,88,88	<mark>3.73</mark>	17 (28%)
3	KGP	В	302[A]	-	47,58,58	1.30	3 (6%)	60,88,88	1.56	9 (15%)
2	YZS	Е	304[A]	-	28,33,58	1.11	1 (3%)	$35,\!52,\!88$	1.49	6 (17%)
2	YZS	А	301[B]	-	47,58,58	1.28	3 (6%)	60,88,88	<mark>3.27</mark>	12 (20%)
2	YZS	С	303[A]	-	28,33,58	1.33	3 (10%)	35,52,88	1.46	6 (17%)
2	YZS	Е	302[B]	-	47,58,58	1.19	2 (4%)	60,88,88	<mark>3.19</mark>	11 (18%)
3	KGP	А	302[A]	-	47,58,58	1.29	3 (6%)	60,88,88	<mark>3.26</mark>	13 (21%)
2	YZS	F	303[A]	-	28,33,58	1.18	2 (7%)	$35,\!52,\!88$	1.88	10 (28%)
3	KGP	Е	303[A]	-	47,58,58	1.24	2 (4%)	60,88,88	<mark>3.08</mark>	13 (21%)
2	YZS	D	301[B]	-	47,58,58	1.25	3 (6%)	60,88,88	1.36	7 (11%)
2	YZS	В	301[B]	-	47,58,58	1.26	3 (6%)	60,88,88	1.39	10 (16%)

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KGP	С	302[A]	-	-	0/54/77/77	0/3/3/3
5	IMD	D	303	-	-	-	0/1/1/1
3	KGP	F	302[A]	-	-	10/54/77/77	0/3/3/3
2	YZS	С	301[B]	-	-	2/54/77/77	0/3/3/3
3	KGP	D	302[A]	-	-	4/54/77/77	0/3/3/3
2	YZS	F	301[B]	-	-	14/54/77/77	0/3/3/3
3	KGP	В	302[A]	-	-	2/54/77/77	0/3/3/3
2	YZS	Е	304[A]	-	-	4/17/37/77	0/3/3/3
2	YZS	А	301[B]	-	-	6/54/77/77	0/3/3/3
2	YZS	С	303[A]	-	-	0/17/37/77	0/3/3/3
2	YZS	Е	302[B]	-	-	13/54/77/77	0/3/3/3

Continued on next page...



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KGP	А	302[A]	-	-	4/54/77/77	0/3/3/3
2	YZS	F	303[A]	-	-	9/17/37/77	0/3/3/3
3	KGP	Е	303[A]	-	-	11/54/77/77	0/3/3/3
2	YZS	D	301[B]	-	-	6/54/77/77	0/3/3/3
2	YZS	В	301[B]	-	-	3/54/77/77	0/3/3/3

Continued from previous page...

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
3	В	302[A]	KGP	CS1-N	6.97	1.48	1.33
3	Е	303[A]	KGP	CS1-N	6.55	1.47	1.33
2	В	301[B]	YZS	CS1-N	6.49	1.47	1.33
3	С	302[A]	KGP	CS1-N	6.44	1.47	1.33
2	С	301[B]	YZS	CS1-N	6.29	1.47	1.33

The worst 5 of 148 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	Е	302[B]	YZS	CP8-CPA-CPB	-14.03	85.35	108.23
3	Е	303[A]	KGP	CP8-CPA-CPB	-13.96	85.47	108.23
2	F	301[B]	YZS	CP8-CPA-CP7	-13.15	86.02	108.82
2	А	301[B]	YZS	CP8-CPA-CPB	-12.64	87.62	108.23
3	А	302[A]	KGP	CP8-CPA-CPB	-12.63	87.63	108.23

There are no chirality outliers.

 $5~{\rm of}~88$  torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	301[B]	YZS	CP7-CPA-CPB-O7
2	А	301[B]	YZS	OP3-CP7-CPA-CP9
2	А	301[B]	YZS	CP6-CP7-CPA-CP9
2	А	301[B]	YZS	N-CS1-CS2-CS3
2	В	301[B]	YZS	C3'-O3'-P3-O31


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is




within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















6N97



















## 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

#### 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

#### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

#### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

