



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 29, 2023 – 01:30 PM EDT

PDB ID : 3NBB
Title : Crystal structure of mutant Y305F expressed in E. coli in the copper amine oxidase from hansenula polymorpha
Authors : Chen, Z.; Datta, S.; DuBois, J.L.; Klinman, J.P.; Mathews, F.S.
Deposited on : 2010-06-03
Resolution : 2.05 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	694	 4% 81% 13% ..
1	B	694	 3% 81% 13% ..
1	C	694	 2% 79% 15% ..
1	D	694	 5% 80% 14% ..
1	E	694	 2% 80% 14% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	694	 <p>3% 81% 13% . .</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 34758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisomal primary amine oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	661	5263	3345	903	991	24	0	2	0
1	B	661	5263	3345	903	991	24	0	2	0
1	C	661	5263	3345	903	991	24	0	2	0
1	D	661	5263	3345	903	991	24	0	2	0
1	E	661	5263	3345	903	991	24	0	2	0
1	F	661	5263	3345	903	991	24	0	2	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	305	PHE	TYR	engineered mutation	UNP P12807
B	305	PHE	TYR	engineered mutation	UNP P12807
C	305	PHE	TYR	engineered mutation	UNP P12807
D	305	PHE	TYR	engineered mutation	UNP P12807
E	305	PHE	TYR	engineered mutation	UNP P12807
F	305	PHE	TYR	engineered mutation	UNP P12807

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	B	1	Total	Cu	0	0
			1	1		
2	C	1	Total	Cu	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Cu 1 1	0	0
2	E	1	Total Cu 1 1	0	0
2	F	1	Total Cu 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	531	Total O 531 531	0	0
3	B	528	Total O 528 528	0	0
3	C	531	Total O 531 531	0	0
3	D	528	Total O 528 528	0	0
3	E	526	Total O 526 526	0	0
3	F	530	Total O 530 530	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Peroxisomal primary amine oxidase





- Molecule 1: Peroxisomal primary amine oxidase





- Molecule 1: Peroxisomal primary amine oxidase





● Molecule 1: Peroxisomal primary amine oxidase






● Molecule 1: Peroxisomal primary amine oxidase





● Molecule 1: Peroxisomal primary amine oxidase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	94.52Å 232.62Å 104.15Å 90.00° 91.51° 90.00°	Depositor
Resolution (Å)	36.13 – 2.05 59.94 – 2.05	Depositor EDS
% Data completeness (in resolution range)	73.3 (36.13-2.05) 73.2 (59.94-2.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.65 (at 2.05Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.193 , 0.200 0.190 , 0.196	Depositor DCC
R_{free} test set	10829 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtrriage
Anisotropy	0.924	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	34758	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TY9, CU, ME0, TY8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	1/5371 (0.0%)	0.68	2/7307 (0.0%)
1	B	0.35	1/5371 (0.0%)	0.68	2/7307 (0.0%)
1	C	0.35	1/5371 (0.0%)	0.68	2/7307 (0.0%)
1	D	0.35	1/5371 (0.0%)	0.68	2/7307 (0.0%)
1	E	0.35	1/5371 (0.0%)	0.68	2/7307 (0.0%)
1	F	0.35	1/5371 (0.0%)	0.68	2/7307 (0.0%)
All	All	0.35	6/32226 (0.0%)	0.68	12/43842 (0.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	672	VAL	CB-CG1	-6.08	1.40	1.52
1	D	672	VAL	CB-CG1	-6.05	1.40	1.52
1	C	672	VAL	CB-CG1	-6.05	1.40	1.52
1	F	672	VAL	CB-CG1	-6.05	1.40	1.52
1	A	672	VAL	CB-CG1	-6.04	1.40	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	641	LEU	CA-CB-CG	7.54	132.65	115.30
1	D	641	LEU	CA-CB-CG	7.54	132.64	115.30
1	A	641	LEU	CA-CB-CG	7.53	132.62	115.30
1	B	641	LEU	CA-CB-CG	7.53	132.61	115.30
1	E	641	LEU	CA-CB-CG	7.53	132.61	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5263	0	5084	75	0
1	B	5263	0	5084	78	1
1	C	5263	0	5084	82	0
1	D	5263	0	5084	80	0
1	E	5263	0	5084	84	0
1	F	5263	0	5084	82	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	531	0	0	2	0
3	B	528	0	0	4	0
3	C	531	0	0	2	0
3	D	528	0	0	2	0
3	E	526	0	0	2	0
3	F	530	0	0	4	0
All	All	34758	0	30504	461	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 461 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:669:LYS:HD3	1:F:675:GLU:OE2	1.26	1.28
1:E:23:HIS:HD2	1:E:25:LEU:H	1.11	0.97
1:F:23:HIS:HD2	1:F:25:LEU:H	1.11	0.95
1:B:23:HIS:HD2	1:B:25:LEU:H	1.11	0.95
1:D:217:LYS:HB2	1:D:217:LYS:NZ	1.82	0.94

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:356:PRO:O	1:F:675:GLU:O[2_556]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	657/694 (95%)	629 (96%)	25 (4%)	3 (0%)	29	18
1	B	657/694 (95%)	629 (96%)	25 (4%)	3 (0%)	29	18
1	C	657/694 (95%)	629 (96%)	25 (4%)	3 (0%)	29	18
1	D	657/694 (95%)	630 (96%)	24 (4%)	3 (0%)	29	18
1	E	657/694 (95%)	629 (96%)	25 (4%)	3 (0%)	29	18
1	F	657/694 (95%)	629 (96%)	25 (4%)	3 (0%)	29	18
All	All	3942/4164 (95%)	3775 (96%)	149 (4%)	18 (0%)	29	18

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	674	LYS
1	B	674	LYS
1	C	674	LYS
1	D	674	LYS
1	E	674	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	569/593 (96%)	547 (96%)	22 (4%)	32	25
1	B	569/593 (96%)	546 (96%)	23 (4%)	31	24
1	C	569/593 (96%)	547 (96%)	22 (4%)	32	25
1	D	569/593 (96%)	547 (96%)	22 (4%)	32	25
1	E	569/593 (96%)	546 (96%)	23 (4%)	31	24
1	F	569/593 (96%)	547 (96%)	22 (4%)	32	25
All	All	3414/3558 (96%)	3280 (96%)	134 (4%)	32	25

5 of 134 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	140	LEU
1	F	254	MET
1	F	633	LEU
1	C	140	LEU
1	C	127	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 106 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	286	HIS
1	E	70	GLN
1	F	382	ASN
1	D	294	HIS
1	D	452	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

18 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TY9	F	405[B]	-	13,16,17	4.61	8 (61%)	14,21,23	1.97	4 (28%)
1	TY9	A	405[B]	-	13,16,17	4.61	8 (61%)	14,21,23	1.97	4 (28%)
1	TY9	C	405[B]	-	13,16,17	4.61	8 (61%)	14,21,23	1.96	4 (28%)
1	TY9	E	405[B]	-	13,16,17	4.60	8 (61%)	14,21,23	1.96	4 (28%)
1	ME0	E	634[A]	1	5,8,9	0.56	0	2,8,10	0.42	0
1	ME0	D	634[A]	1	5,8,9	0.55	0	2,8,10	0.44	0
1	TY8	C	405[A]	1	13,16,17	4.91	9 (69%)	15,21,23	2.50	6 (40%)
1	TY8	A	405[A]	1	13,16,17	4.91	9 (69%)	15,21,23	2.51	6 (40%)
1	TY9	D	405[B]	-	13,16,17	4.61	8 (61%)	14,21,23	1.96	4 (28%)
1	ME0	F	634[A]	1	5,8,9	0.56	0	2,8,10	0.43	0
1	TY8	B	405[A]	1	13,16,17	4.91	9 (69%)	15,21,23	2.51	6 (40%)
1	TY8	D	405[A]	1	13,16,17	4.91	9 (69%)	15,21,23	2.50	6 (40%)
1	ME0	B	634[A]	1	5,8,9	0.57	0	2,8,10	0.42	0
1	TY8	E	405[A]	1	13,16,17	4.90	9 (69%)	15,21,23	2.51	6 (40%)
1	ME0	A	634[A]	1	5,8,9	0.56	0	2,8,10	0.42	0
1	TY8	F	405[A]	1	13,16,17	4.91	9 (69%)	15,21,23	2.51	6 (40%)
1	TY9	B	405[B]	-	13,16,17	4.61	8 (61%)	14,21,23	1.97	4 (28%)
1	ME0	C	634[A]	1	5,8,9	0.57	0	2,8,10	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TY9	F	405[B]	-	-	0/5/10/12	0/1/1/1
1	TY9	A	405[B]	-	-	0/5/10/12	0/1/1/1
1	TY9	C	405[B]	-	-	0/5/10/12	0/1/1/1
1	TY9	E	405[B]	-	-	0/5/10/12	0/1/1/1
1	ME0	E	634[A]	1	-	1/5/7/9	-
1	ME0	D	634[A]	1	-	1/5/7/9	-
1	TY8	C	405[A]	1	-	0/5/10/12	0/1/1/1
1	TY8	A	405[A]	1	-	0/5/10/12	0/1/1/1
1	TY9	D	405[B]	-	-	0/5/10/12	0/1/1/1
1	ME0	F	634[A]	1	-	1/5/7/9	-
1	TY8	B	405[A]	1	-	0/5/10/12	0/1/1/1
1	TY8	D	405[A]	1	-	0/5/10/12	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ME0	B	634[A]	1	-	1/5/7/9	-
1	TY8	E	405[A]	1	-	0/5/10/12	0/1/1/1
1	ME0	A	634[A]	1	-	1/5/7/9	-
1	TY8	F	405[A]	1	-	0/5/10/12	0/1/1/1
1	TY9	B	405[B]	-	-	0/5/10/12	0/1/1/1
1	ME0	C	634[A]	1	-	1/5/7/9	-

The worst 5 of 102 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	405[A]	TY8	C6-C5	8.60	1.51	1.38
1	B	405[B]	TY9	C6-C5	8.60	1.51	1.38
1	A	405[A]	TY8	C6-C5	8.56	1.51	1.38
1	A	405[B]	TY9	C6-C5	8.56	1.51	1.38
1	E	405[A]	TY8	C6-C5	8.55	1.51	1.38

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	405[A]	TY8	C1-CB-CA	6.78	125.00	114.53
1	E	405[A]	TY8	C1-CB-CA	6.75	124.97	114.53
1	A	405[A]	TY8	C1-CB-CA	6.75	124.96	114.53
1	B	405[A]	TY8	C1-CB-CA	6.74	124.94	114.53
1	C	405[A]	TY8	C1-CB-CA	6.73	124.94	114.53

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	634[A]	ME0	CB-CG-SD-CE
1	B	634[A]	ME0	CB-CG-SD-CE
1	C	634[A]	ME0	CB-CG-SD-CE
1	D	634[A]	ME0	CB-CG-SD-CE
1	E	634[A]	ME0	CB-CG-SD-CE

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	634[A]	ME0	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	659/694 (94%)	0.06	25 (3%) 40 44	18, 26, 46, 99	0
1	B	659/694 (94%)	0.12	23 (3%) 44 48	18, 26, 46, 99	0
1	C	659/694 (94%)	-0.01	14 (2%) 63 67	18, 26, 46, 99	0
1	D	659/694 (94%)	0.20	36 (5%) 25 27	18, 26, 46, 99	0
1	E	659/694 (94%)	-0.08	14 (2%) 63 67	18, 26, 46, 99	0
1	F	659/694 (94%)	0.01	23 (3%) 44 48	18, 26, 46, 99	0
All	All	3954/4164 (94%)	0.05	135 (3%) 45 49	18, 26, 46, 99	0

The worst 5 of 135 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	675	GLU	16.2
1	B	672	VAL	15.9
1	E	674	LYS	14.4
1	C	672	VAL	13.7
1	E	673	HIS	12.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TY8	B	405[A]	16/17	0.84	0.23	25,38,49,53	16
1	TY9	B	405[B]	16/17	0.84	0.23	25,38,49,53	16
1	TY8	F	405[A]	16/17	0.84	0.17	25,38,49,53	16
1	TY9	F	405[B]	16/17	0.84	0.19	25,38,49,53	16

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TY8	D	405[A]	16/17	0.85	0.17	25,38,49,53	16
1	TY8	E	405[A]	16/17	0.87	0.21	25,38,49,53	16
1	TY9	E	405[B]	16/17	0.87	0.22	25,38,49,53	16
1	TY9	C	405[B]	16/17	0.87	0.22	25,38,49,53	16
1	TY9	D	405[B]	16/17	0.87	0.17	25,38,49,53	16
1	TY8	C	405[A]	16/17	0.88	0.22	25,38,49,53	16
1	TY9	A	405[B]	16/17	0.88	0.19	25,38,49,53	16
1	TY8	A	405[A]	16/17	0.89	0.19	25,38,49,53	16
1	ME0	C	634[A]	9/10	0.92	0.16	24,26,42,47	9
1	ME0	B	634[A]	9/10	0.94	0.13	24,26,42,47	9
1	ME0	A	634[A]	9/10	0.94	0.18	24,26,42,47	9
1	ME0	E	634[A]	9/10	0.95	0.15	24,26,42,47	9
1	ME0	F	634[A]	9/10	0.96	0.10	24,26,42,47	9
1	ME0	D	634[A]	9/10	0.97	0.10	24,26,42,47	9

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CU	C	701	1/1	0.99	0.09	26,26,26,26	0
2	CU	B	701	1/1	1.00	0.12	26,26,26,26	0
2	CU	A	701	1/1	1.00	0.12	26,26,26,26	0
2	CU	D	701	1/1	1.00	0.11	26,26,26,26	0
2	CU	E	701	1/1	1.00	0.12	26,26,26,26	0
2	CU	F	701	1/1	1.00	0.08	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.