



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 05:16 pm BST

PDB ID : 5ND6  
Title : Crystal structure of apo transketolase from *Chlamydomonas reinhardtii*  
Authors : Fermani, S.; Zaffagnini, M.; Francia, F.; Pasquini, M.  
Deposited on : 2017-03-07  
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	693	 4% 88% 7% . .
1	B	693	 5% 87% 9% . .

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 11118 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	669	5132	3250	884	965	33	0	4	0
1	B	670	5141	3255	888	965	33	0	4	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	MET	-	initiating methionine	UNP A8IAN1
A	27	HIS	-	expression tag	UNP A8IAN1
A	28	HIS	-	expression tag	UNP A8IAN1
A	29	HIS	-	expression tag	UNP A8IAN1
A	30	HIS	-	expression tag	UNP A8IAN1
A	31	HIS	-	expression tag	UNP A8IAN1
A	32	HIS	-	expression tag	UNP A8IAN1
A	33	HIS	-	expression tag	UNP A8IAN1
A	34	MET	-	expression tag	UNP A8IAN1
A	35	ALA	-	expression tag	UNP A8IAN1
B	26	MET	-	initiating methionine	UNP A8IAN1
B	27	HIS	-	expression tag	UNP A8IAN1
B	28	HIS	-	expression tag	UNP A8IAN1
B	29	HIS	-	expression tag	UNP A8IAN1
B	30	HIS	-	expression tag	UNP A8IAN1
B	31	HIS	-	expression tag	UNP A8IAN1
B	32	HIS	-	expression tag	UNP A8IAN1
B	33	HIS	-	expression tag	UNP A8IAN1
B	34	MET	-	expression tag	UNP A8IAN1
B	35	ALA	-	expression tag	UNP A8IAN1

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C O	0	0
			4	2 2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	430	Total	O	0	0
			430	430		
3	B	411	Total	O	0	0
			411	411		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transketolase



- Molecule 1: Transketolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	165.21Å 74.31Å 133.72Å 90.00° 119.30° 90.00°	Depositor
Resolution (Å)	116.60 – 1.58 49.57 – 1.58	Depositor EDS
% Data completeness (in resolution range)	98.2 (116.60-1.58) 98.2 (49.57-1.58)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 1.58Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.159 , 0.192 0.174 , 0.204	Depositor DCC
$R_{free}$ test set	9518 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.5	Xtrriage
Anisotropy	0.094	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 42.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11118	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.03	0/5270	1.00	15/7151 (0.2%)
1	B	1.07	3/5280 (0.1%)	1.05	23/7163 (0.3%)
All	All	1.05	3/10550 (0.0%)	1.02	38/14314 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	470	CYS	CB-SG	-5.66	1.72	1.81
1	B	162	GLU	CD-OE1	-5.66	1.19	1.25
1	B	162	GLU	CD-OE2	5.49	1.31	1.25

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	433	LEU	CA-CB-CG	9.98	138.25	115.30
1	B	495	ARG	NE-CZ-NH1	-8.64	115.98	120.30
1	A	573	ARG	NE-CZ-NH2	-8.54	116.03	120.30
1	B	91	LEU	CB-CG-CD1	8.01	124.62	111.00
1	B	109	ASP	CB-CG-OD2	-7.77	111.31	118.30
1	B	199	ASP	CB-CG-OD1	6.98	124.58	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	185	LEU	CB-CG-CD1	6.84	122.64	111.00
1	B	143	ARG	NE-CZ-NH1	-6.79	116.91	120.30
1	A	143	ARG	NE-CZ-NH1	-6.76	116.92	120.30
1	A	63	ARG	NE-CZ-NH1	6.56	123.58	120.30
1	B	143	ARG	NE-CZ-NH2	6.51	123.56	120.30
1	A	68	ASP	CB-CG-OD2	-6.49	112.46	118.30
1	B	499	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	A	279	ARG	NE-CZ-NH2	6.22	123.41	120.30
1	A	143	ARG	NE-CZ-NH2	6.10	123.35	120.30
1	B	193	ASP	CB-CG-OD1	6.08	123.77	118.30
1	B	240	ILE	N-CA-C	5.99	127.17	111.00
1	B	409	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	A	495	ARG	NE-CZ-NH2	-5.87	117.37	120.30
1	B	108	ARG	NE-CZ-NH1	5.77	123.18	120.30
1	B	253	ASP	CB-CG-OD1	5.76	123.49	118.30
1	B	632	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	249	SER	N-CA-C	5.57	126.03	111.00
1	B	499	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	B	536	ARG	NE-CZ-NH1	5.48	123.04	120.30
1	A	573	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	446	ASP	CB-CG-OD1	5.40	123.16	118.30
1	B	354	ARG	NE-CZ-NH2	-5.37	117.62	120.30
1	A	257	ARG	NE-CZ-NH2	-5.35	117.62	120.30
1	A	495	ARG	NE-CZ-NH1	-5.33	117.64	120.30
1	A	462	ARG	NE-CZ-NH1	5.30	122.95	120.30
1	B	193	ASP	CB-CG-OD2	-5.28	113.55	118.30
1	A	109	ASP	CB-CG-OD2	-5.16	113.66	118.30
1	A	499	ARG	NE-CZ-NH2	-5.08	117.76	120.30
1	B	657	ASP	CB-CG-OD1	5.08	122.87	118.30
1	B	279	ARG	NE-CZ-NH2	5.07	122.84	120.30
1	A	657	ASP	CB-CG-OD1	5.07	122.86	118.30
1	B	608	MET	CA-CB-CG	5.05	121.88	113.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	239	LYS	Peptide
1	A	89	TYR	Sidechain
1	B	240	ILE	Peptide
1	B	432	ASP	Peptide



## 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5132	0	5038	36	0
1	B	5141	0	5048	30	1
2	A	4	0	6	0	0
3	A	430	0	0	4	0
3	B	411	0	0	7	0
All	All	11118	0	10092	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:ILE:HD13	1:A:312:HIS:CD2	1.98	0.97
1:A:248:ILE:HD11	1:B:458:ARG:HG3	1.66	0.75
1:B:417:ASN:HD21	1:B:442:LYS:H	1.39	0.70
1:A:267:HIS:HD2	3:A:1045:HOH:O	1.73	0.69
1:A:476:HIS:HD2	1:A:478:SER:OG	1.78	0.67
1:A:238:ASN:OD1	1:A:241:SER:OG	2.08	0.64
1:B:56:GLU:OE2	3:B:801:HOH:O	2.14	0.64
1:A:243:ASP:OD1	1:B:433:LEU:HB3	1.98	0.63
1:B:229:GLY:O	3:B:802:HOH:O	2.15	0.63
1:B:476:HIS:HD2	1:B:478:SER:OG	1.82	0.61
1:B:438:LEU:HA	3:B:805:HOH:O	2.00	0.60
1:B:409:ARG:HD3	3:B:921:HOH:O	2.01	0.60
1:B:267:HIS:HD2	3:B:1012:HOH:O	1.85	0.59
1:A:417:ASN:HD21	1:A:442:LYS:H	1.52	0.57
1:A:242:ILE:HD13	1:A:312:HIS:NE2	2.20	0.56
1:B:193:ASP:OD2	1:B:372:LYS:NZ	2.38	0.56
1:A:674:LYS:CE	3:A:1295:HOH:O	2.54	0.56
1:A:242:ILE:HD12	1:A:245:HIS:HA	1.89	0.55
1:B:155:ASN:HD21	1:B:504[A]:SER:HB2	1.72	0.55
1:B:437:ASN:O	3:B:804:HOH:O	2.18	0.55
1:B:663:SER:OG	1:B:682:HIS:HD2	1.89	0.55
1:A:245:HIS:O	1:A:246:THR:HG22	2.08	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:433:LEU:HB3	1:A:437:ASN:OD1	2.08	0.53
1:A:674:LYS:HE2	3:A:1295:HOH:O	2.09	0.53
1:A:712:ALA:O	1:A:715:THR:HG22	2.09	0.52
1:A:663:SER:OG	1:A:682:HIS:HD2	1.92	0.52
1:A:155:ASN:HD21	1:A:504:SER:HB2	1.75	0.51
1:A:245:HIS:HB3	1:A:248:ILE:HG23	1.92	0.51
1:A:208:ASP:OD2	1:A:241:SER:OG	2.29	0.50
1:A:242:ILE:HG13	1:A:243:ASP:OD1	2.12	0.50
1:A:240:ILE:O	1:A:240:ILE:HG22	2.11	0.50
1:B:200:HIS:NE2	1:B:476:HIS:HE1	2.10	0.49
1:B:562:ASN:HD21	1:B:565:ARG:HH11	1.59	0.49
1:A:674:LYS:HE3	3:A:1295:HOH:O	2.13	0.48
1:A:200:HIS:NE2	1:A:476:HIS:HE1	2.11	0.48
1:A:437:ASN:O	1:A:438:LEU:HB2	2.12	0.48
1:A:245:HIS:CG	1:A:248:ILE:HD13	2.49	0.47
1:A:340:VAL:HB	1:A:345:TYR:CE2	2.50	0.46
1:B:562:ASN:ND2	1:B:565:ARG:HE	2.14	0.46
1:A:409:ARG:NH1	1:A:436:SER:O	2.49	0.46
1:B:245:HIS:CE1	3:B:1051:HOH:O	2.68	0.46
1:B:335:TYR:CE2	1:B:341:PRO:HG3	2.51	0.46
1:B:97:LYS:NZ	1:B:199:ASP:O	2.49	0.45
1:A:97:LYS:HE3	1:A:107:ASN:O	2.15	0.45
1:A:660:ALA:HB1	1:A:715:THR:HG23	1.98	0.45
1:B:598:LYS:NZ	1:B:630:ASN:HD21	2.15	0.45
1:B:712:ALA:O	1:B:715:THR:HG22	2.17	0.45
1:B:240:ILE:HG23	1:B:245:HIS:HA	2.00	0.44
1:A:242:ILE:HA	1:A:243:ASP:HA	1.64	0.43
1:A:437:ASN:O	1:A:438:LEU:CB	2.66	0.43
1:B:239:LYS:C	1:B:240:ILE:HG12	2.39	0.42
1:B:242:ILE:O	1:B:242:ILE:HG22	2.19	0.42
1:A:77:HIS:CD2	1:A:316:GLY:HA2	2.55	0.42
1:B:195:LYS:HE2	1:B:368:GLU:OE1	2.19	0.42
1:A:274:ASP:OD2	1:A:277:GLY:HA3	2.19	0.41
1:A:155:ASN:H	1:A:155:ASN:HD22	1.68	0.41
1:A:429:GLY:O	1:A:484:CYS:HA	2.21	0.41
1:B:77:HIS:CD2	1:B:316:GLY:HA2	2.55	0.41
1:B:429:GLY:HA3	1:B:457:LEU:O	2.21	0.40
1:B:274:ASP:OD2	1:B:277:GLY:HA3	2.21	0.40
1:B:246:THR:C	1:B:248:ILE:H	2.25	0.40
1:A:248:ILE:CD1	1:B:458:ARG:HG3	2.44	0.40
1:A:245:HIS:HB3	1:A:248:ILE:HG12	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:710:GLU:OE2	1:B:710:GLU:OE2[2_555]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	671/693 (97%)	645 (96%)	20 (3%)	6 (1%)	17	4
1	B	672/693 (97%)	651 (97%)	19 (3%)	2 (0%)	41	21
All	All	1343/1386 (97%)	1296 (96%)	39 (3%)	8 (1%)	25	8

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	240	ILE
1	A	242	ILE
1	A	246	THR
1	A	249	SER
1	B	433	LEU
1	A	239	LYS
1	A	438	LEU
1	B	247	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	538/548 (98%)	526 (98%)	12 (2%)	52	25
1	B	539/548 (98%)	525 (97%)	14 (3%)	46	19
All	All	1077/1096 (98%)	1051 (98%)	26 (2%)	49	22

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ILE
1	A	242	ILE
1	A	243	ASP
1	A	248	ILE
1	A	331	LEU
1	A	436	SER
1	A	449	LYS
1	A	573	ARG
1	A	598	LYS
1	A	608	MET
1	A	636	PHE
1	A	717	GLN
1	B	49	SER
1	B	53	ASP
1	B	91	LEU
1	B	185	LEU
1	B	239	LYS
1	B	240	ILE
1	B	245	HIS
1	B	248	ILE
1	B	313	ASP
1	B	331	LEU
1	B	388	LEU
1	B	391	ASN
1	B	495	ARG
1	B	636	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	A	102	ASN
1	A	155	ASN
1	A	170	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	267	HIS
1	A	312	HIS
1	A	413	GLN
1	A	417	ASN
1	A	471	ASN
1	A	476	HIS
1	A	496	ASN
1	A	562	ASN
1	A	682	HIS
1	B	102	ASN
1	B	155	ASN
1	B	245	HIS
1	B	267	HIS
1	B	391	ASN
1	B	413	GLN
1	B	417	ASN
1	B	437	ASN
1	B	464	HIS
1	B	471	ASN
1	B	476	HIS
1	B	496	ASN
1	B	562	ASN
1	B	630	ASN
1	B	682	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	801	-	3,3,3	0.66	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	801	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	669/693 (96%)	-0.04	27 (4%) 38 39	12, 22, 45, 175	0
1	B	670/693 (96%)	0.19	32 (4%) 30 30	12, 23, 46, 128	0
All	All	1339/1386 (96%)	0.07	59 (4%) 34 34	12, 22, 45, 175	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	242	ILE	27.4
1	B	433	LEU	18.7
1	B	242	ILE	15.5
1	B	240	ILE	13.8
1	B	436	SER	13.2
1	A	245	HIS	12.4
1	A	243	ASP	12.2
1	A	240	ILE	10.9
1	B	241	SER	10.2
1	A	241	SER	9.6
1	B	435	PRO	9.0
1	B	437	ASN	8.8
1	A	437	ASN	8.7
1	B	244	GLY	8.7
1	A	435	PRO	7.7
1	A	244	GLY	7.5
1	B	434	ALA	7.3
1	B	245	HIS	6.6
1	A	247	ASP	6.3
1	A	434	ALA	5.9
1	A	436	SER	5.8
1	B	246	THR	5.6
1	A	248	ILE	5.4
1	B	243	ASP	5.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	433	LEU	5.2
1	B	247	ASP	4.8
1	A	246	THR	4.7
1	B	718	HIS	4.1
1	B	304	GLY	3.6
1	B	312	HIS	3.3
1	B	321	PRO	3.3
1	B	49	SER	3.3
1	A	312	HIS	3.2
1	A	321	PRO	3.0
1	B	336	GLY	3.0
1	B	332	ASN	3.0
1	A	364	LYS	2.9
1	B	248	ILE	2.8
1	A	432	ASP	2.8
1	A	313	ASP	2.7
1	A	329	LYS	2.7
1	A	332	ASN	2.6
1	B	335	TYR	2.6
1	A	194	VAL	2.5
1	B	320	GLY	2.5
1	A	334	PRO	2.4
1	A	320	GLY	2.4
1	A	300	LEU	2.4
1	A	325	ALA	2.4
1	B	329	LYS	2.3
1	B	599	ALA	2.2
1	B	535	PHE	2.2
1	A	306	PRO	2.2
1	B	194	VAL	2.2
1	B	497	ALA	2.1
1	B	503	LEU	2.1
1	B	404	LYS	2.1
1	B	508	VAL	2.0
1	B	324	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	A	801	4/4	0.82	0.09	46,47,47,48	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.