



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2023 – 11:15 AM EDT

PDB ID : 6NDL
Title : Crystal structure of Staphylococcus aureus biotin protein ligase in complex with a sulfonamide inhibitor
Authors : Marshall, A.C.; Polyak, S.W.; Bruning, J.B.; Lee, K.
Deposited on : 2018-12-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3216 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Biotin Protein Ligase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2652	1690	449	507	6	0	7	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0

- Molecule 3 is 1-[4-(6-aminopurin-9-yl)butylsulfamoyl]-3-[4-[(4 {S})-2-oxidanylidene-1,3,3 {a},4,6,6 {a}-hexahydrothieno[3,4-d]imidazol-4-yl]butyl]urea (three-letter code: BQX) (formula: C₁₉H₃₀N₁₀O₄S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	70	38	20	8	4	0	1

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	464	464	464	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.03Å 94.03Å 130.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.66 – 2.00	Depositor
% Data completeness (in resolution range)	99.9 (46.66-2.00)	Depositor
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.170 , 0.222	Depositor
Wilson B-factor (Å ²)	51.1	Xtrriage
Anisotropy	0.201	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3216	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BQX	A	406[A]	-	35,38,38	5.86	19 (54%)	41,53,53	2.86	15 (36%)
2	GOL	A	404	-	5,5,5	0.90	0	5,5,5	0.97	0
2	GOL	A	401	-	5,5,5	0.97	0	5,5,5	0.97	0
2	GOL	A	405	-	5,5,5	0.91	0	5,5,5	0.96	0
2	GOL	A	402	-	5,5,5	0.95	0	5,5,5	0.99	0
3	BQX	A	406[B]	-	35,38,38	6.08	19 (54%)	41,53,53	3.26	17 (41%)
2	GOL	A	403	-	5,5,5	0.95	0	5,5,5	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BQX	A	406[A]	-	-	6/21/43/43	0/4/4/4
2	GOL	A	404	-	-	2/4/4/4	-
2	GOL	A	401	-	-	0/4/4/4	-
2	GOL	A	405	-	-	0/4/4/4	-
2	GOL	A	402	-	-	1/4/4/4	-
3	BQX	A	406[B]	-	-	8/21/43/43	0/4/4/4
2	GOL	A	403	-	-	4/4/4/4	-

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	406[B]	BQX	C08-S07	-20.78	1.49	1.82
3	A	406[A]	BQX	C08-S07	-19.48	1.51	1.82
3	A	406[B]	BQX	C01-N05	16.68	1.63	1.35
3	A	406[A]	BQX	C01-N05	16.41	1.62	1.35
3	A	406[B]	BQX	C01-N02	11.26	1.54	1.35
3	A	406[A]	BQX	C01-N02	10.99	1.53	1.35
3	A	406[B]	BQX	C04-N05	-9.23	1.29	1.45
3	A	406[A]	BQX	C04-N05	-8.97	1.29	1.45
3	A	406[A]	BQX	C15-N14	7.22	1.50	1.35
3	A	406[B]	BQX	C15-N14	6.74	1.49	1.35
3	A	406[B]	BQX	C08-C04	6.43	1.67	1.53
3	A	406[A]	BQX	C08-C04	6.17	1.66	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	406[B]	BQX	S18-N21	6.17	1.68	1.61
3	A	406[A]	BQX	C03-N02	-5.92	1.37	1.46
3	A	406[B]	BQX	C03-N02	-5.83	1.37	1.46
3	A	406[B]	BQX	C06-C03	-5.73	1.42	1.53
3	A	406[A]	BQX	S18-N21	5.53	1.68	1.61
3	A	406[A]	BQX	C06-C03	-5.47	1.43	1.53
3	A	406[A]	BQX	C03-C04	5.35	1.70	1.55
3	A	406[B]	BQX	C03-C04	5.31	1.70	1.55
3	A	406[A]	BQX	C32-N31	5.09	1.40	1.32
3	A	406[B]	BQX	C32-N31	5.04	1.40	1.32
3	A	406[B]	BQX	O19-S18	5.03	1.50	1.43
3	A	406[B]	BQX	C10-C08	4.48	1.63	1.52
3	A	406[A]	BQX	C10-C08	4.16	1.62	1.52
3	A	406[B]	BQX	O17-C15	-4.12	1.14	1.23
3	A	406[A]	BQX	O19-S18	4.12	1.49	1.43
3	A	406[B]	BQX	C15-N16	3.93	1.48	1.39
3	A	406[A]	BQX	O17-C15	-3.91	1.15	1.23
3	A	406[A]	BQX	C15-N16	3.79	1.48	1.39
3	A	406[A]	BQX	O20-S18	3.08	1.47	1.43
3	A	406[B]	BQX	O20-S18	2.86	1.47	1.43
3	A	406[B]	BQX	C34-N35	2.60	1.43	1.34
3	A	406[A]	BQX	C34-N35	2.59	1.43	1.34
3	A	406[B]	BQX	C12-C13	2.42	1.61	1.51
3	A	406[A]	BQX	C12-C13	2.37	1.61	1.51
3	A	406[A]	BQX	C32-N33	2.14	1.37	1.33
3	A	406[B]	BQX	C32-N33	2.12	1.37	1.33

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	406[B]	BQX	O20-S18-O19	-13.34	100.44	120.40
3	A	406[A]	BQX	O20-S18-O19	-10.35	104.91	120.40
3	A	406[B]	BQX	C03-C04-N05	6.42	109.57	102.67
3	A	406[A]	BQX	C03-C04-N05	6.07	109.20	102.67
3	A	406[B]	BQX	C08-C04-C03	-5.62	102.42	108.94
3	A	406[A]	BQX	C08-C04-C03	-5.24	102.85	108.94
3	A	406[B]	BQX	C04-N05-C01	-5.17	107.81	112.62
3	A	406[A]	BQX	C04-N05-C01	-4.78	108.16	112.62
3	A	406[A]	BQX	N31-C32-N33	-4.68	121.36	128.68
3	A	406[B]	BQX	C04-C03-N02	4.58	107.30	102.43
3	A	406[B]	BQX	N31-C32-N33	-4.51	121.64	128.68
3	A	406[A]	BQX	C04-C03-N02	4.34	107.04	102.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	406[B]	BQX	C06-S07-C08	4.19	98.49	89.89
3	A	406[A]	BQX	C06-S07-C08	3.83	97.76	89.89
3	A	406[B]	BQX	O19-S18-N21	3.66	113.17	106.73
3	A	406[B]	BQX	N16-C15-N14	3.58	120.80	114.25
3	A	406[A]	BQX	C06-C03-C04	-3.19	105.89	108.66
3	A	406[B]	BQX	C03-N02-C01	-3.07	108.47	112.46
3	A	406[A]	BQX	N16-C15-N14	2.85	119.46	114.25
3	A	406[B]	BQX	C27-C28-N29	-2.82	106.46	109.40
3	A	406[A]	BQX	O20-S18-N21	2.77	111.61	106.73
3	A	406[B]	BQX	C06-C03-C04	-2.71	106.31	108.66
3	A	406[A]	BQX	C27-C28-N29	-2.69	106.60	109.40
3	A	406[A]	BQX	C03-N02-C01	-2.65	109.01	112.46
3	A	406[A]	BQX	C06-C03-N02	-2.62	109.71	113.03
3	A	406[B]	BQX	C13-N14-C15	-2.61	116.54	121.75
3	A	406[B]	BQX	O17-C15-N14	-2.36	118.40	122.50
3	A	406[B]	BQX	O20-S18-N21	2.34	110.85	106.73
3	A	406[A]	BQX	N05-C01-N02	-2.33	106.57	108.76
3	A	406[A]	BQX	C10-C08-C04	-2.27	108.11	114.73
3	A	406[B]	BQX	C10-C08-C04	-2.25	108.18	114.73
3	A	406[B]	BQX	N05-C01-N02	-2.04	106.85	108.76

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	403	GOL	C1-C2-C3-O3
3	A	406[A]	BQX	N21-C22-C23-C24
3	A	406[A]	BQX	C15-N16-S18-O19
3	A	406[A]	BQX	C22-N21-S18-N16
3	A	406[A]	BQX	C22-N21-S18-O19
3	A	406[B]	BQX	C15-N16-S18-O19
3	A	406[B]	BQX	C15-N16-S18-O20
3	A	406[B]	BQX	C11-C12-C13-N14
2	A	403	GOL	O1-C1-C2-C3
2	A	404	GOL	C1-C2-C3-O3
2	A	403	GOL	O2-C2-C3-O3
2	A	403	GOL	O1-C1-C2-O2
3	A	406[B]	BQX	C22-C23-C24-C25
3	A	406[B]	BQX	C22-N21-S18-O20
3	A	406[B]	BQX	O17-C15-N16-S18
3	A	406[B]	BQX	N14-C15-N16-S18
3	A	406[B]	BQX	C08-C10-C11-C12

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	406[A]	BQX	C22-N21-S18-O20
2	A	404	GOL	O2-C2-C3-O3
2	A	402	GOL	C1-C2-C3-O3
3	A	406[A]	BQX	C08-C10-C11-C12

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.