



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2023 – 06:03 PM EST

PDB ID : 1NR9  
Title : Crystal Structure of Escherichia coli 1262 (APC5008), Putative Isomerase  
Authors : Kim, Y.; Joachimiak, A.; Edwards, A.; Skarina, T.; Savchenko, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2003-01-24  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.32.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	<div style="display: flex; align-items: center;"> <div style="width: 20px; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 20px; position: relative;"> <div style="width: 53%; height: 100%; background-color: green;"></div> <div style="width: 39%; height: 100%; background-color: yellow;"></div> <div style="width: 5%; height: 100%; background-color: orange;"></div> <div style="width: 3%; height: 100%; background-color: grey;"></div> </div> <div style="margin-left: 5px;">5%</div> </div> <p style="text-align: center;">53%      39%      • 5%</p>
1	B	223	<div style="width: 100%; height: 20px; position: relative;"> <div style="width: 48%; height: 100%; background-color: green;"></div> <div style="width: 42%; height: 100%; background-color: yellow;"></div> <div style="width: 5%; height: 100%; background-color: orange;"></div> <div style="width: 6%; height: 100%; background-color: grey;"></div> </div> <p style="text-align: center;">48%      42%      5%      6%</p>
1	C	223	<div style="display: flex; align-items: center;"> <div style="width: 20px; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 20px; position: relative;"> <div style="width: 48%; height: 100%; background-color: green;"></div> <div style="width: 41%; height: 100%; background-color: yellow;"></div> <div style="width: 6%; height: 100%; background-color: orange;"></div> <div style="width: 6%; height: 100%; background-color: grey;"></div> </div> <div style="margin-left: 5px;">5%</div> </div> <p style="text-align: center;">48%      41%      6%      6%</p>
1	D	223	<div style="width: 100%; height: 20px; position: relative;"> <div style="width: 44%; height: 100%; background-color: green;"></div> <div style="width: 45%; height: 100%; background-color: yellow;"></div> <div style="width: 6%; height: 100%; background-color: orange;"></div> <div style="width: 5%; height: 100%; background-color: grey;"></div> </div> <p style="text-align: center;">44%      45%      6%      5%</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein YCGM.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	211	1606	1020	274	305	3	4	0	0	0
1	B	210	1599	1017	271	304	3	4	0	0	0
1	C	210	1602	1018	273	304	3	4	0	0	0
1	D	211	1609	1023	274	305	3	4	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	cloning artifact	UNP P76004
A	0	HIS	-	cloning artifact	UNP P76004
A	1	MSE	MET	modified residue	UNP P76004
A	33	MSE	MET	modified residue	UNP P76004
A	111	MSE	MET	modified residue	UNP P76004
A	165	MSE	MET	modified residue	UNP P76004
A	176	MSE	MET	modified residue	UNP P76004
A	220	GLY	-	cloning artifact	UNP P76004
A	221	SER	-	cloning artifact	UNP P76004
B	-1	GLY	-	cloning artifact	UNP P76004
B	0	HIS	-	cloning artifact	UNP P76004
B	1	MSE	MET	modified residue	UNP P76004
B	33	MSE	MET	modified residue	UNP P76004
B	111	MSE	MET	modified residue	UNP P76004
B	165	MSE	MET	modified residue	UNP P76004
B	176	MSE	MET	modified residue	UNP P76004
B	220	GLY	-	cloning artifact	UNP P76004
B	221	SER	-	cloning artifact	UNP P76004
C	-1	GLY	-	cloning artifact	UNP P76004
C	0	HIS	-	cloning artifact	UNP P76004
C	1	MSE	MET	modified residue	UNP P76004

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	33	MSE	MET	modified residue	UNP P76004
C	111	MSE	MET	modified residue	UNP P76004
C	165	MSE	MET	modified residue	UNP P76004
C	176	MSE	MET	modified residue	UNP P76004
C	220	GLY	-	cloning artifact	UNP P76004
C	221	SER	-	cloning artifact	UNP P76004
D	-1	GLY	-	cloning artifact	UNP P76004
D	0	HIS	-	cloning artifact	UNP P76004
D	1	MSE	MET	modified residue	UNP P76004
D	33	MSE	MET	modified residue	UNP P76004
D	111	MSE	MET	modified residue	UNP P76004
D	165	MSE	MET	modified residue	UNP P76004
D	176	MSE	MET	modified residue	UNP P76004
D	220	GLY	-	cloning artifact	UNP P76004
D	221	SER	-	cloning artifact	UNP P76004

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0

- Molecule 3 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	60	Total O 60 60	0	0
3	B	81	Total O 81 81	0	0
3	C	60	Total O 60 60	0	0
3	D	76	Total O 76 76	0	0





- Molecule 1: Protein YCGM

Chain D: 44% 45% 6% 5%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.37Å 83.55Å 94.90Å 90.00° 93.08° 90.00°	Depositor
Resolution (Å)	47.38 – 2.70 47.38 – 2.67	Depositor EDS
% Data completeness (in resolution range)	97.0 (47.38-2.70) 97.9 (47.38-2.67)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.04 (at 2.65Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.206 , 0.275 0.212 , 0.206	Depositor DCC
$R_{free}$ test set	2614 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.2	Xtrriage
Anisotropy	0.504	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 44.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6697	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1638	0.67	0/2220
1	B	0.39	0/1631	0.68	0/2214
1	C	0.39	0/1634	0.67	0/2215
1	D	0.39	0/1641	0.67	0/2226
All	All	0.39	0/6544	0.68	0/8875

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1606	0	1596	88	0
1	B	1599	0	1594	100	0
1	C	1602	0	1593	110	0
1	D	1609	0	1601	110	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	60	0	0	4	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	81	0	0	8	0
3	C	60	0	0	6	0
3	D	76	0	0	5	0
All	All	6697	0	6384	386	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 386 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:146:THR:HG23	1:B:210:ASP:HB2	1.31	1.12
1:D:146:THR:HG23	1:D:210:ASP:HB2	1.29	1.10
1:C:173:ILE:HA	1:C:176:MSE:HE2	1.32	1.09
1:C:181:THR:HG23	1:D:125:ASP:OD2	1.55	1.06
1:D:173:ILE:HA	1:D:176:MSE:HE2	1.40	1.00

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/223 (93%)	192 (93%)	13 (6%)	2 (1%)	15	37
1	B	206/223 (92%)	191 (93%)	13 (6%)	2 (1%)	15	37
1	C	206/223 (92%)	184 (89%)	18 (9%)	4 (2%)	8	20
1	D	207/223 (93%)	190 (92%)	14 (7%)	3 (1%)	11	28
All	All	826/892 (93%)	757 (92%)	58 (7%)	11 (1%)	12	30

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	210	ASP
1	B	166	ILE
1	C	166	ILE
1	C	169	ILE
1	D	196	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/178 (98%)	162 (93%)	12 (7%)	15	35
1	B	174/178 (98%)	161 (92%)	13 (8%)	13	31
1	C	174/178 (98%)	160 (92%)	14 (8%)	12	27
1	D	175/178 (98%)	158 (90%)	17 (10%)	8	19
All	All	697/712 (98%)	641 (92%)	56 (8%)	12	27

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	103	THR
1	D	216	THR
1	C	206	THR
1	D	206	THR
1	D	156	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	167	HIS
1	D	108	GLN
1	D	145	ASN
1	D	88	HIS
1	C	8	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/223 (92%)	-0.25	3 (1%) 75 77	13, 29, 53, 96	0
1	B	206/223 (92%)	-0.24	1 (0%) 91 92	14, 30, 56, 80	0
1	C	206/223 (92%)	-0.18	2 (0%) 82 83	15, 34, 60, 88	0
1	D	207/223 (92%)	-0.20	1 (0%) 91 92	16, 33, 60, 94	0
All	All	826/892 (92%)	-0.22	7 (0%) 86 87	13, 32, 59, 96	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	0	HIS	3.8
1	A	0	HIS	3.2
1	B	37	VAL	2.7
1	A	220	GLY	2.6
1	A	137	ALA	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	D	304	1/1	0.91	0.20	24,24,24,24	0
2	MG	B	302	1/1	0.94	0.21	23,23,23,23	0
2	MG	A	301	1/1	0.95	0.20	23,23,23,23	0
2	MG	C	303	1/1	0.98	0.16	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.