

Full wwPDB X-ray Structure Validation Report (i)

Sep 12, 2023 – 10:22 PM EDT

PDB ID : 4NSY

Title : Wild-type lysobacter enzymogenes lysc endoproteinase covalently inhibited by

TLCK

Authors: Asztalos, P.; Muller, A.; Holke, W.; Sobek, H.; Rudolph, M.G.

Deposited on : 2013-11-29

Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			43%		
1	A	275	92%	٠	•
			41%		
1	В	275	91%	5%	•

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	303	_	_	X	_



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8686 atoms, of which 3741 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Lysyl endopeptidase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
1	A	264	Total 3802	C 1205	H 1841	N 349	O 395	S 12	0	7	0
1	В	265	Total 3826	C 1213	H 1854	N 350	O 397	S 12	0	8	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	267	GLY	-	expression tag	UNP Q7M135
A	268	SER	-	expression tag	UNP Q7M135
A	269	GLY	-	expression tag	UNP Q7M135
A	270	HIS	-	expression tag	UNP Q7M135
A	271	HIS	-	expression tag	UNP Q7M135
A	272	HIS	-	expression tag	UNP Q7M135
A	273	HIS	-	expression tag	UNP Q7M135
A	274	HIS	-	expression tag	UNP Q7M135
A	275	HIS	_	expression tag	UNP Q7M135
В	267	GLY	-	expression tag	UNP Q7M135
В	268	SER	-	expression tag	UNP Q7M135
В	269	GLY	_	expression tag	UNP Q7M135
В	270	HIS	-	expression tag	UNP Q7M135
В	271	HIS	-	expression tag	UNP Q7M135
В	272	HIS	-	expression tag	UNP Q7M135
В	273	HIS	-	expression tag	UNP Q7M135
В	274	HIS	-	expression tag	UNP Q7M135
В	275	HIS	_	expression tag	UNP Q7M135

• Molecule 2 is N-[(2S,3S)-7-amino-1-chloro-2-hydroxyheptan-3-yl]-4-methylbenzenesulfonami de (Bound Form) (three-letter code: 2OY) (formula: C₁₄H₂₃ClN₂O₃S).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	Λ	1	Total	С	Н	N	О	S	0	0
	A	1	43	14	23	2	3	1	0	
2	D	1	Total	С	Н	N	О	S	0	0
2	Ъ	D 1	43	14	23	2	3	1		

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Cl 2 2	0	0

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	2	Total Ca 2 2	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	484	Total O 486 486	0	2
5	В	480	Total O 482 482	0	2




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lysyl endopeptidase

GLY SER GLY HIS HIS HIS HIS HIS





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	39.58Å 135.81Å 45.59Å	Depositor
a, b, c, α , β , γ	90.00° 97.19° 90.00°	Depositor
Resolution (Å)	33.99 - 1.10	Depositor
Resolution (A)	37.64 - 1.10	EDS
% Data completeness	94.8 (33.99-1.10)	Depositor
(in resolution range)	87.4 (37.64-1.10)	EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.30 (at 1.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1520)	Depositor
D D.	0.164 , 0.208	Depositor
R, R_{free}	0.164 , 0.208	DCC
R_{free} test set	9150 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	7.2	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35 , 48.8	EDS
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8686	wwPDB-VP
Average B, all atoms $(Å^2)$	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.15% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 2OY, CL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.38	0/2027	0.60	0/2760	
1	В	0.38	0/2041	0.59	0/2779	
All	All	0.38	0/4068	0.60	0/5539	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1961	1841	1841	7	0
1	В	1972	1854	1854	9	0
2	A	20	23	20	0	0
2	В	20	23	20	0	0
3	A	2	0	0	6	0
4	В	2	0	0	0	0
5	A	486	0	0	7	2
5	В	482	0	0	5	1
All	All	4945	3741	3735	21	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
3:A:303:CL:CL	5:A:760:HOH:O	2.15	1.01
3:A:303:CL:CL	5:A:859:HOH:O	2.37	0.77
3:A:303:CL:CL	5:A:866:HOH:O	2.39	0.77
3:A:303:CL:CL	5:A:860:HOH:O	2.41	0.75
1:A:151:ASN:OD1	3:A:303:CL:CL	2.47	0.70
3:A:303:CL:CL	5:A:780:HOH:O	2.47	0.68
1:B:204:ARG:NH2	5:B:683:HOH:O	2.27	0.68
1:B:230:VAL:HG11	5:B:755:HOH:O	2.01	0.60
1:B:28:TYR:CZ	1:B:36[B]:CYS:HB3	2.43	0.54
1:A:28:TYR:CZ	1:A:36[B]:CYS:HB3	2.45	0.52
1:A:152:VAL:O	5:A:741:HOH:O	2.21	0.47
1:A:97:ALA:O	5:A:696:HOH:O	2.21	0.44
1:B:136:ARG:N	5:B:755:HOH:O	2.50	0.43
1:B:40[B]:LEU:HG	1:B:131:TRP:CD1	2.52	0.43
1:B:259:ASP:OD2	5:B:699:HOH:O	2.21	0.43
1:A:198:ILE:HB	1:A:227:TYR:CE2	2.54	0.43
1:A:40[A]:LEU:HG	1:A:73:TRP:HH2	1.84	0.43
1:A:40[B]:LEU:HG	1:A:131:TRP:CD1	2.55	0.42
1:B:40[A]:LEU:HD11	1:B:129:LEU:HD13	2.02	0.41
1:B:30:LYS:NZ	5:B:823:HOH:O	2.53	0.41
1:B:53[B]:LEU:HG	1:B:116:LEU:HD13	2.03	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
5:B:715:HOH:O	5:B:726:HOH:O[1_455]	2.13	0.07
5:A:735:HOH:O	5:A:759:HOH:O[1_455]	2.17	0.03
5:A:865:HOH:O	5:A:866:HOH:O[1_455]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$269/275 \ (98\%)$	263 (98%)	6 (2%)	0	100	100
1	В	271/275 (98%)	264 (97%)	7 (3%)	0	100	100
All	All	540/550 (98%)	527 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	208/209 (100%)	205 (99%)	3 (1%)	67	30	
1	В	209/209 (100%)	206 (99%)	3 (1%)	67	30	
All	All	417/418 (100%)	411 (99%)	6 (1%)	65	30	

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	114	PHE
1	A	152	VAL
1	В	76	GLN
1	В	114	PHE
1	В	152	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mal Tyma Chain		Res	Dog	Dog	Dec	Dog	Dog	Dog	Dog	Dog	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2								
2	2OY	В	301	1	19,20,21	0.95	1 (5%)	23,27,28	1.85	2 (8%)								
2	2OY	A	301	1	19,20,21	1.07	1 (5%)	23,27,28	1.84	2 (8%)								

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2OY	В	301	1	-	1/20/20/22	0/1/1/1
2	2OY	A	301	1	-	1/20/20/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
2	A	301	2OY	S-N	2.45	1.65	1.61
2	В	301	2OY	S-N	2.15	1.65	1.61

All (4) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	301	2OY	OS2-S-OS1	-7.37	110.50	119.55
2	A	301	2OY	OS2-S-OS1	-7.25	110.63	119.55
2	В	301	2OY	OS2-S-C7	2.78	111.39	107.97
2	A	301	2OY	OS2-S-C7	2.52	111.07	107.97

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	В	301	2OY	CA-N-S-OS1
2	A	301	2OY	CA-N-S-OS1


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$OWAB(Å^2)$	Q < 0.9	
1	A	$264/275 \ (96\%)$	2.08	119 (45%)	0	1	5, 9, 18, 32	0
1	В	265/275~(96%)	2.10	114 (43%)	0	1	5, 9, 20, 43	0
All	All	529/550 (96%)	2.09	233 (44%)	0	1	5, 9, 20, 43	0

All (233) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	265	GLY	8.2
1	A	251	ALA	7.6
1	В	173	THR	6.3
1	A	254	GLY	5.6
1	A	264	THR	5.4
1	В	251	ALA	5.3
1	В	253	THR	5.1
1	В	254	GLY	4.9
1	A	173	THR	4.7
1	В	264	THR	4.3
1	A	36[A]	CYS	4.3
1	A	172	ALA	4.3
1	В	65	ILE	4.3
1	В	53[A]	LEU	4.3
1	A	35	TRP	4.1
1	В	36[A]	CYS	4.1
1	В	178[A]	LEU	4.1
1	В	255	ALA	4.1
1	В	40[A]	LEU	4.0
1	В	35	TRP	3.9
1	A	221	ALA	3.8
1	В	63	ALA	3.8
1	A	216	CYS	3.8
1	В	162	VAL	3.7



Continued from previous page...

Continued from previous page Mol Chain Res Type RSRZ							
Mol	Chain	Res	0 1				
1	В	82 ALA		3.7			
1	A	178[A]	LEU	3.7			
1	В	247	TRP	3.7			
1	В	261	LEU	3.7			
1	A	166	ILE	3.6			
1	В	70	VAL	3.6			
1	В	96	LEU	3.6			
1	A	90	ALA	3.6			
1	A	10	VAL	3.5			
1	В	10	VAL	3.5			
1	A	122	ALA	3.5			
1	В	166	ILE	3.5			
1	A	131	TRP	3.5			
1	A	234	TRP	3.5			
1	В	141	PHE	3.5			
1	A	253	THR	3.5			
1	A	258	ILE	3.5			
1	A	162	VAL	3.5			
1	В	58[A]	CYS	3.4			
1	A	161	THR	3.4			
1	A	147	ILE	3.4			
1	В	97	ALA	3.3			
1	В	110	ALA	3.3			
1	A	181	PHE	3.3			
1	В	257	PHE	3.3			
1	A	152	VAL	3.3			
1	В	181					
1	A	261	LEU	3.2			
1	В	209	LEU	3.2			
1	В	215	SER	3.2			
1	В	258	ILE	3.2			
1	A	255	ALA	3.2			
1	A	235	THR	3.2			
1	В	147	ILE	3.2			
1	В	114	PHE	3.2			
1	В	131	TRP	3.2			
1	A	73	TRP	3.1			
1	A	220	GLY	3.1			
1	A	141	PHE	3.1			
1	A	58[A]	CYS	3.1			
1	A	8	ILE	3.1			
1	A	202[A]	GLU	3.1			



Continued from previous page...

Continued from previous page Mol Chain Res Type RSRZ							
Mol	Chain	Res	V -				
1	A	129	LEU	3.1			
1	A	209	LEU	3.1			
1	В	161	THR	3.1			
1	В	52	PHE	3.1			
1	В	231	PHE	3.1			
1	В	25	VAL	3.1			
1	A	206	LEU	3.1			
1	A	219	THR	3.1			
1	В	126	ALA	3.0			
1	A	25	VAL	3.0			
1	A	59	GLY	3.0			
1	A	188	VAL	3.0			
1	В	51	TYR	3.0			
1	В	227	TYR	3.0			
1	В	230	VAL	3.0			
1	В	157	ILE	3.0			
1	A	97	ALA	2.9			
1	A	250	ALA	2.9			
1	A	80	CYS	2.9			
1	В	59	GLY	2.9			
1	В	234	TRP	2.9			
1	A	28	TYR	2.9			
1	В	28	TYR	2.9			
1	A	2 VAL		2.9			
1	В	129	LEU	2.9			
1	В	221	ALA	2.9			
1	A	51	TYR	2.9			
1	В	75	TYR	2.9			
1	A	230	VAL	2.8			
1	A	53	LEU	2.8			
1	A	40[A]	LEU	2.8			
1	В	216	CYS	2.8			
1	В	72	TYR	2.8			
1	В	73	TRP	2.8			
1	В	206	LEU	2.8			
1	A	215	SER	2.8			
1	В	185	SER	2.8			
1	A	153	ALA	2.8			
1	A	198	ILE	2.8			
1	В	8	ILE	2.8			
1	В	22	ILE	2.8			
1	A	247					



Continued from previous page...

Mol	$oxed{ ext{Mol Chain Res Type RS}}$					
1	В	21 VAL		2.7		
1	В	116	LEU	2.7		
1	A	199	TYR	2.7		
1	В	172	ALA	2.7		
1	В	71	VAL	2.7		
1	В	152	VAL	2.7		
1	A	182	TRP	2.7		
1	A	244	LEU	2.7		
1	В	61	THR	2.7		
1	В	199	TYR	2.7		
1	A	231	PHE	2.7		
1	В	191	PRO	2.6		
1	В	134	TRP	2.6		
1	A	227	TYR	2.6		
1	В	146	ALA	2.6		
1	A	22	ILE	2.6		
1	A	157	ILE	2.6		
1	A	197	PRO	2.6		
1	A	87	SER	2.6		
1	A	205	VAL	2.6		
1	A	37	THR	2.6		
1	A	119	LEU	2.6		
1	A	144	ALA	2.6		
1	В	142	ALA	2.6		
1	A	134	TRP	2.6		
1	В	219	THR	2.6		
1	A	55	ALA	2.6		
1	В	26 ALA		2.6		
1	В	144	ALA	2.6		
1	A	17	GLY	2.6		
1	A	192	GLY	2.6		
1	В	12	CYS	2.6		
1	A	72	TYR	2.6		
1	В	33	THR	2.5		
1	A	65	ILE	2.5		
1	A	41	VAL	2.5		
1	В	244	LEU	2.5		
1	A	27	ALA	2.5		
1	A	163	ALA	2.5		
1	В	202[A]	GLU	2.5		
1	В	37	THR	2.5		
1	A	110	ALA	2.5		



Continued from previous page...

	nued from			DCDZ	
Mol	Chain	Res	Type	RSRZ	
1	В	182	TRP	2.5	
1	A	75	TYR	2.5	
1	A	169	TYR	2.5	
1	A	86	SER	2.5	
1	A	21			
1	A	38 GLY		2.4	
1	A	125	PRO	2.4	
1	A	54	THR	2.4	
1	В	101	THR	2.4	
1	В	115	THR	2.4	
1	A	127	TYR	2.4	
1	A	52	PHE	2.4	
1	A	257	PHE	2.4	
1	В	90	ALA	2.4	
1	В	80	CYS	2.4	
1	В	2	VAL	2.4	
1	A	82	ALA	2.4	
1	A	44	SER	2.4	
1	В	130	PHE	2.4	
1	В	205	VAL	2.4	
1	В	220	220 GLY		
1	A	146 ALA		2.4	
1	A	187			
1	A	236			
1	В	38			
1	В	45	ALA	2.3	
1	В	198	ILE	2.3	
1	A	42	ASN	2.3	
1	A	60	MET	2.3	
1	A	116	LEU	2.3	
1	A	207	GLY	2.3	
1	A	6	CYS	2.3	
1	A	26	ALA	2.3	
1	В	188	VAL	2.3	
1	В	217	SER	2.3	
1	В	84	GLY	2.3	
1	В	187	GLY	2.3	
1	В	207	GLY	2.3	
1	В	226	TYR	2.3	
1	A	114	PHE	2.3	
1	В	87	SER	2.2	
1	A	66			



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	RSRZ	
1	A	133 GLY		2.2	
1	В	54	THR	2.2	
1	A	83	PRO	2.2	
1	A	117	LEU	2.2	
1	В	68	SER	2.2	
1	A	130	PHE	2.2	
1	В	11	VAL	2.2	
1	A	226	TYR	2.2	
1	В	127	TYR	2.2	
1	В	117	LEU	2.2	
1	A	150	PRO	2.2	
1	A	191	PRO	2.2	
1	A	12	CYS	2.2	
1	В	6	CYS	2.2	
1	A	246	ASP	2.2	
1	A	124	ASN	2.1	
1	A	11	VAL	2.1	
1	A	68[A]	SER	2.1	
1	A	70	VAL	2.1	
1	A	105	VAL	2.1	
1	В	125	PRO	2.1	
1	A	111	ALA	2.1	
1	В	55			
1	В	232	THR	2.1	
1	В	24[A]	SER	2.1	
1	В	60	MET	2.1	
1	A	180	VAL	2.1	
1	A	45	ALA	2.1	
1	A	33	THR	2.1	
1	A	96	LEU	2.1	
1	В	46	ASN	2.1	
1	В	104	VAL	2.1	
1	В	189	THR	2.1	
1	В	102	GLY	2.1	
1	В	34			
1	В	123	ALA	2.0	
1	В	235	THR	2.0	
1	A	71	VAL	2.0	
1	В	5	SER	2.0	
1	A	92	GLY	2.0	
1	В	57	HIS	2.0	



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	2OY	A	301	20/21	0.95	0.16	6,8,15,17	0
2	2OY	В	301	20/21	0.95	0.17	6,9,13,16	0
4	CA	В	302	1/1	0.95	0.12	32,32,32,32	0
3	CL	A	303	1/1	0.98	0.12	14,14,14,14	0
4	CA	В	303	1/1	0.99	0.17	16,16,16,16	0
3	CL	A	302	1/1	1.00	0.11	12,12,12,12	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 2OY A 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 2OY B 301:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

