

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 01:24 PM EDT

PDB ID : 6NUB

Title: Pyruvate Kinase M2 Mutant - S437Y in Complex with L-serine

Authors: Srivastava, D.; Nandi, S.; Dey, M.

Deposited on : 2019-01-31

Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as 541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 9203 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Pyruvate kinase PKM.

\mathbf{Mol}	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
1	A	522	Total 4059	C 2568	N 714	O 749	S 28	51	18	0
1	В	519	Total 4020	C 2546	N 704	O 741	S 29	3	17	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	initiating methionine	UNP P14618
A	-17	GLY	_	expression tag	UNP P14618
A	-16	SER	-	expression tag	UNP P14618
A	-15	SER	-	expression tag	UNP P14618
A	-14	HIS	-	expression tag	UNP P14618
A	-13	HIS	-	expression tag	UNP P14618
A	-12	HIS	-	expression tag	UNP P14618
A	-11	HIS	-	expression tag	UNP P14618
A	-10	HIS	-	expression tag	UNP P14618
A	-9	HIS	-	expression tag	UNP P14618
A	-8	SER	-	expression tag	UNP P14618
A	-7	SER	-	expression tag	UNP P14618
A	-6	GLY	-	expression tag	UNP P14618
A	-5	LEU	-	expression tag	UNP P14618
A	-4	VAL	-	expression tag	UNP P14618
A	-3	PRO	-	expression tag	UNP P14618
A	-2	ARG	-	expression tag	UNP P14618
A	-1	GLY	-	expression tag	UNP P14618
A	0	SER	-	expression tag	UNP P14618
A	437	TYR	SER	engineered mutation	UNP P14618
В	-18	MET	-	initiating methionine	UNP P14618
В	-17	GLY	-	expression tag	UNP P14618
В	-16	SER	-	expression tag	UNP P14618
В	-15	SER	-	expression tag	UNP P14618
В	-14	HIS	-	expression tag	UNP P14618


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	-13	HIS	-	expression tag	UNP P14618
В	-12	HIS	-	expression tag	UNP P14618
В	-11	HIS	-	expression tag	UNP P14618
В	-10	HIS	-	expression tag	UNP P14618
В	-9	HIS	-	expression tag	UNP P14618
В	-8	SER	-	expression tag	UNP P14618
В	-7	SER	-	expression tag	UNP P14618
В	-6	GLY	-	expression tag	UNP P14618
В	-5	LEU	-	expression tag	UNP P14618
В	-4	VAL	-	expression tag	UNP P14618
В	-3	PRO	-	expression tag	UNP P14618
В	-2	ARG	-	expression tag	UNP P14618
В	-1	GLY	-	expression tag	UNP P14618
В	0	SER	-	expression tag	UNP P14618
В	437	TYR	SER	engineered mutation	UNP P14618

 \bullet Molecule 2 is SERINE (three-letter code: SER) (formula: $\mathrm{C_3H_7NO_3}).$




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total C N 7 3 1		0	0
2	В	1	Total C N 7 3 1	O 3	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0

 \bullet Molecule 4 is OXALATE ION (three-letter code: OXL) (formula: $\mathrm{C}_2\mathrm{O}_4).$




Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 2 4	0	0
4	В	1	Total C O 6 2 4	0	0

 \bullet Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total K 1 1	0	0
5	В	1	Total K 1 1	0	0


• Molecule 6 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N S	0	0
0	Λ	1	3 1 1 1	U	
6	Δ	1	Total C N S	0	0
0	Λ	1	3 1 1 1	0	U
6	В	1	Total C N S	0	0
	Ъ	1	3 1 1 1		
6	В	1	Total C N S	0	0
	D	1	3 1 1 1	0	0
6	R	1	Total C N S	0	0
	ם	ı L	3 1 1 1		U


 $\bullet \ \ Molecule\ 7\ is\ DI(HYDROXYETHYL)ETHER\ (three-letter\ code:\ PEG)\ (formula:\ C_4H_{10}O_3).$





\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 7 4 3	0	0


• Molecule 8 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYL AMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total 19			0	0
8	В	1	Total 19		N 2	0	0


 \bullet Molecule 9 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $\mathrm{C_8H_{18}O_5}).$





Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
9	В	1	Total 13	C 8	O 5	0	0

 \bullet Molecule 10 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	В	1	Total C O 4 2 2	0	0

• Molecule 11 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	494	Total O 494 494	0	0
11	В	523	Total O 523 523	0	0

Mol Probity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	110.11Å 94.23Å 109.13Å	Depositor	
a, b, c, α , β , γ	90.00° 95.63° 90.00°	Depositor	
Resolution (Å)	36.81 - 1.70	Depositor	
% Data completeness	99.6 (36.81-1.70)	Depositor	
(in resolution range)	, , ,	-	
R_{merge}	0.05	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	3.76 (at 1.70Å)	Xtriage	
Refinement program	PHENIX (1.15.2_3472: ???)	Depositor	
R, R_{free}	0.154 , 0.174	Depositor	
Wilson B-factor (A^2)	13.2	Xtriage	
Anisotropy	0.519	Xtriage	
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	9203	wwPDB-VP	
Average B, all atoms (\mathring{A}^2)	17.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Res Link	Вс	Bond lengths			ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	PEG	A	607	-	6,6,6	0.49	0	5,5,5	0.70	0
9	PG4	В	608	_	12,12,12	0.51	0	11,11,11	0.62	0
8	ВЗР	A	608	-	18,18,18	0.66	0	21,23,23	1.35	3 (14%)
10	EDO	В	609	-	3,3,3	0.36	0	2,2,2	0.98	0
8	ВЗР	В	610	-	18,18,18	0.61	0	21,23,23	1.34	2 (9%)
2	SER	В	601	-	5,6,6	0.91	0	5,7,7	1.33	0
6	SCN	В	607	-	1,2,2	0.43	0	0,1,1	-	-
2	SER	A	601	-	5,6,6	1.01	0	5,7,7	1.41	1 (20%)
6	SCN	В	605	-	1,2,2	0.21	0	0,1,1	-	-
6	SCN	A	605	-	1,2,2	0.60	0	0,1,1	-	-
4	OXL	В	603	3	5,5,5	1.28	0	6,6,6	1.61	2 (33%)
4	OXL	A	603	3	5,5,5	1.22	0	6,6,6	1.61	2 (33%)
6	SCN	В	606	-	1,2,2	0.36	0	0,1,1	-	-
6	SCN	A	606	-	1,2,2	0.37	0	0,1,1	-	=


In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PEG	A	607	-	-	4/4/4/4	-
9	PG4	В	608	-	-	5/10/10/10	-
8	ВЗР	A	608	-	-	13/28/28/28	-
10	EDO	В	609	-	-	0/1/1/1	-
8	ВЗР	В	610	-	-	4/28/28/28	-
2	SER	A	601	-	-	0/6/6/6	-
2	SER	В	601	-	-	0/6/6/6	-
4	OXL	В	603	3	-	1/4/4/4	-
4	OXL	A	603	3	-	1/4/4/4	-

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mo	l Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$ \mathbf{Ideal}(^o) $
8	В	610	ВЗР	C2-N2-C8	-3.73	110.79	116.08





Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
8	A	608	ВЗР	C2-N2-C8	-3.62	110.94	116.08
8	В	610	ВЗР	C3-N1-C4	-2.75	112.17	116.08
4	A	603	OXL	O3-C1-C2	2.71	121.21	113.16
8	A	608	ВЗР	C3-N1-C4	-2.63	112.34	116.08

There are no chirality outliers.

5 of 28 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	A	608	ВЗР	C5-C4-N1-C3
8	A	608	ВЗР	C6-C4-N1-C3
8	A	608	ВЗР	C7-C4-N1-C3
8	A	608	ВЗР	C9-C8-N2-C2
8	A	608	ВЗР	C11-C8-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

