



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 30, 2023 – 05:18 AM EDT

PDB ID : 3NWD
Title : Glycoprotein B from Herpes simplex virus type 1, Y179S mutant, low-pH
Authors : Stampfer, S.D.; Lou, H.; Cohen, G.H.; Eisenberg, R.J.; Heldwein, E.E.
Deposited on : 2010-07-09
Resolution : 2.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	703	 6% 61% 25% 14%
1	B	703	 7% 60% 24% 15%
1	C	703	 8% 55% 30% 15%
1	D	703	 6% 65% 21% 14%
2	E	2	 50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	E	1	X	-	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 19481 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	597	4767	3000	835	910	22	0	1	0
1	A	606	4843	3056	843	922	22	0	1	0
1	C	601	4766	3003	831	910	22	0	0	0
1	D	606	4829	3044	846	918	21	0	1	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	28	ASP	-	expression tag	UNP P06437
B	29	PRO	-	expression tag	UNP P06437
B	58	ALA	PRO	SEE REMARK 999	UNP P06437
B	179	SER	TYR	engineered mutation	UNP P06437
B	313	SER	THR	SEE REMARK 999	UNP P06437
B	443	LEU	GLN	SEE REMARK 999	UNP P06437
A	28	ASP	-	expression tag	UNP P06437
A	29	PRO	-	expression tag	UNP P06437
A	58	ALA	PRO	SEE REMARK 999	UNP P06437
A	179	SER	TYR	engineered mutation	UNP P06437
A	313	SER	THR	SEE REMARK 999	UNP P06437
A	443	LEU	GLN	SEE REMARK 999	UNP P06437
C	28	ASP	-	expression tag	UNP P06437
C	29	PRO	-	expression tag	UNP P06437
C	58	ALA	PRO	SEE REMARK 999	UNP P06437
C	179	SER	TYR	engineered mutation	UNP P06437
C	313	SER	THR	SEE REMARK 999	UNP P06437
C	443	LEU	GLN	SEE REMARK 999	UNP P06437
D	28	ASP	-	expression tag	UNP P06437
D	29	PRO	-	expression tag	UNP P06437
D	58	ALA	PRO	SEE REMARK 999	UNP P06437

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	179	SER	TYR	engineered mutation	UNP P06437
D	313	SER	THR	SEE REMARK 999	UNP P06437
D	443	LEU	GLN	SEE REMARK 999	UNP P06437

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
2	E	2	28	16	2	10	0	0	0
2	F	2	28	16	2	10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	B	1	14	8	1	5	0	0
3	B	1	14	8	1	5	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		


- Molecule 5 is MESO-ERYTHRITOL (three-letter code: MRY) (formula: C₄H₁₀O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			8	4	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	16	Total	O	0	0
			16	16		
6	A	25	Total	O	0	0
			25	25		
6	C	18	Total	O	0	0
			18	18		
6	D	25	Total	O	0	0
			25	25		




● Molecule 1: Envelope glycoprotein B





- Molecule 1: Envelope glycoprotein B



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	117.76Å 117.76Å 318.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.56 – 2.88 48.56 – 2.88	Depositor EDS
% Data completeness (in resolution range)	88.0 (48.56-2.88) 88.0 (48.56-2.88)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.86Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.238 , 0.283 0.235 , 0.284	Depositor DCC
R_{free} test set	5310 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	64.3	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for -h,-k,l 0.468 for h,-h-k,-l 0.022 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	19481	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 73.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8395e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CL, MRY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/4966	0.39	0/6757
1	B	0.22	0/4879	0.38	0/6631
1	C	0.22	0/4880	0.38	0/6639
1	D	0.22	0/4948	0.38	0/6731
All	All	0.22	0/19673	0.38	0/26758

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4843	0	4604	113	0
1	B	4767	0	4540	123	0
1	C	4766	0	4514	144	0
1	D	4829	0	4602	111	0
2	E	28	0	25	2	0
2	F	28	0	25	3	0
3	A	42	0	39	0	0
3	B	28	0	26	0	0
3	C	42	0	39	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	14	0	13	0	0
4	A	1	0	0	0	0
4	D	1	0	0	0	0
5	C	8	0	10	1	0
6	A	25	0	0	0	0
6	B	16	0	0	0	0
6	C	18	0	0	1	0
6	D	25	0	0	0	0
All	All	19481	0	18437	491	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 491 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:MET:H	1:A:184:GLY:HA3	1.20	1.05
1:A:183:MET:N	1:A:184:GLY:HA3	1.88	0.88
1:B:259:VAL:HG22	1:B:260:GLU:H	1.45	0.82
1:C:256:PRO:HD3	1:C:266:GLY:HA2	1.64	0.79
1:D:403:PRO:HG3	1:D:476:PRO:HB3	1.65	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	603/703 (86%)	543 (90%)	56 (9%)	4 (1%)	22	52
1	B	590/703 (84%)	523 (89%)	64 (11%)	3 (0%)	29	59
1	C	595/703 (85%)	532 (89%)	58 (10%)	5 (1%)	19	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	601/703 (86%)	548 (91%)	52 (9%)	1 (0%)	47	76
All	All	2389/2812 (85%)	2146 (90%)	230 (10%)	13 (0%)	29	59

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	259	VAL
1	B	413	ILE
1	A	238	ASN
1	C	429	TYR
1	B	106	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	517/593 (87%)	505 (98%)	12 (2%)	50	79
1	B	509/593 (86%)	501 (98%)	8 (2%)	62	85
1	C	506/593 (85%)	501 (99%)	5 (1%)	76	91
1	D	514/593 (87%)	509 (99%)	5 (1%)	76	91
All	All	2046/2372 (86%)	2016 (98%)	30 (2%)	65	86

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	323	ASP
1	D	384	PHE
1	A	401	GLU
1	D	625	LEU
1	C	631	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	468	HIS
1	D	472	GLN
1	A	620	ASN
1	C	140	GLN
1	C	270	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	1,2	14,14,15	0.62	0	17,19,21	1.09	1 (5%)
2	NAG	E	2	2	14,14,15	0.48	0	17,19,21	0.70	0
2	NAG	F	1	1,2	14,14,15	0.55	0	17,19,21	1.46	4 (23%)
2	NAG	F	2	2	14,14,15	0.52	0	17,19,21	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	1/1/5/7	3/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	1	1,2	-	3/6/23/26	0/1/1/1
2	NAG	F	2	2	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	NAG	C2-N2-C7	-3.44	118.00	122.90
2	F	1	NAG	C1-O5-C5	2.54	115.63	112.19
2	F	1	NAG	O7-C7-C8	-2.45	117.51	122.06
2	E	1	NAG	C2-N2-C7	2.41	126.34	122.90
2	F	1	NAG	O7-C7-N2	2.28	126.15	121.95

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	E	1	NAG	C1

5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	1	NAG	C3-C2-N2-C7
2	F	1	NAG	C8-C7-N2-C2
2	F	1	NAG	O7-C7-N2-C2
2	E	1	NAG	C4-C5-C6-O6
2	F	2	NAG	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	NAG	2	0
2	F	1	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	1430	1	14,14,15	0.59	0	17,19,21	1.30	2 (11%)
5	MRY	C	2001	-	7,7,7	0.38	0	8,8,8	0.34	0
3	NAG	D	1398	1	14,14,15	0.56	0	17,19,21	0.59	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	1141	1	14,14,15	0.51	0	17,19,21	0.71	0
3	NAG	A	1430	1	14,14,15	0.56	0	17,19,21	0.92	1 (5%)
3	NAG	C	1430	1	14,14,15	0.48	0	17,19,21	1.16	1 (5%)
3	NAG	B	1398	1	14,14,15	0.55	0	17,19,21	0.64	0
3	NAG	A	1674	1	14,14,15	0.52	0	17,19,21	0.64	0
3	NAG	C	1674	1	14,14,15	0.53	0	17,19,21	0.63	0
3	NAG	C	1141	1	14,14,15	0.54	0	17,19,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1430	1	-	4/6/23/26	0/1/1/1
5	MRY	C	2001	-	-	0/8/8/8	-
3	NAG	D	1398	1	-	3/6/23/26	0/1/1/1
3	NAG	A	1141	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1430	1	-	2/6/23/26	0/1/1/1
3	NAG	C	1430	1	-	0/6/23/26	0/1/1/1
3	NAG	B	1398	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1674	1	-	0/6/23/26	0/1/1/1
3	NAG	C	1674	1	-	0/6/23/26	0/1/1/1
3	NAG	C	1141	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1430	NAG	C1-O5-C5	3.27	116.63	112.19
3	B	1430	NAG	O5-C1-C2	3.19	116.33	111.29
3	B	1430	NAG	C1-O5-C5	2.82	116.02	112.19
3	A	1430	NAG	C1-O5-C5	2.36	115.39	112.19

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1430	NAG	C3-C2-N2-C7
3	B	1430	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	1430	NAG	O7-C7-N2-C2
3	C	1141	NAG	C3-C2-N2-C7
3	C	1141	NAG	C8-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	2001	MRY	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	606/703 (86%)	0.48	42 (6%) 16 13	20, 66, 145, 550	0
1	B	597/703 (84%)	0.70	52 (8%) 10 7	23, 71, 174, 550	0
1	C	601/703 (85%)	0.60	53 (8%) 10 7	20, 76, 176, 550	0
1	D	606/703 (86%)	0.58	45 (7%) 14 11	22, 65, 151, 550	0
All	All	2410/2812 (85%)	0.59	192 (7%) 12 9	20, 69, 164, 550	0

The worst 5 of 192 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	600	PRO	23.9
1	C	595	ALA	22.8
1	D	600	PRO	16.5
1	B	612	LEU	15.9
1	D	604	PHE	14.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	E	1	14/15	0.91	0.10	104,104,104,104	0
2	NAG	F	2	14/15	0.92	0.14	91,91,91,91	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	E	2	14/15	0.94	0.09	89,89,89,89	0
2	NAG	F	1	14/15	0.96	0.12	77,77,77,77	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	A	1430	14/15	0.81	0.18	118,118,118,118	0
3	NAG	C	1141	14/15	0.81	0.26	113,113,113,113	0
3	NAG	C	1430	14/15	0.84	0.17	117,117,117,117	0
3	NAG	B	1430	14/15	0.85	0.20	117,117,117,117	0
3	NAG	D	1398	14/15	0.88	0.16	87,87,87,87	0
3	NAG	C	1674	14/15	0.89	0.12	77,77,77,77	0
4	CL	A	2	1/1	0.89	0.11	78,78,78,78	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	B	1398	14/15	0.90	0.14	92,92,92,92	0
4	CL	D	1	1/1	0.90	0.15	59,59,59,59	0
3	NAG	A	1674	14/15	0.92	0.14	89,89,89,89	0
3	NAG	A	1141	14/15	0.94	0.25	120,120,120,120	0
5	MRY	C	2001	8/8	0.95	0.17	69,69,69,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.