



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2021 – 05:13 am BST

PDB ID : 7ODA
Title : OXA-48-like Beta-lactamase OXA-436
Authors : Lund, B.A.; Thomassen, A.M.; Carlsen, T.J.W.; Leiros, H.K.S.
Deposited on : 2021-04-29
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	 6% 93% 6%
1	B	244	 5% 91% 7%
1	C	244	 5% 93% 5%
1	D	244	 4% 95%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16976 atoms, of which 7822 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	241	3950	1279	1945	353	365	8	0	12	0
1	B	241	3937	1276	1937	350	367	7	0	12	0
1	C	241	3968	1284	1956	352	368	8	0	16	0
1	D	241	3939	1275	1943	350	364	7	0	10	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLY	-	expression tag	UNP A0A0S2C593
B	22	GLY	-	expression tag	UNP A0A0S2C593
C	22	GLY	-	expression tag	UNP A0A0S2C593
D	22	GLY	-	expression tag	UNP A0A0S2C593

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	H			O
2	A	1	10	2	6	2	0	0
2	A	1	9	2	5	2	0	0
2	B	1	10	2	6	2	0	0
2	B	1	10	2	6	2	0	0
2	C	1	10	2	6	2	0	0
2	C	1	10	2	6	2	0	0
2	D	1	10	2	6	2	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	B	1	1	1	0	0
3	C	1	1	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	240	Total 240	O 240	0	0
4	B	282	Total 282	O 282	0	0
4	C	289	Total 289	O 289	0	0
4	D	300	Total 300	O 300	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-lactamase




- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.12Å 94.57Å 87.76Å 90.00° 109.89° 90.00°	Depositor
Resolution (Å)	44.26 – 1.80 44.26 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.9 (44.26-1.80) 89.0 (44.26-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.58 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.183 , 0.227 0.186 , 0.228	Depositor DCC
R_{free} test set	4637 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtriage
Anisotropy	0.426	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16976	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2092	0.54	1/2832 (0.0%)
1	B	0.47	0/2098	0.57	1/2843 (0.0%)
1	C	0.46	0/2130	0.57	1/2887 (0.0%)
1	D	0.46	0/2081	0.58	0/2818
All	All	0.45	0/8401	0.57	3/11380 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	221	TRP	CA-CB-CG	5.30	123.78	113.70
1	A	221	TRP	CA-CB-CG	5.30	123.77	113.70
1	B	221	TRP	CA-CB-CG	5.12	123.42	113.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2005	1945	1906	10	0
1	B	2000	1937	1894	17	0
1	C	2012	1956	1891	8	0
1	D	1996	1943	1910	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	8	11	12	4	0
2	B	8	12	12	0	0
2	C	8	12	12	0	0
2	D	4	6	6	0	0
3	B	1	0	0	1	0
3	C	1	0	0	1	0
4	A	240	0	0	2	4
4	B	282	0	0	6	5
4	C	289	0	0	5	4
4	D	300	0	0	3	2
All	All	9154	7822	7643	43	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:MET:SD	4:B:644:HOH:O	2.07	1.12
3:C:301:CL:CL	4:D:603:HOH:O	2.05	1.11
4:A:582:HOH:O	3:B:301:CL:CL	2.16	1.00
1:A:174:ARG:NH2	2:A:302:EDO:O2	2.08	0.86
1:C:137:LYS:NZ	4:C:402:HOH:O	2.17	0.74
1:A:140[B]:HIS:ND1	1:A:147:GLU:OE1	2.19	0.73
1:D:25:TRP:N	4:D:402:HOH:O	2.22	0.72
1:C:132:GLU:OE1	4:C:401:HOH:O	2.08	0.71
1:B:192:LYS:HA	1:B:195:MET:HE3	1.72	0.71
1:C:259:LYS:NZ	1:C:265:PRO:O	2.24	0.71
1:A:245:GLU:OE2	4:A:401:HOH:O	2.10	0.70
1:B:255:LYS:NZ	4:B:404:HOH:O	2.24	0.70
1:B:192:LYS:HA	1:B:195:MET:CE	2.23	0.66
1:B:82:ASP:O	4:B:401:HOH:O	2.14	0.66
1:B:60:LYS:NZ	4:B:409:HOH:O	2.34	0.60
1:B:107:ARG:NH1	4:B:406:HOH:O	2.24	0.59
1:C:140:HIS:ND1	1:C:147:GLU:OE1	2.31	0.58
1:C:107:ARG:NH2	4:C:406:HOH:O	2.38	0.56
1:D:180:LYS:HA	1:D:180:LYS:HE2	1.92	0.52
1:D:70:SER:OG	4:D:401:HOH:O	2.17	0.51
1:B:53:GLN:HG3	1:B:55:PHE:CE1	2.47	0.50
1:B:39:LYS:O	4:B:402:HOH:O	2.19	0.50
1:B:226[B]:ILE:HD11	1:B:258:LEU:CD1	2.43	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:VAL:HG21	2:A:301:EDO:H11	1.95	0.48
1:C:82:ASP:OD2	4:C:403:HOH:O	2.20	0.48
1:A:51:THR:OG1	1:A:53:GLN:HG2	2.15	0.46
1:D:140:HIS:ND1	1:D:147:GLU:OE1	2.44	0.45
1:B:65:ALA:HB1	1:B:163:ARG:HB3	1.99	0.45
1:A:120:VAL:HB	1:A:121:PRO:HD3	1.98	0.45
1:B:192:LYS:HG2	1:B:195:MET:CE	2.46	0.45
1:C:27:GLU:OE2	1:C:29:LYS:NZ	2.51	0.44
1:B:176:LEU:HD21	1:B:195:MET:HE1	2.00	0.44
1:A:53:GLN:HG3	1:A:55:PHE:CE1	2.54	0.43
1:A:171[B]:ALA:HA	2:A:302:EDO:H21	2.01	0.43
1:A:226[B]:ILE:HD11	1:A:258:LEU:CD1	2.49	0.43
1:D:226[B]:ILE:HD11	1:D:258:LEU:CD1	2.50	0.42
1:A:171[A]:ALA:HA	2:A:302:EDO:H21	2.01	0.42
1:B:226[B]:ILE:CD1	1:B:258:LEU:HD13	2.50	0.41
1:B:74:ILE:HB	1:B:75:PRO:CD	2.51	0.41
1:B:192:LYS:HA	1:B:195:MET:HE2	2.00	0.41
1:C:167[A]:THR:HG21	4:C:421:HOH:O	2.21	0.40
1:D:120:VAL:HB	1:D:121:PRO:HD3	2.03	0.40
1:B:226[B]:ILE:HD11	1:B:258:LEU:HD13	2.03	0.40

All (8) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:617:HOH:O	4:B:585:HOH:O[1_455]	2.08	0.12
4:B:586:HOH:O	4:D:671:HOH:O[1_554]	2.09	0.11
4:B:654:HOH:O	4:C:675:HOH:O[1_554]	2.09	0.11
4:A:410:HOH:O	4:C:402:HOH:O[2_546]	2.12	0.08
4:A:595:HOH:O	4:B:579:HOH:O[2_545]	2.13	0.07
4:C:572:HOH:O	4:C:631:HOH:O[2_556]	2.14	0.06
4:A:608:HOH:O	4:C:441:HOH:O[2_546]	2.18	0.02
4:B:628:HOH:O	4:D:573:HOH:O[1_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/244 (102%)	242 (98%)	6 (2%)	0	100	100
1	B	248/244 (102%)	243 (98%)	5 (2%)	0	100	100
1	C	252/244 (103%)	247 (98%)	5 (2%)	0	100	100
1	D	246/244 (101%)	241 (98%)	5 (2%)	0	100	100
All	All	994/976 (102%)	973 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/208 (102%)	213 (100%)	0	100	100
1	B	214/208 (103%)	213 (100%)	1 (0%)	88	87
1	C	217/208 (104%)	217 (100%)	0	100	100
1	D	213/208 (102%)	213 (100%)	0	100	100
All	All	857/832 (103%)	856 (100%)	1 (0%)	93	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	82	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	64	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	B	73	1	7,11,12	1.39	1 (14%)	4,12,14	0.21	0
1	KCX	D	73	1	7,11,12	1.21	1 (14%)	4,12,14	0.26	0
1	KCX	C	73	1	7,11,12	1.30	1 (14%)	4,12,14	0.68	0
1	KCX	A	73	1	7,11,12	1.31	1 (14%)	4,12,14	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	B	73	1	-	0/7/10/12	-
1	KCX	D	73	1	-	0/7/10/12	-
1	KCX	C	73	1	-	0/7/10/12	-
1	KCX	A	73	1	-	0/7/10/12	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	KCX	CE-NZ	2.73	1.51	1.45
1	B	73	KCX	CE-NZ	2.65	1.51	1.45
1	D	73	KCX	CE-NZ	2.36	1.50	1.45
1	C	73	KCX	CE-NZ	2.25	1.50	1.45

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	302	-	3,3,3	0.40	0	2,2,2	0.54	0
2	EDO	B	302	-	3,3,3	0.50	0	2,2,2	0.74	0
2	EDO	A	301	-	3,3,3	0.33	0	2,2,2	0.98	0
2	EDO	B	303	-	3,3,3	0.43	0	2,2,2	0.31	0
2	EDO	C	302	-	3,3,3	0.26	0	2,2,2	0.65	0
2	EDO	C	303	-	3,3,3	0.44	0	2,2,2	0.56	0
2	EDO	D	301	-	3,3,3	0.45	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	302	-	-	0/1/1/1	-
2	EDO	B	302	-	-	0/1/1/1	-
2	EDO	A	301	-	-	1/1/1/1	-
2	EDO	B	303	-	-	0/1/1/1	-
2	EDO	C	302	-	-	0/1/1/1	-
2	EDO	C	303	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	D	301	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	302	EDO	3	0
2	A	301	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/244 (98%)	0.63	15 (6%) 20 15	10, 20, 40, 56	0
1	B	240/244 (98%)	0.70	12 (5%) 28 23	9, 17, 43, 72	0
1	C	240/244 (98%)	0.56	11 (4%) 32 26	9, 17, 33, 62	0
1	D	240/244 (98%)	0.53	9 (3%) 40 35	10, 17, 37, 59	0
All	All	960/976 (98%)	0.61	47 (4%) 29 24	9, 18, 38, 72	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	40	THR	5.4
1	B	39	LYS	5.1
1	B	25	TRP	4.9
1	A	31	TRP	4.1
1	C	39	LYS	4.0
1	B	245	GLU	3.9
1	B	36	SER	3.8
1	A	25	TRP	3.6
1	C	92[A]	VAL	3.4
1	D	215	ILE	3.3
1	D	245	GLU	3.2
1	A	53	GLN	3.2
1	A	60	LYS	3.0
1	C	245	GLU	2.9
1	B	41	GLN	2.9
1	D	182	HIS	2.9
1	A	95	TRP	2.9
1	B	38	HIS	2.8
1	A	182	HIS	2.7
1	B	40	THR	2.7
1	D	39	LYS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	41	GLN	2.7
1	D	41	GLN	2.7
1	C	26	GLN	2.6
1	B	246	GLY	2.6
1	D	25	TRP	2.6
1	B	243	THR	2.6
1	B	31	TRP	2.5
1	A	263	ILE	2.5
1	D	101	ASP	2.5
1	A	99	THR	2.4
1	C	41	GLN	2.4
1	C	36	SER	2.4
1	C	28	ASN	2.3
1	A	104[A]	ALA	2.3
1	B	211	TYR	2.2
1	A	59	LEU	2.2
1	D	26	GLN	2.2
1	C	215	ILE	2.2
1	C	226[B]	ILE	2.1
1	B	259	LYS	2.1
1	A	30	SER	2.1
1	A	44	VAL	2.1
1	A	243	THR	2.0
1	D	242	PRO	2.0
1	A	140[A]	HIS	2.0
1	C	232[A]	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	B	73	12/13	0.89	0.12	9,12,15,15	0
1	KCX	D	73	12/13	0.90	0.13	10,12,16,17	0
1	KCX	C	73	12/13	0.92	0.12	10,15,15,16	0
1	KCX	A	73	12/13	0.93	0.12	11,12,19,20	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	B	302	4/4	0.47	0.30	21,25,26,27	0
2	EDO	A	302	4/4	0.67	0.29	21,23,26,27	0
2	EDO	D	301	4/4	0.76	0.13	21,26,26,27	0
2	EDO	A	301	4/4	0.81	0.25	21,25,26,27	0
3	CL	C	301	1/1	0.81	0.13	35,35,35,35	0
2	EDO	C	302	4/4	0.82	0.17	20,24,25,26	0
2	EDO	C	303	4/4	0.86	0.13	21,25,26,27	0
3	CL	B	301	1/1	0.94	0.08	30,30,30,30	0
2	EDO	B	303	4/4	0.95	0.13	20,24,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.