



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2023 – 01:18 PM EDT

PDB ID : 1OW0  
Title : Crystal structure of human FcαRI bound to IgA1-Fc  
Authors : Herr, A.B.; Ballister, E.R.; Bjorkman, P.J.  
Deposited on : 2003-03-27  
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	
1	B	214	
2	C	207	
2	D	207	
3	E	8	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	8	
5	G	5	
5	J	5	
6	H	3	
6	K	3	
7	I	6	
7	L	6	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	E	1	-	-	-	X
3	NAG	E	2	-	-	-	X
3	NAG	E	5	-	-	X	X
4	NAG	F	1	-	-	-	X
4	NAG	F	2	-	-	-	X
4	NAG	F	5	-	-	X	X
4	FUC	F	8	X	-	-	-
5	NAG	G	1	-	-	-	X
5	NAG	G	2	-	-	-	X
5	MAN	G	4	-	-	-	X
5	NAG	J	1	-	-	-	X
5	NAG	J	2	-	-	-	X
5	MAN	J	4	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig alpha-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	209	1435	913	245	270	7	0	0	0
1	B	209	1435	913	245	270	7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	241	ALA	CYS	engineered mutation	UNP P01876
B	241	ALA	CYS	engineered mutation	UNP P01876

- Molecule 2 is a protein called Immunoglobulin alpha Fc receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	190	1414	914	239	255	6	0	0	0
2	D	190	1414	914	239	255	6	0	0	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	196	ALA	-	expression tag	UNP P24071
C	197	ILE	-	expression tag	UNP P24071
C	198	ASP	-	expression tag	UNP P24071
C	199	GLY	-	expression tag	UNP P24071
C	200	ARG	-	expression tag	UNP P24071
C	201	ALA	-	expression tag	UNP P24071
C	202	HIS	-	expression tag	UNP P24071
C	203	HIS	-	expression tag	UNP P24071
C	204	HIS	-	expression tag	UNP P24071
C	205	HIS	-	expression tag	UNP P24071

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	206	HIS	-	expression tag	UNP P24071
C	207	HIS	-	expression tag	UNP P24071
D	196	ALA	-	expression tag	UNP P24071
D	197	ILE	-	expression tag	UNP P24071
D	198	ASP	-	expression tag	UNP P24071
D	199	GLY	-	expression tag	UNP P24071
D	200	ARG	-	expression tag	UNP P24071
D	201	ALA	-	expression tag	UNP P24071
D	202	HIS	-	expression tag	UNP P24071
D	203	HIS	-	expression tag	UNP P24071
D	204	HIS	-	expression tag	UNP P24071
D	205	HIS	-	expression tag	UNP P24071
D	206	HIS	-	expression tag	UNP P24071
D	207	HIS	-	expression tag	UNP P24071

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	8	105	59	4	42	0	0	0


- Molecule 4 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	F	8	105	59	4	42	0	0	0


- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[beta-D-mannopyranose-

se-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	G	5	Total	C	N	O	0	0	0
			61	34	2	25			
5	J	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	H	3	Total	C	N	O	0	0	0
			39	22	2	15			
6	K	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 7 is an oligosaccharide called beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	I	6	Total	C	N	O	0	0	0
			72	40	2	30			
7	L	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	C	1	Total 14	8	1	5	0	0
8	D	1	Total 14	8	1	5	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ig alpha-1 chain C region




- Molecule 1: Ig alpha-1 chain C region





- Molecule 2: Immunoglobulin alpha Fc receptor







- Molecule 2: Immunoglobulin alpha Fc receptor




- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: alpha-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: alpha-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 33% 67%




- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 33% 67%




- Molecule 7: beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 17% 67% 17%



- Molecule 7: beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 17% 67% 17%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	142.99Å 142.99Å 67.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.69 – 3.10 29.69 – 3.06	Depositor EDS
% Data completeness (in resolution range)	95.2 (29.69-3.10) 93.7 (29.69-3.06)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.32 (at 3.06Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.252 , 0.284 0.249 , 0.281	Depositor DCC
$R_{free}$ test set	1300 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	83.2	Xtrriage
Anisotropy	0.637	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 68.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for -h,-k,l 0.458 for h,-h-k,-l 0.026 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6280	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	98.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, FUC, FUL, NAG, BMA, GAL, NDG, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1474	0.74	0/2041
1	B	0.45	0/1474	0.73	0/2041
2	C	0.47	0/1454	0.70	1/1989 (0.1%)
2	D	0.45	0/1454	0.70	1/1989 (0.1%)
All	All	0.46	0/5856	0.72	2/8060 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	PRO	N-CA-CB	6.07	110.59	103.30
2	D	6	PRO	N-CA-CB	6.04	110.55	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1435	0	1279	130	0
1	B	1435	0	1279	128	0
2	C	1414	0	1295	126	0
2	D	1414	0	1295	125	0
3	E	105	0	89	20	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	105	0	89	22	0
5	G	61	0	52	3	0
5	J	61	0	52	3	0
6	H	39	0	33	1	0
6	K	39	0	33	1	0
7	I	72	0	61	2	0
7	L	72	0	61	2	0
8	C	14	0	13	2	0
8	D	14	0	13	2	0
All	All	6280	0	5644	523	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 44.

The worst 5 of 523 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:345:PHE:H	1:B:375:SER:HB2	1.16	1.06
1:A:345:PHE:H	1:A:375:SER:HB2	1.14	1.05
1:A:345:PHE:H	1:A:375:SER:CB	1.78	0.96
1:B:345:PHE:H	1:B:375:SER:CB	1.80	0.94
1:B:391:PRO:HA	4:F:6:GAL:H62	1.48	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	207/214 (97%)	173 (84%)	22 (11%)	12 (6%)	<b>1</b> <b>10</b>
1	B	207/214 (97%)	174 (84%)	21 (10%)	12 (6%)	<b>1</b> <b>10</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	188/207 (91%)	143 (76%)	32 (17%)	13 (7%)	1	7
2	D	188/207 (91%)	143 (76%)	31 (16%)	14 (7%)	1	6
All	All	790/842 (94%)	633 (80%)	106 (13%)	51 (6%)	1	8

5 of 51 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	375	SER
1	A	442	ALA
1	B	375	SER
1	B	442	ALA
2	C	31	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/181 (74%)	119 (89%)	15 (11%)	6	24
1	B	134/181 (74%)	120 (90%)	14 (10%)	7	27
2	C	138/180 (77%)	129 (94%)	9 (6%)	17	47
2	D	138/180 (77%)	127 (92%)	11 (8%)	12	40
All	All	544/722 (75%)	495 (91%)	49 (9%)	9	34

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	447	THR
2	C	156	ASN
2	C	18	ILE
2	C	59	GLU
2	C	175	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	177	ASN
2	D	27	GLN
2	D	177	ASN
2	D	80	GLN
1	B	436	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

44 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	E	1	3	14,14,15	0.62	0	17,19,21	0.68	0
3	NAG	E	2	3	14,14,15	0.74	0	17,19,21	0.67	0
3	BMA	E	3	3	11,11,12	0.65	0	15,15,17	0.46	0
3	BMA	E	4	3	11,11,12	1.10	1 (9%)	15,15,17	0.88	1 (6%)
3	NAG	E	5	3	14,14,15	0.80	0	17,19,21	0.99	1 (5%)
3	GAL	E	6	3	11,11,12	0.79	0	15,15,17	0.73	0
3	SIA	E	7	3	20,20,21	0.85	1 (5%)	24,28,31	0.80	1 (4%)
3	FUL	E	8	3	10,10,11	0.58	0	14,14,16	0.74	0
4	NAG	F	1	4	14,14,15	0.61	0	17,19,21	0.69	0
4	NAG	F	2	4	14,14,15	0.75	0	17,19,21	0.68	0
4	BMA	F	3	4	11,11,12	0.63	0	15,15,17	0.48	0
4	BMA	F	4	4	11,11,12	1.10	1 (9%)	15,15,17	0.89	1 (6%)
4	NAG	F	5	4	14,14,15	0.78	0	17,19,21	0.99	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GAL	F	6	4	11,11,12	0.79	0	15,15,17	0.73	0
4	SIA	F	7	4	20,20,21	0.86	1 (5%)	24,28,31	0.81	1 (4%)
4	FUC	F	8	4	10,10,11	0.58	0	14,14,16	0.74	0
5	NAG	G	1	2,5	14,14,15	0.60	0	17,19,21	0.67	0
5	NAG	G	2	5	14,14,15	0.82	1 (7%)	17,19,21	0.59	0
5	MAN	G	3	5	11,11,12	0.75	0	15,15,17	0.97	1 (6%)
5	MAN	G	4	5	11,11,12	0.60	0	15,15,17	0.80	1 (6%)
5	BMA	G	5	5	11,11,12	0.53	0	15,15,17	0.54	0
6	NAG	H	1	6,2	14,14,15	0.74	0	17,19,21	0.69	0
6	NDG	H	2	6	14,14,15	0.86	1 (7%)	17,19,21	1.09	1 (5%)
6	BMA	H	3	6	11,11,12	0.52	0	15,15,17	0.43	0
7	NAG	I	1	2,7	14,14,15	0.85	1 (7%)	17,19,21	0.89	0
7	NAG	I	2	7	14,14,15	0.71	0	17,19,21	1.07	2 (11%)
7	BMA	I	3	7	11,11,12	0.61	0	15,15,17	0.40	0
7	MAN	I	4	7	11,11,12	0.86	0	15,15,17	0.79	1 (6%)
7	BMA	I	5	7	11,11,12	0.73	0	15,15,17	0.47	0
7	BMA	I	6	7	11,11,12	0.70	0	15,15,17	0.29	0
5	NAG	J	1	2,5	14,14,15	0.60	0	17,19,21	0.66	0
5	NAG	J	2	5	14,14,15	0.83	1 (7%)	17,19,21	0.59	0
5	MAN	J	3	5	11,11,12	0.76	0	15,15,17	0.96	1 (6%)
5	MAN	J	4	5	11,11,12	0.60	0	15,15,17	0.78	1 (6%)
5	BMA	J	5	5	11,11,12	0.59	0	15,15,17	0.53	0
6	NAG	K	1	6,2	14,14,15	0.76	0	17,19,21	0.69	0
6	NDG	K	2	6	14,14,15	0.87	1 (7%)	17,19,21	1.09	1 (5%)
6	BMA	K	3	6	11,11,12	0.52	0	15,15,17	0.42	0
7	NAG	L	1	2,7	14,14,15	0.84	1 (7%)	17,19,21	0.91	0
7	NAG	L	2	7	14,14,15	0.71	0	17,19,21	1.07	2 (11%)
7	BMA	L	3	7	11,11,12	0.61	0	15,15,17	0.41	0
7	MAN	L	4	7	11,11,12	0.86	0	15,15,17	0.78	1 (6%)
7	BMA	L	5	7	11,11,12	0.73	0	15,15,17	0.48	0
7	BMA	L	6	7	11,11,12	0.67	0	15,15,17	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3	-	4/6/23/26	0/1/1/1

Continued on next page...



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	2	3	-	3/6/23/26	0/1/1/1
3	BMA	E	3	3	-	2/2/19/22	0/1/1/1
3	BMA	E	4	3	-	1/2/19/22	0/1/1/1
3	NAG	E	5	3	-	4/6/23/26	0/1/1/1
3	GAL	E	6	3	-	1/2/19/22	0/1/1/1
3	SIA	E	7	3	-	9/18/34/38	0/1/1/1
3	FUL	E	8	3	-	-	0/1/1/1
4	NAG	F	1	4	-	5/6/23/26	0/1/1/1
4	NAG	F	2	4	-	3/6/23/26	0/1/1/1
4	BMA	F	3	4	-	2/2/19/22	0/1/1/1
4	BMA	F	4	4	-	1/2/19/22	0/1/1/1
4	NAG	F	5	4	-	4/6/23/26	0/1/1/1
4	GAL	F	6	4	-	1/2/19/22	0/1/1/1
4	SIA	F	7	4	-	9/18/34/38	0/1/1/1
4	FUC	F	8	4	1/1/4/5	-	0/1/1/1
5	NAG	G	1	2,5	-	3/6/23/26	0/1/1/1
5	NAG	G	2	5	-	4/6/23/26	0/1/1/1
5	MAN	G	3	5	-	2/2/19/22	0/1/1/1
5	MAN	G	4	5	-	2/2/19/22	0/1/1/1
5	BMA	G	5	5	-	0/2/19/22	0/1/1/1
6	NAG	H	1	6,2	-	2/6/23/26	0/1/1/1
6	NDG	H	2	6	-	4/6/23/26	0/1/1/1
6	BMA	H	3	6	-	2/2/19/22	0/1/1/1
7	NAG	I	1	2,7	-	3/6/23/26	0/1/1/1
7	NAG	I	2	7	-	4/6/23/26	0/1/1/1
7	BMA	I	3	7	-	2/2/19/22	0/1/1/1
7	MAN	I	4	7	-	0/2/19/22	1/1/1/1
7	BMA	I	5	7	-	2/2/19/22	0/1/1/1
7	BMA	I	6	7	-	2/2/19/22	0/1/1/1
5	NAG	J	1	2,5	-	3/6/23/26	0/1/1/1
5	NAG	J	2	5	-	4/6/23/26	0/1/1/1
5	MAN	J	3	5	-	2/2/19/22	0/1/1/1
5	MAN	J	4	5	-	2/2/19/22	0/1/1/1
5	BMA	J	5	5	-	0/2/19/22	0/1/1/1
6	NAG	K	1	6,2	-	2/6/23/26	0/1/1/1
6	NDG	K	2	6	-	4/6/23/26	0/1/1/1

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BMA	K	3	6	-	2/2/19/22	0/1/1/1
7	NAG	L	1	2,7	-	3/6/23/26	0/1/1/1
7	NAG	L	2	7	-	4/6/23/26	0/1/1/1
7	BMA	L	3	7	-	2/2/19/22	0/1/1/1
7	MAN	L	4	7	-	0/2/19/22	1/1/1/1
7	BMA	L	5	7	-	2/2/19/22	0/1/1/1
7	BMA	L	6	7	-	2/2/19/22	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	4	BMA	C2-C3	2.33	1.55	1.52
3	E	4	BMA	C2-C3	2.30	1.55	1.52
7	L	1	NAG	C1-C2	2.25	1.55	1.52
7	I	1	NAG	C1-C2	2.25	1.55	1.52
3	E	7	SIA	C2-C1	2.15	1.54	1.52

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	3	MAN	C6-C5-C4	2.95	119.92	113.00
5	J	3	MAN	C6-C5-C4	2.92	119.84	113.00
7	I	4	MAN	C1-O5-C5	2.67	115.81	112.19
7	L	4	MAN	C1-O5-C5	2.64	115.77	112.19
7	L	2	NAG	C4-C3-C2	-2.59	107.22	111.02

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	F	8	FUC	C1

5 of 113 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2
3	E	5	NAG	C8-C7-N2-C2


All (2) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	I	4	MAN	C1-C2-C3-C4-C5-O5
7	L	4	MAN	C1-C2-C3-C4-C5-O5


30 monomers are involved in 54 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	5	BMA	2	0
3	E	1	NAG	6	0
5	J	2	NAG	1	0
4	F	7	SIA	5	0
5	G	1	NAG	1	0
4	F	1	NAG	6	0
5	G	2	NAG	1	0
4	F	5	NAG	8	0
7	I	4	MAN	1	0
7	L	6	BMA	1	0
3	E	4	BMA	1	0
5	J	1	NAG	1	0
7	I	6	BMA	1	0
3	E	6	GAL	3	0
5	G	3	MAN	2	0
6	H	1	NAG	1	0
4	F	4	BMA	1	0
4	F	3	BMA	2	0
7	L	5	BMA	2	0
7	L	4	MAN	1	0
3	E	3	BMA	2	0
4	F	6	GAL	3	0
6	K	1	NAG	1	0
5	J	3	MAN	2	0
5	J	5	BMA	2	0
4	F	2	NAG	1	0
3	E	2	NAG	1	0
3	E	5	NAG	7	0
3	E	7	SIA	4	0
7	I	5	BMA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	NAG	C	710	2	14,14,15	0.57	0	17,19,21	0.60	0
8	NAG	D	810	2	14,14,15	0.59	0	17,19,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	C	710	2	-	3/6/23/26	0/1/1/1
8	NAG	D	810	2	-	3/6/23/26	0/1/1/1

There are no bond length outliers.



There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	C	710	NAG	C8-C7-N2-C2
8	C	710	NAG	O7-C7-N2-C2
8	D	810	NAG	C8-C7-N2-C2
8	D	810	NAG	O7-C7-N2-C2
8	C	710	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	C	710	NAG	2	0
8	D	810	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	209/214 (97%)	0.05	1 (0%) 91 81	41, 86, 142, 201	0
1	B	209/214 (97%)	0.03	1 (0%) 91 81	41, 86, 142, 201	0
2	C	190/207 (91%)	0.04	1 (0%) 91 81	54, 88, 128, 153	0
2	D	190/207 (91%)	0.03	1 (0%) 91 81	54, 88, 128, 153	0
All	All	798/842 (94%)	0.04	4 (0%) 91 81	41, 87, 140, 201	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	113	LEU	4.0
2	C	113	LEU	3.5
1	A	299	CYS	3.2
1	B	299	CYS	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	G	2	14/15	0.30	0.72	200,200,200,200	0
7	BMA	L	6	11/12	0.33	0.22	177,177,177,177	0
5	NAG	J	2	14/15	0.36	0.73	200,200,200,200	0


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	BMA	I	6	11/12	0.37	0.24	177,177,177,177	0
4	NAG	F	2	14/15	0.45	0.56	189,197,200,200	0
3	NAG	E	2	14/15	0.53	0.43	189,197,200,200	0
5	MAN	G	3	11/12	0.55	0.31	200,200,200,200	0
5	MAN	G	4	11/12	0.62	0.47	189,197,200,200	0
3	BMA	E	3	11/12	0.64	0.30	197,200,200,200	0
5	MAN	J	3	11/12	0.64	0.34	200,200,200,200	0
7	MAN	L	4	11/12	0.65	0.33	195,200,200,200	0
4	NAG	F	1	14/15	0.66	0.46	178,193,200,200	0
4	BMA	F	3	11/12	0.66	0.26	197,200,200,200	0
7	NAG	L	1	14/15	0.67	0.19	138,138,138,138	0
6	NAG	H	1	14/15	0.67	0.21	196,196,196,196	0
3	NAG	E	1	14/15	0.67	0.44	178,193,200,200	0
5	NAG	J	1	14/15	0.68	0.72	184,199,200,200	0
5	MAN	J	4	11/12	0.69	0.48	189,197,200,200	0
5	NAG	G	1	14/15	0.69	0.75	184,199,200,200	0
6	NAG	K	1	14/15	0.70	0.22	196,196,196,196	0
7	NAG	I	1	14/15	0.72	0.18	138,138,138,138	0
7	MAN	I	4	11/12	0.72	0.29	195,200,200,200	0
3	GAL	E	6	11/12	0.74	0.22	143,151,157,173	0
4	NAG	F	5	14/15	0.74	0.57	200,200,200,200	0
3	SIA	E	7	20/21	0.76	0.19	177,198,200,200	0
4	GAL	F	6	11/12	0.76	0.20	143,151,157,173	0
3	BMA	E	4	11/12	0.77	0.23	146,159,165,173	0
4	SIA	F	7	20/21	0.77	0.19	177,198,200,200	0
7	BMA	I	3	11/12	0.78	0.18	163,163,163,163	0
3	NAG	E	5	14/15	0.78	0.65	200,200,200,200	0
6	BMA	H	3	11/12	0.80	0.41	183,191,198,198	0
3	FUL	E	8	10/11	0.80	0.23	200,200,200,200	0
4	BMA	F	4	11/12	0.80	0.20	146,159,165,173	0
4	FUC	F	8	10/11	0.81	0.23	200,200,200,200	0
7	BMA	L	5	11/12	0.81	0.37	200,200,200,200	0
7	BMA	L	3	11/12	0.81	0.20	163,163,163,163	0
5	BMA	J	5	11/12	0.83	0.17	116,120,132,133	0
7	BMA	I	5	11/12	0.83	0.39	200,200,200,200	0
5	BMA	G	5	11/12	0.84	0.18	116,120,132,133	0
6	NDG	K	2	14/15	0.87	0.29	194,200,200,200	0
6	NDG	H	2	14/15	0.88	0.26	194,200,200,200	0
6	BMA	K	3	11/12	0.88	0.45	183,191,198,198	0
7	NAG	I	2	14/15	0.89	0.20	146,146,146,146	0
7	NAG	L	2	14/15	0.90	0.21	146,146,146,146	0




The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain J:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain K:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain L:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	NAG	D	810	14/15	0.82	0.12	151,162,167,167	0
8	NAG	C	710	14/15	0.84	0.13	151,162,167,167	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.