

# wwPDB X-ray Structure Validation Summary Report (i)

### Nov 5, 2023 – 04:16 AM EST

PDB ID : 6P02

Title: Crystal structure of Mtb aspartate decarboxylase, 6-Chlorine pyrazinoic acid

complex

Authors: Sun, Q.; Li, X.; Sacchettini, J.C.; TB Structural Genomics Consortium (TB-

SGC)

Deposited on : 2019-05-16

Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \text{Entries})$	Similar resolution $(\#\text{Entries, resolution range}(\mathring{A}))$			
$R_{free}$	130704	1377 (2.26-2.26)			
Clashscore	141614	1487 (2.26-2.26)			
Ramachandran outliers	138981	1449 (2.26-2.26)			
Sidechain outliers	138945	1450 (2.26-2.26)			
RSRZ outliers	127900	1356 (2.26-2.26)			

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	24	79%	21%
1	С	24	12%	21%
1	Е	24	83%	12% •
1	G	24	71%	29%



Continued from previous page...

Mol	Chain	$oxed{ \mathbf{Length} }$	Quality of chain	
1	I	24	12%	17%
1	K	24	92%	8%
1	M	24	83%	17%
1	О	24	8%	12%
1	Q	24	12%	8%
1	S	24	96%	•
1	U	24	88%	12%
1	W	24	79%	21%
2	В	123	63% 11%	26%
2	D	123	67% 6% •	26%
2	F	123	65% 9%	26%
2	Н	123	65% 9%	26%
2	J	123	67% 6% •	26%
2	L	123	64% 9% •	26%
2	N	123	66% 8%	26%
2	Р	123	63% 11% •	26%
2	R	123	71%	26%
2	Т	123	71%	26%
2	V	123	65% 9%	26%
2	X	123	66% 8%	26%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

$\mathbf{Mol}$	Type	Chain	$\operatorname{Res}$	Chirality	Geometry	Clashes	Electron density
3	NMJ	F	201	-	X	-	-



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 10543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Aspartate 1-decarboxylase beta chain.

Mol	Chain	Residues		Ato	ms			ZeroOcc	AltConf	Trace
1	A	24	Total	С	N	О	S	0	0	0
1	A	24	190	119	36	32	3	0	U	0
1	С	24	Total	С	N	О	S	0	0	0
1		24	190	119	36	32	3	0	U	0
1	Е	24	Total	С	N	О	S	0	0	0
1	שנ	24	190	119	36	32	3	0	U	
1	G	24	Total	С	N	О	S	0	0	0
1	G	24	190	119	36	32	3	0	U	U
1	I	24	Total	С	N	О	S	0	0	0
1	1	24	190	119	36	32	3	0	U	U
1	K	24	Total	С	N	Ο	S	0	0	0
1	11	24	190	119	36	32	3	0	O	U
1	M	24	Total	С	N	Ο	S	0	0	0
1	1/1	24	190	119	36	32	3	0	U	
1	О	24	Total	С	N	Ο	S	0	0	0
1	U	24	190	119	36	32	3	0	U	U
1	Q	24	Total	С	N	Ο	S	0	0	0
1	Q	24	190	119	36	32	3	0	U	U
1	S	24	Total	С	N	О	S	0	0	0
1	D	24	190	119	36	32	3	0	U	U
1	U	24	Total	С	N	О	S	0	0	0
_ 1		Δ≒	190	119	36	32	3		U	U
1	W	24	Total	С	N	О	S	0	0	0
1	٧٧	24	190	119	36	32	3		U	U

• Molecule 2 is a protein called Aspartate 1-decarboxylase alpha chain.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
2	D	91	Total	С	N	О	S	0	0	0
2 D	91	678	427	115	134	2	0	U		
2	D	91	Total	С	N	О	S	0	0	0
	ע	91	678	427	115	134	2	0	U	U



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	F	91	Total	С	N	О	S	0	0	0
2	I.	91	678	427	115	134	2	0		U
2	Н	91	Total	С	N	О	S	0	0	0
	11	91	678	427	115	134	2	0	U	
2	J	91	Total	С	N	Ο	S	0	0	0
		91	678	427	115	134	2	0	U	
2	L	91	Total	С	N	О	S	0	0	0
	П	91	678	427	115	134	2	0	U	
2	N	91	Total	С	N	О	S	0	0	0
	11	91	678	427	115	134	2		0	
2	Р	91	Total	С	N	О	S	0	0	0
	1	91	678	427	115	134	2	0	U	
2	R	91	Total	С	N	О	S	0	0	0
	16	91	678	427	115	134	2	0	U	U
2	Т	91	Total	С	N	О	S	0	0	0
	1	91	678	427	115	134	2	0	U	U
2	V	91	Total	С	N	О	S	0	0	0
<u> </u>	v	91	678	427	115	134	2		U	
2	X	91	Total	С	N	О	S	0	0	0
	Λ	91	678	427	115	134	2	U	U	U

There are 108 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	25	PYR	SER	conflict	UNP P9WIL3
В	140	LEU	-	expression tag	UNP P9WIL3
В	141	GLU	-	expression tag	UNP P9WIL3
В	142	HIS	-	expression tag	UNP P9WIL3
В	143	HIS	-	expression tag	UNP P9WIL3
В	144	HIS	-	expression tag	UNP P9WIL3
В	145	HIS	-	expression tag	UNP P9WIL3
В	146	HIS	-	expression tag	UNP P9WIL3
В	147	HIS	-	expression tag	UNP P9WIL3
D	25	PYR	SER	conflict	UNP P9WIL3
D	140	LEU	-	expression tag	UNP P9WIL3
D	141	GLU	-	expression tag	UNP P9WIL3
D	142	HIS	-	expression tag	UNP P9WIL3
D	143	HIS	-	expression tag	UNP P9WIL3
D	144	HIS	-	expression tag	UNP P9WIL3
D	145	HIS	-	expression tag	UNP P9WIL3
D	146	HIS	-	expression tag	UNP P9WIL3
D	147	HIS	-	expression tag	UNP P9WIL3
F	25	PYR	SER	conflict	UNP P9WIL3



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
F	140	LEU	-	expression tag	UNP P9WIL3
F	141	GLU	_	expression tag	UNP P9WIL3
F	142	HIS	-	expression tag	UNP P9WIL3
F	143	HIS	-	expression tag	UNP P9WIL3
F	144	HIS	-	expression tag	UNP P9WIL3
F	145	HIS	_	expression tag	UNP P9WIL3
F	146	HIS	_	expression tag	UNP P9WIL3
F	147	HIS	-	expression tag	UNP P9WIL3
Н	25	PYR	SER	conflict	UNP P9WIL3
Н	140	LEU	-	expression tag	UNP P9WIL3
Н	141	GLU	-	expression tag	UNP P9WIL3
Н	142	HIS	-	expression tag	UNP P9WIL3
Н	143	HIS	-	expression tag	UNP P9WIL3
Н	144	HIS	-	expression tag	UNP P9WIL3
Н	145	HIS	-	expression tag	UNP P9WIL3
Н	146	HIS	_	expression tag	UNP P9WIL3
Н	147	HIS	-	expression tag	UNP P9WIL3
J	25	PYR	SER	conflict	UNP P9WIL3
J	140	LEU	-	expression tag	UNP P9WIL3
J	141	GLU	-	expression tag	UNP P9WIL3
J	142	HIS	-	expression tag	UNP P9WIL3
J	143	HIS	-	expression tag	UNP P9WIL3
J	144	HIS	-	expression tag	UNP P9WIL3
J	145	HIS	-	expression tag	UNP P9WIL3
J	146	HIS	-	expression tag	UNP P9WIL3
J	147	HIS	-	expression tag	UNP P9WIL3
L	25	PYR	SER	conflict	UNP P9WIL3
L	140	LEU	-	expression tag	UNP P9WIL3
L	141	GLU	-	expression tag	UNP P9WIL3
L	142	HIS	-	expression tag	UNP P9WIL3
L	143	HIS	-	expression tag	UNP P9WIL3
L	144	HIS	-	expression tag	UNP P9WIL3
L	145	HIS	-	expression tag	UNP P9WIL3
L	146	HIS	-	expression tag	UNP P9WIL3
L	147	HIS	-	expression tag	UNP P9WIL3
N	25	PYR	SER	conflict	UNP P9WIL3
N	140	LEU		expression tag	UNP P9WIL3
N	141	GLU		expression tag	UNP P9WIL3
N	142	HIS		expression tag	UNP P9WIL3
N	143	HIS		expression tag	UNP P9WIL3
N	144	HIS	-	expression tag	UNP P9WIL3
N	145	HIS	_	expression tag	UNP P9WIL3



 $Continued\ from\ previous\ page...$ 


Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
N	146	HIS	-	expression tag	UNP P9WIL3
N	147	HIS	-	expression tag	UNP P9WIL3
Р	25	PYR	SER	conflict	UNP P9WIL3
Р	140	LEU	-	expression tag	UNP P9WIL3
Р	141	GLU	-	expression tag	UNP P9WIL3
Р	142	HIS	-	expression tag	UNP P9WIL3
Р	143	HIS	-	expression tag	UNP P9WIL3
Р	144	HIS	-	expression tag	UNP P9WIL3
Р	145	HIS	-	expression tag	UNP P9WIL3
Р	146	HIS	-	expression tag	UNP P9WIL3
Р	147	HIS	-	expression tag	UNP P9WIL3
R	25	PYR	SER	conflict	UNP P9WIL3
R	140	LEU	-	expression tag	UNP P9WIL3
R	141	GLU	-	expression tag	UNP P9WIL3
R	142	HIS	-	expression tag	UNP P9WIL3
R	143	HIS	-	expression tag	UNP P9WIL3
R	144	HIS	-	expression tag	UNP P9WIL3
R	145	HIS	-	expression tag	UNP P9WIL3
R	146	HIS	-	expression tag	UNP P9WIL3
R	147	HIS	-	expression tag	UNP P9WIL3
Т	25	PYR	SER	conflict	UNP P9WIL3
Т	140	LEU	-	expression tag	UNP P9WIL3
Т	141	GLU	-	expression tag	UNP P9WIL3
Т	142	HIS	-	expression tag	UNP P9WIL3
Т	143	HIS	-	expression tag	UNP P9WIL3
Т	144	HIS	-	expression tag	UNP P9WIL3
Т	145	HIS	-	expression tag	UNP P9WIL3
Т	146	HIS	-	expression tag	UNP P9WIL3
Т	147	HIS	-	expression tag	UNP P9WIL3
V	25	PYR	SER	conflict	UNP P9WIL3
V	140	LEU	-	expression tag	UNP P9WIL3
V	141	GLU		expression tag	UNP P9WIL3
V	142	HIS	_	expression tag	UNP P9WIL3
V	143	HIS	_	expression tag	UNP P9WIL3
V	144	HIS	_	expression tag	UNP P9WIL3
V	145	HIS	_	expression tag	UNP P9WIL3
V	146	HIS	_	expression tag	UNP P9WIL3
V	147	HIS	-	expression tag	UNP P9WIL3
X	25	PYR	SER	conflict	UNP P9WIL3
X	140	LEU	-	expression tag	UNP P9WIL3
X	141	GLU	-	expression tag	UNP P9WIL3
X	142	HIS	_	expression tag	UNP P9WIL3



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
X	143	HIS	-	expression tag	UNP P9WIL3
X	144	HIS	-	expression tag	UNP P9WIL3
X	145	HIS	-	expression tag	UNP P9WIL3
X	146	HIS	-	expression tag	UNP P9WIL3
X	147	HIS	-	expression tag	UNP P9WIL3

• Molecule 3 is 6-chloropyrazine-2-carboxylic acid (three-letter code: NMJ) (formula:  $C_5H_3ClN_2O_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	
3	В	1	Total	С	Cl	N	О	0	0	
3	Б	1	10	5	1	2	2	0	U	
3	D	1	Total	С	Cl	N	О	0	0	
J		D	1	10	5	1	2	2	U	U
3	F	1	Total	С	$\operatorname{Cl}$	N	Ο	0	0	
	I.	1	10	5	1	2	2	O	U	
3	H	1	Total	С	$\operatorname{Cl}$	N	Ο	0	0	
	11	1	10	5	1	2	2	0		
3	Ţ	J	1	Total	С	Cl	N	Ο	0	0
	3	1	10	5	1	2	2	0	O	
3	L	1	Total	С	Cl	N	Ο	0	0	
	L	1	10	5	1	2	2	0	0	
3	N	1	Total	С	Cl	N	Ο	0	0	
	11	1N 1	10	5	1	2	2	0	U	
3	Р	1	Total	С	$\operatorname{Cl}$	N	Ο	0	0	
	P	I I	10	5	1	2	2		0	



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	R	1	Total C Cl N O	0	0	
)	$\mathbf{n}$	1	10  5  1  2  2	0		
3	Т	1	Total C Cl N O	0	0	
3	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	1	10  5  1  2  2	0		
2	V	1	Total C Cl N O	0	0	
3 V	1	10  5  1  2  2	0	U		
2	V	V 1	Total C Cl N O	0	0	
3	$\Lambda$	1	10  5  1  2  2		U	

### • Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	2	Total O 2 2	0	0
4	D	2	Total O 2 2	0	0
4	N	1	Total O 1 1	0	0
4	О	1	Total O 1 1	0	0
4	R	1	Total O 1 1	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aspartate 1-decarboxylase beta chain




• Molecule 1: Aspartate 1-decarboxylase beta chain











• Molecule 2: Aspartate 1-decarboxylase alpha chain Chain P: 26% 11% • Molecule 2: Aspartate 1-decarboxylase alpha chain Chain R: 26% • Molecule 2: Aspartate 1-decarboxylase alpha chain Chain T: 71% 26% • Molecule 2: Aspartate 1-decarboxylase alpha chain Chain V: 65% 9% 26% • Molecule 2: Aspartate 1-decarboxylase alpha chain Chain X: 66% 26%



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	143.68Å 143.68Å 59.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	21.89 - 2.25 $21.89 - 2.25$	Depositor EDS
% Data completeness	99.5 (21.89-2.25)	Depositor
(in resolution range)	99.6 (21.89-2.25)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.52 (at 2.26Å)	Xtriage
Refinement program	REFMAC 5.8.0232	Depositor
D D.	0.134 , 0.164	Depositor
$R, R_{free}$	0.135 , $0.162$	DCC
$R_{free}$ test set	3466 reflections (5.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.1	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.35, 24.3	EDS
L-test for twinning <sup>2</sup>	$< L >=0.35, < L^2>=0.17$	Xtriage
Estimated twinning fraction	0.408 for -h,-k,l 0.217 for h,-h-k,-l 0.216 for -k,-h,-l	Xtriage
Reported twinning fraction	0.086 for H, K, L 0.074 for -h,-k,l 0.483 for K, H, -L 0.357 for -K, -H, -L	Depositor
Outliers	0 of 65249 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10543	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.12% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NMJ, PYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Chair		Bond	lengths	Bond angles		
Mol	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.70	0/192	0.94	0/255	
1	С	0.72	0/192	0.85	0/255	
1	Е	0.71	0/192	1.01	1/255~(0.4%)	
1	G	0.66	0/192	0.89	0/255	
1	I	0.74	0/192	0.87	0/255	
1	K	0.74	0/192	0.92	0/255	
1	M	0.66	0/192	0.93	0/255	
1	О	0.78	0/192	0.93	0/255	
1	Q	0.72	0/192	0.86	0/255	
1	S	0.71	0/192	0.88	0/255	
1	U	0.67	0/192	0.90	0/255	
1	W	0.74	0/192	0.92	0/255	
2	В	0.72	0/682	0.82	0/931	
2	D	0.73	0/682	0.83	0/931	
2	F	0.72	0/682	0.90	1/931 (0.1%)	
2	Н	0.71	0/682	0.78	0/931	
2	J	0.75	0/682	0.85	0/931	
2	L	0.72	0/682	0.83	1/931 (0.1%)	
2	N	0.73	0/682	0.85	0/931	
2	Р	0.70	0/682	0.81	0/931	
2	R	0.67	0/682	0.80	0/931	
2	Т	0.70	0/682	0.81	0/931	
2	V	0.67	0/682	0.80	0/931	
2	X	0.68	0/682	0.79	0/931	
All	All	0.71	0/10488	0.84	3/14232 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.



Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	1
2	F	0	1
2	Н	0	1
2	J	0	1
2	L	0	1
2	Р	0	1
2	R	0	1
2	V	0	1
2	X	0	1
All	All	0	9

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	L	104	ARG	CG-CD-NE	-6.27	98.64	111.80
2	F	50	ASP	CB-CG-OD2	-6.05	112.85	118.30
1	Е	1	MET	CG-SD-CE	5.95	109.71	100.20

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	25	PYR	Mainchain
2	F	25	PYR	Mainchain
2	Н	25	PYR	Mainchain
2	J	25	PYR	Mainchain
2	L	25	PYR	Mainchain

# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	190	0	205	6	0
1	С	190	0	205	10	0
1	Е	190	0	205	4	0
1	G	190	0	205	8	0



Continued from previous page...

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
1	I	190	0	205	9	0
1	K	190	0	205	4	0
1	M	190	0	205	5	0
1	О	190	0	205	5	0
1	Q	190	0	205	3	0
1	S	190	0	205	1	0
1	U	190	0	205	3	0
1	W	190	0	205	5	0
2	В	678	0	672	25	0
2	D	678	0	672	14	0
2	F	678	0	672	10	1
2	Н	678	0	672	15	0
2	J	678	0	672	14	0
2	L	678	0	672	19	1
2	N	678	0	672	13	0
2	Р	678	0	672	13	0
2	R	678	0	672	6	0
2	Τ	678	0	672	2	0
2	V	678	0	672	14	0
2	X	678	0	672	13	0
3	В	10	0	0	1	0
3	D	10	0	0	0	0
3	F	10	0	0	0	0
3	Н	10	0	0	1	0
3	J	10	0	0	1	0
3	L	10	0	0	0	0
3	N	10	0	0	0	0
3	Р	10	0	0	0	0
3	R	10	0	0	0	0
3	Т	10	0	0	0	0
3	V	10	0	0	2	0
3	X	10	0	0	0	0
4	В	2	0	0	0	0
4	D	2	0	0	0	0
4	N	1	0	0	0	0
4	О	1	0	0	0	0
4	R	1	0	0	0	0
All	All	10543	0	10524	130	1

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 6.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:7:LYS:HE2	2:P:42:GLU:OE2	1.54	1.08
1:Q:7:LYS:HE2	2:R:42:GLU:OE2	1.62	0.96
1:E:7:LYS:HE2	2:F:42:GLU:OE1	1.68	0.93
2:B:94:ASP:OD2	2:L:64:ARG:NH1	2.01	0.93
1:G:7:LYS:HE2	2:H:42:GLU:OE2	1.71	0.90

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$	
2:F:99:ARG:O	2:L:99:ARG:NH2[1_554]	2.07	0.13	

# 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$22/24\ (92\%)$	22 (100%)	0	0	100	100
1	С	22/24~(92%)	22 (100%)	0	0	100	100
1	E	22/24~(92%)	21 (96%)	1 (4%)	0	100	100
1	G	22/24~(92%)	21 (96%)	1 (4%)	0	100	100
1	I	22/24 (92%)	22 (100%)	0	0	100	100
1	K	22/24 (92%)	22 (100%)	0	0	100	100
1	M	22/24 (92%)	22 (100%)	0	0	100	100
1	О	22/24 (92%)	22 (100%)	0	0	100	100
1	Q	22/24 (92%)	22 (100%)	0	0	100	100
1	S	22/24 (92%)	22 (100%)	0	0	100	100
1	U	22/24 (92%)	22 (100%)	0	0	100	100
1	W	22/24 (92%)	22 (100%)	0	0	100	100
2	В	89/123 (72%)	89 (100%)	0	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
2	D	89/123 (72%)	89 (100%)	0	0	100	100
2	F	89/123 (72%)	89 (100%)	0	0	100	100
2	Н	89/123 (72%)	89 (100%)	0	0	100	100
2	J	89/123 (72%)	89 (100%)	0	0	100	100
2	L	89/123 (72%)	89 (100%)	0	0	100	100
2	N	89/123 (72%)	88 (99%)	1 (1%)	0	100	100
2	Р	89/123 (72%)	89 (100%)	0	0	100	100
2	R	89/123 (72%)	89 (100%)	0	0	100	100
2	Т	89/123 (72%)	89 (100%)	0	0	100	100
2	V	89/123 (72%)	89 (100%)	0	0	100	100
2	X	89/123 (72%)	89 (100%)	0	0	100	100
All	All	1332/1764 (76%)	1329 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	21/21~(100%)	21 (100%)	0	100	100
1	C	21/21~(100%)	21 (100%)	0	100	100
1	E	21/21~(100%)	19 (90%)	2 (10%)	8	6
1	G	21/21~(100%)	21 (100%)	0	100	100
1	I	$21/21\ (100\%)$	21 (100%)	0	100	100
1	K	21/21~(100%)	21 (100%)	0	100	100
1	M	21/21~(100%)	21 (100%)	0	100	100
1	О	21/21~(100%)	21 (100%)	0	100	100
1	Q	21/21 (100%)	21 (100%)	0	100	100
1	S	$21/21\ (100\%)$	21 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	U	21/21 (100%)	20 (95%)	1 (5%)	25	28
1	W	21/21 (100%)	21 (100%)	0	100	100
2	В	69/95~(73%)	69 (100%)	0	100	100
2	D	69/95~(73%)	68 (99%)	1 (1%)	67	76
2	F	69/95 (73%)	69 (100%)	0	100	100
2	Н	69/95 (73%)	68 (99%)	1 (1%)	67	76
2	J	69/95 (73%)	69 (100%)	0	100	100
2	L	69/95~(73%)	69 (100%)	0	100	100
2	N	69/95 (73%)	68 (99%)	1 (1%)	67	76
2	Р	69/95~(73%)	67 (97%)	2 (3%)	42	51
2	R	69/95 (73%)	69 (100%)	0	100	100
2	Т	69/95~(73%)	68 (99%)	1 (1%)	67	76
2	V	69/95 (73%)	69 (100%)	0	100	100
2	X	69/95 (73%)	69 (100%)	0	100	100
All	All	1080/1392 (78%)	1071 (99%)	9 (1%)	81	88

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	Т	94	ASP
1	U	12	ARG
2	Н	49	ILE
2	N	94	ASP
2	Р	94	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	21	HIS
1	K	21	HIS
1	S	21	HIS

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Вс	ond leng	ths	Bond angles		
WIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NMJ	R	201	-	10,10,10	1.19	1 (10%)	13,13,13	2.95	6 (46%)
3	NMJ	Р	201	-	10,10,10	1.13	1 (10%)	13,13,13	2.49	4 (30%)
3	NMJ	F	201	-	10,10,10	2.32	3 (30%)	13,13,13	3.90	9 (69%)
3	NMJ	X	201	ı	10,10,10	1.75	2 (20%)	13,13,13	1.75	2 (15%)
3	NMJ	D	201	ı	10,10,10	1.09	0	13,13,13	2.09	3 (23%)
3	NMJ	В	201	ı	10,10,10	1.57	2 (20%)	13,13,13	2.59	5 (38%)
3	NMJ	V	201	-	10,10,10	0.93	0	13,13,13	1.76	5 (38%)
3	NMJ	Н	201	-	10,10,10	1.25	2 (20%)	13,13,13	3.10	4 (30%)
3	NMJ	J	201	-	10,10,10	1.00	1 (10%)	13,13,13	1.95	6 (46%)
3	NMJ	L	201	-	10,10,10	1.04	0	13,13,13	2.19	5 (38%)
3	NMJ	N	201	-	10,10,10	1.30	2 (20%)	13,13,13	2.69	5 (38%)
3	NMJ	Т	201	-	10,10,10	0.92	0	13,13,13	2.10	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NMJ	R	201	-	-	4/4/4/4	0/1/1/1
3	NMJ	Р	201	-	-	4/4/4/4	0/1/1/1
3	NMJ	F	201	-	-	3/4/4/4	0/1/1/1
3	NMJ	X	201	-	-	1/4/4/4	0/1/1/1
3	NMJ	D	201	-	-	4/4/4/4	0/1/1/1
3	NMJ	В	201	-	-	0/4/4/4	0/1/1/1
3	NMJ	V	201	-	-	3/4/4/4	0/1/1/1
3	NMJ	Н	201	-	-	0/4/4/4	0/1/1/1
3	NMJ	J	201	-	-	3/4/4/4	0/1/1/1
3	NMJ	L	201	-	-	2/4/4/4	0/1/1/1
3	NMJ	N	201	-	-	0/4/4/4	0/1/1/1
3	NMJ	Т	201	-	-	2/4/4/4	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	Ideal(A)
3	F	201	NMJ	C02-C08	-6.14	1.42	1.50
3	X	201	NMJ	C02-C08	-3.75	1.45	1.50
3	В	201	NMJ	C02-C08	-3.42	1.46	1.50
3	N	201	NMJ	O10-C08	-2.91	1.21	1.30
3	X	201	NMJ	C02-N03	-2.56	1.30	1.34

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	Н	201	NMJ	C05-C04-CL07	9.08	124.76	119.52
3	N	201	NMJ	C04-C05-N06	-7.31	117.49	120.43
3	F	201	NMJ	C01-C02-N03	6.33	130.86	120.77
3	В	201	NMJ	C04-C05-N06	6.20	122.93	120.43
3	F	201	NMJ	C05-C04-CL07	6.02	122.99	119.52

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	201	NMJ	N03-C02-C08-O10
3	J	201	NMJ	C01-C02-C08-O10
3	J	201	NMJ	N03-C02-C08-O10
3	Р	201	NMJ	C01-C02-C08-O10
3	Р	201	NMJ	N03-C02-C08-O10




There are no ring outliers.


4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	201	NMJ	1	0
3	V	201	NMJ	2	0
3	Н	201	NMJ	1	0
3	J	201	NMJ	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























































# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	A	24/24~(100%)	0.12	2 (8%) 11 12	38, 51, 98, 123	0
1	C	24/24~(100%)	0.15	3 (12%) 3 3	38, 47, 124, 136	0
1	E	24/24~(100%)	-0.35	1 (4%) 36 38	39, 46, 65, 79	0
1	G	24/24~(100%)	0.32	2 (8%) 11 12	41, 60, 108, 148	0
1	I	24/24~(100%)	0.09	3 (12%) 3 3	39, 49, 115, 127	0
1	K	$24/24\ (100\%)$	-0.12	1 (4%) 36 38	39, 52, 103, 105	0
1	M	24/24~(100%)	-0.33	0 100 100	37, 48, 63, 92	0
1	О	$24/24\ (100\%)$	0.30	2 (8%) 11 12	38, 52, 120, 151	0
1	Q	24/24~(100%)	0.15	3 (12%) 3 3	47, 55, 131, 158	0
1	S	$24/24\ (100\%)$	-0.03	1 (4%) 36 38	46, 53, 75, 100	0
1	U	24/24~(100%)	0.83	4 (16%) 1 1	47, 62, 137, 154	0
1	W	$24/24\ (100\%)$	-0.18	2 (8%) 11 12	40, 51, 79, 108	0
2	В	90/123~(73%)	-0.35	0 100 100	34, 48, 66, 77	0
2	D	90/123~(73%)	-0.45	0 100 100	34, 46, 69, 80	0
2	F	90/123~(73%)	-0.47	0 100 100	37, 49, 65, 83	0
2	Н	90/123~(73%)	-0.37	0 100 100	38, 52, 72, 96	0
2	J	90/123~(73%)	-0.41	0 100 100	35, 47, 69, 78	0
2	L	90/123~(73%)	-0.42	0 100 100	36, 51, 76, 89	0
2	N	90/123 (73%)	-0.38	0 100 100	32, 46, 72, 83	0
2	Р	90/123~(73%)	-0.51	0 100 100	37, 52, 68, 77	0
2	R	90/123 (73%)	-0.33	1 (1%) 80 82	40, 54, 71, 94	0
2	Т	90/123 (73%)	-0.37	0 100 100	41, 54, 80, 89	0
2	V	90/123 (73%)	-0.35	0 100 100	40, 58, 78, 83	0
2	X	90/123~(73%)	-0.51	0 100 100	39, 49, 74, 83	0



Continued from previous page...

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\# \mathrm{RSRZ}{>}2$		$OWAB(Å^2)$	Q<0.9	
All	All	1368/1764 (77%)	-0.31	25 (1%)	68	71	32, 51, 79, 158	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	U	24	GLY	9.9
1	U	23	VAL	9.0
1	O	23	VAL	8.3
1	G	22	TYR	6.9
1	G	23	VAL	6.5

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


# 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	NMJ	R	201	10/10	0.89	0.13	50,64,68,69	0
3	NMJ	Τ	201	10/10	0.91	0.12	53,58,63,72	0
3	NMJ	X	201	10/10	0.92	0.17	48,58,70,74	0
3	NMJ	F	201	10/10	0.94	0.10	37,53,62,69	0
3	NMJ	V	201	10/10	0.95	0.12	51,61,65,87	0
3	NMJ	D	201	10/10	0.96	0.11	42,52,60,62	0
3	NMJ	J	201	10/10	0.96	0.11	38,56,62,63	0
3	NMJ	N	201	10/10	0.97	0.08	35,60,65,69	0
3	NMJ	Р	201	10/10	0.97	0.11	46,55,63,64	0
3	NMJ	В	201	10/10	0.97	0.12	47,56,58,74	0
3	NMJ	Н	201	10/10	0.98	0.14	51,56,62,66	0
3	NMJ	L	201	10/10	0.98	0.11	41,51,65,70	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







### Electron density around NMJ T 201:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around NMJ X 201:


 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)















# Electron density around NMJ V 201: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)




# Electron density around NMJ D 201:

 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around NMJ J 201:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)





















### Electron density around NMJ H 201:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around NMJ L 201:

 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









# 6.5 Other polymers (i)

There are no such residues in this entry.

