



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2022 – 12:07 pm GMT

PDB ID : 7P0T  
Title : CRYSTAL STRUCTURE OF THE MURINE CLASS I MAJOR HISTOCOMPATIBILITY COMPLEX H-2DB IN COMPLEX WITH LCMV-DERIVED GP33 PEPTIDE with D-AMINOACID  
Authors : Brogini, L.; Ricagno, S.  
Deposited on : 2021-06-30  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 5% 73% 14% 11%
1	D	276	 7% 70% 22% 7%
2	B	101	 % 72% 26%
2	E	101	 81% 14%
3	C	9	 67% 33%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	9	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '11%', a green segment in the middle labeled '56%', and a yellow segment on the right labeled '44%'. The segments are stacked horizontally, with the red segment starting from the left edge of the bar.</p>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, D-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	247	Total	C	N	O	S	0	2	0
			1959	1233	356	362	8			
1	D	258	Total	C	N	O	S	0	0	0
			1929	1228	334	358	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	Total	C	N	O	S	0	0	0
			770	493	127	144	6			
2	E	99	Total	C	N	O	S	0	0	0
			785	503	135	140	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	MET	-	initiating methionine	UNP P01887
B	0	GLY	-	expression tag	UNP P01887
B	85	ASP	ALA	conflict	UNP P01887
E	-1	MET	-	initiating methionine	UNP P01887
E	0	GLY	-	expression tag	UNP P01887
E	85	ASP	ALA	conflict	UNP P01887

- Molecule 3 is a protein called Derived peptide gp33-41 from LCMV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	9	Total	C	N	O	S	0	0	0
			73	48	11	13	1			
3	F	9	Total	C	N	O	S	0	0	0
			73	48	11	13	1			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0


- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	42	Total 42	O 42	0	0
6	B	15	Total 15	O 15	0	0
6	C	2	Total 2	O 2	0	0
6	D	6	Total 6	O 6	0	0
6	E	8	Total 8	O 8	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin

Chain E:  81% 14% . .



- Molecule 3: Derived peptide gp33-41 from LCMV

Chain C:  67% 33%



- Molecule 3: Derived peptide gp33-41 from LCMV

Chain F:  11% 56% 44%





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.51Å 124.60Å 92.92Å 90.00° 126.87° 90.00°	Depositor
Resolution (Å)	48.20 – 2.60 48.20 – 2.61	Depositor EDS
% Data completeness (in resolution range)	66.2 (48.20-2.60) 66.2 (48.20-2.61)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.72 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, $R_{free}$	0.197 , 0.244 0.199 , 0.243	Depositor DCC
$R_{free}$ test set	2017 reflections (9.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.5	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5703	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, DPN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2014	0.61	2/2736 (0.1%)
1	D	0.28	0/1988	0.51	0/2722
2	B	0.30	0/796	0.55	0/1091
2	E	0.32	0/811	0.59	1/1105 (0.1%)
3	C	0.34	0/61	0.58	0/78
3	F	0.28	0/61	0.47	0/78
All	All	0.31	0/5731	0.56	3/7810 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	193	PRO	CB-CA-C	10.71	138.76	112.00
1	A	193	PRO	C-N-CA	6.45	137.82	121.70
2	E	31	HIS	CB-CA-C	5.24	120.88	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	193	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1959	0	1749	35	0
1	D	1929	0	1639	53	0
2	B	770	0	692	22	0
2	E	785	0	736	14	0
3	C	73	0	73	6	0
3	F	73	0	73	9	0
4	A	15	0	0	0	0
4	B	10	0	0	0	0
4	D	5	0	0	0	0
4	E	10	0	0	0	0
5	A	1	0	0	0	0
6	A	42	0	0	1	0
6	B	15	0	0	1	0
6	C	2	0	0	0	0
6	D	6	0	0	0	0
6	E	8	0	0	0	0
All	All	5703	0	4962	116	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 116 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:73:THR:HB	2:B:76:ASP:HB2	1.68	0.75
2:B:1:ILE:N	6:B:201:HOH:O	2.18	0.74
1:A:62:ARG:HH11	3:C:1:LYS:HE2	1.55	0.72
2:B:4:THR:HG22	2:B:86:SER:HB2	1.72	0.70
1:D:199:VAL:C	1:D:248:VAL:HG23	2.12	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/276 (87%)	229 (96%)	8 (3%)	2 (1%)	19	39
1	D	250/276 (91%)	241 (96%)	9 (4%)	0	100	100
2	B	97/101 (96%)	90 (93%)	6 (6%)	1 (1%)	15	32
2	E	97/101 (96%)	94 (97%)	2 (2%)	1 (1%)	15	32
3	C	6/9 (67%)	5 (83%)	1 (17%)	0	100	100
3	F	6/9 (67%)	6 (100%)	0	0	100	100
All	All	695/772 (90%)	665 (96%)	26 (4%)	4 (1%)	25	47

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	97	ARG
2	E	97	ARG
1	A	41	GLU
1	A	193	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/234 (79%)	178 (96%)	7 (4%)	33	59
1	D	168/234 (72%)	164 (98%)	4 (2%)	49	74
2	B	81/95 (85%)	81 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	84/95 (88%)	81 (96%)	3 (4%)	35	61
3	C	6/6 (100%)	6 (100%)	0	100	100
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	530/670 (79%)	516 (97%)	14 (3%)	44	72

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	38	SER
1	D	45	TYR
2	E	99	MET
2	E	44	LYS
2	E	52	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	F	5	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	A	301	-	4,4,4	0.15	0	6,6,6	0.06	0
4	SO4	D	301	-	4,4,4	0.14	0	6,6,6	0.04	0
4	SO4	E	101	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	A	302	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	A	303	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	B	101	-	4,4,4	0.15	0	6,6,6	0.06	0
4	SO4	B	102	-	4,4,4	0.15	0	6,6,6	0.05	0
4	SO4	E	102	-	4,4,4	0.13	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	247/276 (89%)	0.12	15 (6%) 21 16	20, 68, 146, 222	0
1	D	258/276 (93%)	0.11	20 (7%) 13 9	50, 102, 148, 174	0
2	B	99/101 (98%)	-0.13	1 (1%) 82 80	33, 75, 120, 143	0
2	E	99/101 (98%)	-0.35	0 100 100	34, 67, 102, 113	0
3	C	8/9 (88%)	0.06	0 100 100	33, 42, 71, 84	0
3	F	8/9 (88%)	0.45	1 (12%) 3 2	93, 101, 120, 126	0
All	All	719/772 (93%)	0.02	37 (5%) 28 22	20, 81, 143, 222	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	4	TYR	4.1
1	D	197	GLY	4.1
1	D	274	TRP	3.8
1	D	191	HIS	3.3
1	D	256	ASN	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	DPN	F	6	11/12	0.90	0.25	95,103,120,122	0
3	DPN	C	6	11/12	0.97	0.17	31,46,68,78	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	A	302	5/5	0.87	0.14	126,133,134,137	0
4	SO4	A	303	5/5	0.88	0.18	142,151,152,152	0
4	SO4	E	102	5/5	0.89	0.16	138,140,140,141	0
5	CL	A	304	1/1	0.89	0.31	73,73,73,73	0
4	SO4	E	101	5/5	0.90	0.18	127,132,138,145	0
4	SO4	B	101	5/5	0.91	0.13	114,118,121,123	0
4	SO4	B	102	5/5	0.93	0.09	147,151,153,155	0
4	SO4	A	301	5/5	0.94	0.11	107,122,126,130	0
4	SO4	D	301	5/5	0.95	0.05	145,147,149,149	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.