

wwPDB X-ray Structure Validation Summary Report (i)

Jan 11, 2022 - 09:16 pm GMT

PDB ID	:	7PUH
Title	:	Crystal Structure of Two-Domain Laccase mutant $H165A/R240H$ from Strep-
		tomyces griseoflavus
Authors	:	Kolyadenko, I.; Tishchenko, S.; Gabdulkhakov, A.
Deposited on		
Resolution	:	1.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity Mogul Xtriage (Phenix) EDS buster-report	: : :	1.8.4, CSD as541be (2020) 1.13 2.24
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.24

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	278	2% 98%	
			2%	•
1	В	278	96%	•
1	С	278	94% 5%	<mark></mark>
1	D	278	95% ·	-
1	Е	278	^{3%} 96%	•



Continued from previous page...

Mol	Chain	Length	Quality of chain
1	Б	079	3%
L	Г	210	98% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CU	В	405	-	-	-	Х
5	PG4	А	406	-	-	Х	-



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 15938 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	278	Total	С	Ν	0	S	0	39	0
	A	210	2428	1507	450	454	17	0		0
1	В	277	Total	С	Ν	0	S	0	18	0
	D	211	2249	1407	406	421	15	0	10	0
1	С	277	Total	С	Ν	0	S	0	23	0
		211	2313	1439	421	435	18			
1	D	278	Total	С	Ν	0	S	0	21	0
	D	210	2301	1432	420	432	17	0	21	U
1	Е	278	Total	С	Ν	Ο	\mathbf{S}	0	47	0
1	Ľ	210	2492	1553	453	468	18	0	41	0
1	F	278	Total	С	Ν	Ο	S	0	36	0
	Ľ	210	2394	1488	434	456	16	0	- 30	U

• Molecule 1 is a protein called Two-domain laccase.

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	165	ALA	HIS	engineered mutation	UNP A0A0M4FJ81
А	240	HIS	ARG	engineered mutation	UNP A0A0M4FJ81
В	165	ALA	HIS	engineered mutation	UNP A0A0M4FJ81
В	240	HIS	ARG	engineered mutation	UNP A0A0M4FJ81
С	165	ALA	HIS	engineered mutation	UNP A0A0M4FJ81
С	240	HIS	ARG	engineered mutation	UNP A0A0M4FJ81
D	165	ALA	HIS	engineered mutation	UNP A0A0M4FJ81
D	240	HIS	ARG	engineered mutation	UNP A0A0M4FJ81
E	165	ALA	HIS	engineered mutation	UNP A0A0M4FJ81
Е	240	HIS	ARG	engineered mutation	UNP A0A0M4FJ81
F	165	ALA	HIS	engineered mutation	UNP A0A0M4FJ81
F	240	HIS	ARG	engineered mutation	UNP A0A0M4FJ81

• Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	3	Total Cu 3 3	0	0
2	В	5	Total Cu 5 5	0	0
2	С	4	Total Cu 4 4	0	0
2	D	3	Total Cu 3 3	0	0
2	Ε	4	Total Cu 4 4	0	0
2	F	5	Total Cu 5 5	0	0

• Molecule 3 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O₂) (labeled as "Ligand of Interest" by depositor).

OXY	
01 🔵 🔤 🔘 02	


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total O 2 2	0	0
3	В	1	Total O 2 2	0	0
3	С	1	Total O 2 2	0	0
3	D	1	Total O 2 2	0	0
3	Е	1	Total O 2 2	0	0



Continued from previous page...

Mo	Chain	hain Residues Atoms		ZeroOcc	AltConf	
3	F	1	Total O 2 2		0	0


• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0


• Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	А	1	Total 13	$\begin{array}{c} \mathrm{C} \\ \mathrm{8} \end{array}$	O 5	0	0

• Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 4 & 4 \end{array}$	0	1
6	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 4 & 4 \end{array}$	0	1

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	\mathbf{F}	1	Total 4	${ m C} 2$	O 2	0	0

• Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	С	1	Total C O 14 8 6	0	1


• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	280	Total O 280 280	0	0
8	В	287	Total O 287 287	0	0
8	С	259	Total O 259 259	0	0
8	D	277	Total O 277 277	0	0
8	Е	264	Total O 265 265	0	1
8	F	282	Total O 282 282	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Two-domain laccase

• Molecule 1: Two-domain laccase



Cha	ain	F:	3%	,)										98%	•
A40 G41	L59	<mark>de3</mark>	E91	N98	T208	M224	309	T311	312	313	314 317	315	317		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	74.76Å 94.29Å 120.48Å	Depositor
a, b, c, α , β , γ	90.00° 91.27° 90.00°	Depositor
Resolution (Å)	47.41 - 1.30	Depositor
Resolution (A)	47.41 - 1.30	EDS
% Data completeness	99.2 (47.41-1.30)	Depositor
(in resolution range)	$99.1 \ (47.41 - 1.30)$	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.29 (at 1.30 Å)	Xtriage
Refinement program	REFMAC 5.8.0230, PHENIX 1.19.2_4158	Depositor
D D.	0.133 , 0.157	Depositor
R, R_{free}	0.139 , 0.161	DCC
R_{free} test set	20412 reflections $(5.03%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	15.6	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning ²	$< L > = 0.50, < L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	15938	wwPDB-VP
Average B, all atoms $(Å^2)$	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OXY, EDO, PG4, GOL, PEG, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.34	0/2499	0.59	0/3388	
1	В	0.32	0/2321	0.59	0/3151	
1	С	0.31	0/2378	0.59	0/3222	
1	D	0.35	0/2364	0.60	0/3205	
1	Е	0.31	0/2560	0.59	0/3475	
1	F	0.34	0/2469	0.59	0/3347	
All	All	0.33	0/14591	0.59	0/19788	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2428	0	2296	4	0
1	В	2249	0	2129	7	0
1	С	2313	0	2177	16	0
1	D	2301	0	2163	21	0
1	Е	2492	0	2351	6	0
1	F	2394	0	2255	5	0
2	А	3	0	0	0	0



Mol	Chain	n previous	H(model)	H(added)	Clashes	Symm-Clashes
				. ,		
2	B	5	0	0	0	0
2	С	4	0	0	0	0
2	D	3	0	0	0	0
2	Е	4	0	0	0	0
2	F	5	0	0	0	0
3	А	2	0	0	0	0
3	В	2	0	0	0	0
3	С	2	0	0	0	0
3	D	2	0	0	0	0
3	Е	2	0	0	0	0
3	F	2	0	0	0	0
4	А	6	0	8	0	0
4	В	6	0	8	0	0
4	С	6	0	8	0	0
4	D	6	0	7	2	0
5	А	13	0	18	7	0
6	В	4	0	6	0	0
6	С	8	0	12	0	0
6	Е	8	0	12	1	0
6	F	4	0	6	0	0
7	С	14	0	20	3	0
8	А	280	0	0	1	0
8	В	287	0	0	1	0
8	С	259	0	0	1	0
8	D	277	0	0	1	0
8	Е	265	0	0	1	0
8	F	282	0	0	2	0
All	All	15938	0	13476	62	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:199[B]:MET:HE3	1:D:297[B]:MET:SD	1.28	1.66
1:D:199[B]:MET:CE	1:D:297[B]:MET:HG2	1.48	1.43
1:D:199[B]:MET:CE	1:D:297[B]:MET:CG	2.07	1.30
1:D:199[B]:MET:CE	1:D:297[B]:MET:SD	2.21	1.28
1:D:199[B]:MET:HE1	1:D:297[B]:MET:CG	1.73	1.14



There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	315/278~(113%)	308~(98%)	7(2%)	0	100	100
1	В	293/278~(105%)	291 (99%)	2 (1%)	0	100	100
1	С	299/278~(108%)	295~(99%)	4 (1%)	0	100	100
1	D	297/278~(107%)	294 (99%)	3 (1%)	0	100	100
1	Ε	323/278~(116%)	316~(98%)	7 (2%)	0	100	100
1	F	312/278~(112%)	310 (99%)	2 (1%)	0	100	100
All	All	1839/1668~(110%)	1814 (99%)	25 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	252/218~(116%)	252~(100%)	0	100 100
1	В	235/218~(108%)	232~(99%)	3 (1%)	69 35
1	С	241/218~(111%)	237~(98%)	4 (2%)	60 26
1	D	238/218~(109%)	236~(99%)	2(1%)	81 58
1	Е	259/218~(119%)	255~(98%)	4 (2%)	65 31
1	F	249/218~(114%)	246~(99%)	3 (1%)	71 40



Continued from previous page...

Mol	Chain	Analysed	Analysed Rotameric Ou		Percentiles
All	All	1474/1308~(113%)	1458~(99%)	16 (1%)	81 45

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	309[A]	ASP
1	F	224	MET
1	D	224	MET
1	Е	224[B]	MET
1	D	120	LYS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	А	257	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 43 ligands modelled in this entry, 24 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	gles
MOI	• -		nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	OXY	С	405	2	$1,\!1,\!1$	0.20	0	-		
6	EDO	F	407	-	3,3,3	0.54	0	$2,\!2,\!2$	0.13	0
3	OXY	В	406	2	$1,\!1,\!1$	0.20	0	-		
4	GOL	В	407	-	$5,\!5,\!5$	0.90	0	$5,\!5,\!5$	1.11	0
6	EDO	С	407[A]	-	3, 3, 3	0.47	0	$2,\!2,\!2$	0.22	0
4	GOL	А	405	-	$5,\!5,\!5$	0.89	0	$5,\!5,\!5$	1.04	0
6	EDO	Е	406[A]	-	3, 3, 3	0.40	0	$2,\!2,\!2$	0.54	0
3	OXY	А	404	2	$1,\!1,\!1$	0.16	0	-		
7	PEG	С	408[B]	-	$6,\!6,\!6$	0.44	0	$5,\!5,\!5$	0.52	0
6	EDO	В	408	-	3, 3, 3	0.37	0	$2,\!2,\!2$	0.53	0
3	OXY	F	406	2	$1,\!1,\!1$	0.20	0	-		
3	OXY	Е	405	2	$1,\!1,\!1$	0.26	0	-		
7	PEG	С	408[A]	-	$6,\!6,\!6$	0.43	0	$5,\!5,\!5$	0.35	0
4	GOL	С	406	-	$5,\!5,\!5$	0.85	0	$5,\!5,\!5$	1.03	0
4	GOL	D	405	-	$5,\!5,\!5$	1.10	1 (20%)	$5,\!5,\!5$	1.19	0
5	PG4	А	406	-	12,12,12	0.42	0	11,11,11	0.65	0
3	OXY	D	404	2	$1,\!1,\!1$	0.22	0	-		
6	EDO	Е	406[B]	-	3,3,3	0.55	0	$2,\!2,\!2$	0.16	0
6	EDO	С	407[B]	-	3, 3, 3	0.42	0	$2,\!2,\!2$	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PG4	А	406	-	-	5/10/10/10	-
7	PEG	С	408[B]	-	-	3/4/4/4	-
6	EDO	Е	406[B]	-	-	0/1/1/1	-
7	PEG	С	408[A]	-	-	1/4/4/4	-
4	GOL	С	406	-	-	1/4/4/4	-
4	GOL	D	405	-	-	0/4/4/4	-
4	GOL	В	407	-	-	2/4/4/4	-
6	EDO	С	407[B]	-	-	0/1/1/1	-
6	EDO	С	407[A]	-	-	1/1/1/1	-
6	EDO	F	407	-	-	1/1/1/1	-
6	EDO	В	408	-	-	0/1/1/1	-
4	GOL	А	405	-	-	4/4/4/4	-
6	EDO	Е	406[A]	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	405	GOL	O1-C1	-2.26	1.32	1.42

There are no bond angle outliers.

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	405	GOL	O1-C1-C2-O2
4	А	405	GOL	O1-C1-C2-C3
4	В	407	GOL	C1-C2-C3-O3
5	А	406	PG4	C6-C5-O3-C4
7	С	408[B]	PEG	O2-C3-C4-O4


There are no ring outliers.

5 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	С	408[B]	PEG	2	0
7	С	408[A]	PEG	1	0
4	D	405	GOL	2	0
5	А	406	PG4	7	0
6	Е	406[B]	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	А	278/278~(100%)	-0.27	6 (2%) 62 61	12, 19, 33, 51	1 (0%)
1	В	277/278~(99%)	-0.31	5 (1%) 68 69	12, 17, 31, 56	0
1	С	277/278~(99%)	-0.20	7 (2%) 57 56	12, 21, 36, 64	0
1	D	278/278~(100%)	-0.31	4 (1%) 75 77	11, 19, 33, 55	0
1	Ε	278/278~(100%)	-0.19	9 (3%) 47 45	12, 20, 35, 64	0
1	F	278/278~(100%)	-0.17	9 (3%) 47 45	11, 18, 36, 58	1 (0%)
All	All	1666/1668~(99%)	-0.24	40 (2%) 59 58	11, 19, 34, 64	2 (0%)

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	40	ALA	9.7
1	D	40	ALA	9.6
1	F	40	ALA	9.4
1	Е	316	ASP	6.7
1	А	142[A]	ALA	5.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
2	CU	В	405	1/1	0.35	1.12	$55,\!55,\!55,\!55$	1
7	PEG	С	408[A]	7/7	0.55	0.23	40,42,45,45	7
7	PEG	С	408[B]	7/7	0.55	0.23	39,42,45,46	7
4	GOL	С	406	6/6	0.71	0.18	47,49,50,50	6
4	GOL	В	407	6/6	0.77	0.21	42,45,45,46	6
6	EDO	Е	406[A]	4/4	0.79	0.19	37,37,37,37	4
6	EDO	Е	406[B]	4/4	0.79	0.19	26,28,28,31	4
5	PG4	А	406	13/13	0.81	0.22	38,42,44,45	13
4	GOL	А	405	6/6	0.83	0.18	44,49,50,50	0
4	GOL	D	405	6/6	0.86	0.15	34,35,36,37	6
6	EDO	F	407	4/4	0.86	0.10	26,26,27,27	0
6	EDO	В	408	4/4	0.87	0.12	27,29,29,32	0
6	EDO	С	407[B]	4/4	0.90	0.10	32,33,33,33	4
6	EDO	С	407[A]	4/4	0.90	0.10	23,26,27,28	4
3	OXY	Е	405	2/2	0.94	0.12	16,16,16,17	2
2	CU	F	405	1/1	0.95	0.07	18,18,18,18	1
3	OXY	С	405	2/2	0.95	0.11	12,12,12,18	2
3	OXY	D	404	2/2	0.97	0.09	19,19,19,21	2
2	CU	Е	403	1/1	0.98	0.04	19,19,19,19	1
3	OXY	В	406	2/2	0.98	0.08	19,19,19,24	2
3	OXY	F	406	2/2	0.98	0.09	19,19,19,20	2
2	CU	F	402	1/1	0.98	0.05	19,19,19,19	1
2	CU	С	402	1/1	0.99	0.05	17,17,17,17	1
3	OXY	А	404	2/2	0.99	0.07	23,23,23,25	2
2	CU	В	403	1/1	1.00	0.04	17,17,17,17	1
2	CU	В	404	1/1	1.00	0.03	21,21,21,21	1
2	CU	А	401	1/1	1.00	0.06	14,14,14,14	0
2	CU	С	401	1/1	1.00	0.04	21,21,21,21	1
2	CU	А	402	1/1	1.00	0.03	24,24,24,24	1
2	CU	С	403	1/1	1.00	0.06	14,14,14,14	0
2	CU	С	404	1/1	1.00	0.03	18,18,18,18	0
2	CU	D	401	1/1	1.00	0.05	13,13,13,13	0
2	CU	D	402	1/1	1.00	0.04	16,16,16,16	0
2	CU	D	403	1/1	1.00	0.04	17,17,17,17	1
2	CU	Е	401	1/1	1.00	0.06	13,13,13,13	0
2	CU	Е	402	1/1	1.00	0.05	19,19,19,19	1
2	CU	А	403	1/1	1.00	0.06	17,17,17,17	1




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	$Q{<}0.9$
2	CU	Ε	404	1/1	1.00	0.08	16, 16, 16, 16	1
2	CU	F	401	1/1	1.00	0.03	18,18,18,18	1
2	CU	В	401	1/1	1.00	0.05	13,13,13,13	0
2	CU	F	403	1/1	1.00	0.05	14,14,14,14	0
2	CU	F	404	1/1	1.00	0.04	19,19,19,19	1
2	CU	В	402	1/1	1.00	0.04	19,19,19,19	1


Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















































































































































6.5 Other polymers (i)

There are no such residues in this entry.

