



wwPDB EM Validation Summary Report ⓘ

Mar 10, 2024 – 06:38 am GMT

PDB ID : 8Q72
EMDB ID : EMD-18201
Title : E. coli plasmid-borne JetABCD(E248A) core in a cleavage-competent state
Authors : Roisne-Hamelin, F.; Li, Y.; Gruber, S.
Deposited on : 2023-08-15
Resolution : 4.17 Å (reported)
Based on initial models : 8AS8, .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1096	
1	B	1096	
1	F	1096	
1	G	1096	
2	C	250	
2	D	250	
2	H	250	
2	I	250	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	503	 75% 17% 7%
3	J	503	 78% 15% 7%
4	M	390	 27% 70%
4	N	390	 26% 70%
5	P	40	 70% 30%
5	Q	40	 62% 38%
5	R	40	 72% 28%
5	S	40	 62% 38%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 84390 atoms, of which 41400 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called JetC.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	A	750	11881	3710	5918	1093	1144	16	0	0
1	B	750	11880	3710	5917	1093	1144	16	0	0
1	F	750	11881	3710	5918	1093	1144	16	0	0
1	G	750	11880	3710	5917	1093	1144	16	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	283	LEU	GLN	conflict	UNP A0A6D0I2P0
A	298	SER	ASN	conflict	UNP A0A6D0I2P0
A	386	SER	ILE	conflict	UNP A0A6D0I2P0
A	398	GLU	ALA	conflict	UNP A0A6D0I2P0
A	400	ARG	LEU	conflict	UNP A0A6D0I2P0
A	576	HIS	ARG	conflict	UNP A0A6D0I2P0
A	625	ALA	THR	conflict	UNP A0A6D0I2P0
A	647	LEU	ILE	conflict	UNP A0A6D0I2P0
A	705	ILE	VAL	conflict	UNP A0A6D0I2P0
A	729	LEU	SER	conflict	UNP A0A6D0I2P0
A	817	ALA	THR	conflict	UNP A0A6D0I2P0
A	823	PRO	THR	conflict	UNP A0A6D0I2P0
A	889	ASP	TYR	conflict	UNP A0A6D0I2P0
A	933	VAL	ILE	conflict	UNP A0A6D0I2P0
A	1096	GLY	-	expression tag	UNP A0A6D0I2P0
B	283	LEU	GLN	conflict	UNP A0A6D0I2P0
B	298	SER	ASN	conflict	UNP A0A6D0I2P0
B	386	SER	ILE	conflict	UNP A0A6D0I2P0
B	398	GLU	ALA	conflict	UNP A0A6D0I2P0
B	400	ARG	LEU	conflict	UNP A0A6D0I2P0
B	576	HIS	ARG	conflict	UNP A0A6D0I2P0
B	625	ALA	THR	conflict	UNP A0A6D0I2P0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	647	LEU	ILE	conflict	UNP A0A6D0I2P0
B	705	ILE	VAL	conflict	UNP A0A6D0I2P0
B	729	LEU	SER	conflict	UNP A0A6D0I2P0
B	817	ALA	THR	conflict	UNP A0A6D0I2P0
B	823	PRO	THR	conflict	UNP A0A6D0I2P0
B	889	ASP	TYR	conflict	UNP A0A6D0I2P0
B	933	VAL	ILE	conflict	UNP A0A6D0I2P0
B	1096	GLY	-	expression tag	UNP A0A6D0I2P0
F	283	LEU	GLN	conflict	UNP A0A6D0I2P0
F	298	SER	ASN	conflict	UNP A0A6D0I2P0
F	386	SER	ILE	conflict	UNP A0A6D0I2P0
F	398	GLU	ALA	conflict	UNP A0A6D0I2P0
F	400	ARG	LEU	conflict	UNP A0A6D0I2P0
F	576	HIS	ARG	conflict	UNP A0A6D0I2P0
F	625	ALA	THR	conflict	UNP A0A6D0I2P0
F	647	LEU	ILE	conflict	UNP A0A6D0I2P0
F	705	ILE	VAL	conflict	UNP A0A6D0I2P0
F	729	LEU	SER	conflict	UNP A0A6D0I2P0
F	817	ALA	THR	conflict	UNP A0A6D0I2P0
F	823	PRO	THR	conflict	UNP A0A6D0I2P0
F	889	ASP	TYR	conflict	UNP A0A6D0I2P0
F	933	VAL	ILE	conflict	UNP A0A6D0I2P0
F	1096	GLY	-	expression tag	UNP A0A6D0I2P0
G	283	LEU	GLN	conflict	UNP A0A6D0I2P0
G	298	SER	ASN	conflict	UNP A0A6D0I2P0
G	386	SER	ILE	conflict	UNP A0A6D0I2P0
G	398	GLU	ALA	conflict	UNP A0A6D0I2P0
G	400	ARG	LEU	conflict	UNP A0A6D0I2P0
G	576	HIS	ARG	conflict	UNP A0A6D0I2P0
G	625	ALA	THR	conflict	UNP A0A6D0I2P0
G	647	LEU	ILE	conflict	UNP A0A6D0I2P0
G	705	ILE	VAL	conflict	UNP A0A6D0I2P0
G	729	LEU	SER	conflict	UNP A0A6D0I2P0
G	817	ALA	THR	conflict	UNP A0A6D0I2P0
G	823	PRO	THR	conflict	UNP A0A6D0I2P0
G	889	ASP	TYR	conflict	UNP A0A6D0I2P0
G	933	VAL	ILE	conflict	UNP A0A6D0I2P0
G	1096	GLY	-	expression tag	UNP A0A6D0I2P0

- Molecule 2 is a protein called JetB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	H	N	O		
2	C	195	3173	991	1607	280	295	0	0
2	D	195	3172	991	1606	280	295	0	0
2	H	195	3173	991	1607	280	295	0	0
2	I	195	3172	991	1606	280	295	0	0

- Molecule 3 is a protein called JetA.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
3	E	466	7532	2356	3745	700	723	8	0	0
3	J	466	7532	2356	3745	700	723	8	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-3	GLY	-	expression tag	UNP A0A4V3QHV5
E	-2	PRO	-	expression tag	UNP A0A4V3QHV5
E	-1	ALA	-	expression tag	UNP A0A4V3QHV5
E	0	ALA	-	expression tag	UNP A0A4V3QHV5
E	187	ASP	GLU	conflict	UNP A0A4V3QHV5
E	435	GLU	ALA	engineered mutation	UNP A0A4V3QHV5
E	499	GLY	-	expression tag	UNP A0A4V3QHV5
J	-3	GLY	-	expression tag	UNP A0A4V3QHV5
J	-2	PRO	-	expression tag	UNP A0A4V3QHV5
J	-1	ALA	-	expression tag	UNP A0A4V3QHV5
J	0	ALA	-	expression tag	UNP A0A4V3QHV5
J	187	ASP	GLU	conflict	UNP A0A4V3QHV5
J	435	GLU	ALA	engineered mutation	UNP A0A4V3QHV5
J	499	GLY	-	expression tag	UNP A0A4V3QHV5

- Molecule 4 is a protein called JetD(E248A).

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
4	M	117	1917	611	961	174	168	3	0	0
4	N	117	1917	611	961	174	168	3	0	0


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	74	PHE	TYR	conflict	UNP A0A3T6B0Z0
M	248	ALA	GLU	engineered mutation	UNP A0A3T6B0Z0
M	358	ARG	CYS	conflict	UNP A0A3T6B0Z0
M	374	ALA	VAL	conflict	UNP A0A3T6B0Z0
M	383	GLY	-	expression tag	UNP A0A3T6B0Z0
M	384	SER	-	expression tag	UNP A0A3T6B0Z0
M	385	LEU	-	expression tag	UNP A0A3T6B0Z0
M	386	GLU	-	expression tag	UNP A0A3T6B0Z0
M	387	VAL	-	expression tag	UNP A0A3T6B0Z0
M	388	LEU	-	expression tag	UNP A0A3T6B0Z0
M	389	PHE	-	expression tag	UNP A0A3T6B0Z0
M	390	GLN	-	expression tag	UNP A0A3T6B0Z0
N	74	PHE	TYR	conflict	UNP A0A3T6B0Z0
N	248	ALA	GLU	engineered mutation	UNP A0A3T6B0Z0
N	358	ARG	CYS	conflict	UNP A0A3T6B0Z0
N	374	ALA	VAL	conflict	UNP A0A3T6B0Z0
N	383	GLY	-	expression tag	UNP A0A3T6B0Z0
N	384	SER	-	expression tag	UNP A0A3T6B0Z0
N	385	LEU	-	expression tag	UNP A0A3T6B0Z0
N	386	GLU	-	expression tag	UNP A0A3T6B0Z0
N	387	VAL	-	expression tag	UNP A0A3T6B0Z0
N	388	LEU	-	expression tag	UNP A0A3T6B0Z0
N	389	PHE	-	expression tag	UNP A0A3T6B0Z0
N	390	GLN	-	expression tag	UNP A0A3T6B0Z0


- Molecule 5 is a DNA chain called Circular plasmid DNA (1840-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
5	P	40	Total	C	H	N	O	P	0	0
			1281	400	461	140	240	40		
5	Q	40	Total	C	H	N	O	P	0	0
			1281	400	461	140	240	40		
5	R	40	Total	C	H	N	O	P	0	0
			1281	400	461	140	240	40		
5	S	40	Total	C	H	N	O	P	0	0
			1281	400	461	140	240	40		


- Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					AltConf	
			Total	C	H	N	O		P
6	A	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
6	B	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
6	F	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
6	G	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	




• Molecule 1: JetC







• Molecule 2: JetB




• Molecule 2: JetB



• Molecule 2: JetB





• Molecule 2: JetB








• Molecule 3: JetA



• Molecule 3: JetA



• Molecule 4: JetD(E248A)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	235956	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.153	Depositor
Minimum map value	-0.661	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.019	Depositor
Recommended contour level	0.102	Depositor
Map size (Å)	531.2, 531.2, 531.2	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.6600001, 1.6600001, 1.6600001	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/6059	0.52	0/8179
1	B	0.24	0/6059	0.50	0/8179
1	F	0.25	0/6059	0.52	0/8179
1	G	0.24	0/6059	0.50	0/8179
2	C	0.24	0/1592	0.53	0/2164
2	D	0.23	0/1592	0.50	0/2164
2	H	0.23	0/1592	0.51	0/2164
2	I	0.23	0/1592	0.50	0/2164
3	E	0.24	0/3849	0.53	0/5205
3	J	0.24	0/3849	0.53	0/5205
4	M	0.24	0/982	0.51	0/1337
4	N	0.23	0/982	0.48	0/1337
5	P	0.56	0/919	1.01	0/1416
5	Q	0.56	0/919	0.99	0/1416
5	R	0.55	0/919	0.99	0/1416
5	S	0.55	0/919	1.00	0/1416
All	All	0.28	0/43942	0.58	0/60120

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	126	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5963	5918	5916	75	0
1	B	5963	5917	5916	66	0
1	F	5963	5918	5916	70	0
1	G	5963	5917	5916	72	0
2	C	1566	1607	1605	34	0
2	D	1566	1606	1605	24	0
2	H	1566	1607	1605	22	0
2	I	1566	1606	1605	23	0
3	E	3787	3745	3745	68	0
3	J	3787	3745	3745	57	0
4	M	956	961	961	7	0
4	N	956	961	961	9	0
5	P	820	461	461	11	0
5	Q	820	461	461	11	0
5	R	820	461	461	7	0
5	S	820	461	461	12	0
6	A	27	12	12	3	0
6	B	27	12	12	3	0
6	F	27	12	12	2	0
6	G	27	12	12	4	0
All	All	42990	41400	41388	503	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 503 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:128:GLU:HG2	3:E:350:ARG:HE	1.25	0.98
3:E:68:GLU:OE1	1:F:912:ARG:NH1	2.05	0.90
1:F:172:GLU:OE1	1:F:180:THR:OG1	1.91	0.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:GLU:OE1	1:A:180:THR:OG1	1.92	0.88
2:I:137:ARG:NH2	2:I:210:ILE:O	2.09	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	746/1096 (68%)	720 (96%)	26 (4%)	0	100	100
1	B	746/1096 (68%)	721 (97%)	25 (3%)	0	100	100
1	F	746/1096 (68%)	713 (96%)	32 (4%)	1 (0%)	51	85
1	G	746/1096 (68%)	724 (97%)	22 (3%)	0	100	100
2	C	193/250 (77%)	189 (98%)	4 (2%)	0	100	100
2	D	193/250 (77%)	187 (97%)	6 (3%)	0	100	100
2	H	193/250 (77%)	186 (96%)	7 (4%)	0	100	100
2	I	193/250 (77%)	185 (96%)	8 (4%)	0	100	100
3	E	462/503 (92%)	441 (96%)	20 (4%)	1 (0%)	47	80
3	J	462/503 (92%)	442 (96%)	20 (4%)	0	100	100
4	M	115/390 (30%)	114 (99%)	1 (1%)	0	100	100
4	N	115/390 (30%)	113 (98%)	2 (2%)	0	100	100
All	All	4910/7170 (68%)	4735 (96%)	173 (4%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	258	SER
1	F	99	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	635/929 (68%)	635 (100%)	0	100	100
1	B	635/929 (68%)	633 (100%)	2 (0%)	92	95
1	F	635/929 (68%)	633 (100%)	2 (0%)	92	95
1	G	635/929 (68%)	633 (100%)	2 (0%)	92	95
2	C	171/218 (78%)	171 (100%)	0	100	100
2	D	171/218 (78%)	171 (100%)	0	100	100
2	H	171/218 (78%)	171 (100%)	0	100	100
2	I	171/218 (78%)	171 (100%)	0	100	100
3	E	408/436 (94%)	408 (100%)	0	100	100
3	J	408/436 (94%)	408 (100%)	0	100	100
4	M	103/336 (31%)	103 (100%)	0	100	100
4	N	103/336 (31%)	103 (100%)	0	100	100
All	All	4246/6132 (69%)	4240 (100%)	6 (0%)	93	97

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	1092	ARG
1	G	1011	ARG
1	G	1026	ARG
1	B	1026	ARG
1	B	1011	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	250	GLN
1	G	259	GLN
4	N	55	HIS
2	H	164	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	J	19	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ADP	B	1101	-	24,29,29	1.76	4 (16%)	29,45,45	1.32	3 (10%)
6	ADP	F	1101	-	24,29,29	1.76	3 (12%)	29,45,45	1.37	2 (6%)
6	ADP	A	1101	-	24,29,29	1.79	4 (16%)	29,45,45	1.36	2 (6%)
6	ADP	G	1101	-	24,29,29	1.76	4 (16%)	29,45,45	1.30	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADP	B	1101	-	-	3/12/32/32	0/3/3/3
6	ADP	F	1101	-	-	7/12/32/32	0/3/3/3
6	ADP	A	1101	-	-	6/12/32/32	0/3/3/3
6	ADP	G	1101	-	-	4/12/32/32	0/3/3/3

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	F	1101	ADP	PA-O5'	5.40	1.81	1.59
6	G	1101	ADP	PA-O5'	5.32	1.80	1.59
6	A	1101	ADP	PA-O5'	5.17	1.80	1.59
6	B	1101	ADP	PA-O5'	5.14	1.80	1.59
6	A	1101	ADP	C2'-C1'	4.00	1.59	1.53

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1101	ADP	O5'-PA-O1A	-3.62	94.93	109.07
6	F	1101	ADP	O5'-PA-O1A	-3.59	95.05	109.07
6	B	1101	ADP	O3B-PB-O2B	2.74	118.12	107.64
6	B	1101	ADP	O2A-PA-O5'	-2.73	95.07	107.75
6	G	1101	ADP	O2A-PA-O5'	-2.67	95.34	107.75

There are no chirality outliers.

5 of 20 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	1101	ADP	PA-O3A-PB-O2B
6	A	1101	ADP	C5'-O5'-PA-O3A
6	F	1101	ADP	PA-O3A-PB-O2B
6	F	1101	ADP	C5'-O5'-PA-O2A
6	F	1101	ADP	C5'-O5'-PA-O3A



There are no ring outliers.


4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1101	ADP	3	0
6	F	1101	ADP	2	0
6	A	1101	ADP	3	0
6	G	1101	ADP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-18201. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 155




Y Index: 145




Z Index: 181


6.3.2 Raw map



X Index: 155



Y Index: 175




Z Index: 181


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z

6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.102. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_18201_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 701 nm^3 ; this corresponds to an approximate mass of 633 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.240 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.240 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.17	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.72	7.66	4.81

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.72 differs from the reported value 4.17 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-18201 and PDB model 8Q72. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.102 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.102).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 91% of all backbone atoms, 91% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.102) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9120	 0.3070
A	 0.9010	 0.3600
B	 0.8640	 0.2950
C	 0.9330	 0.2610
D	 0.9290	 0.2080
E	 0.9700	 0.3750
F	 0.9000	 0.3600
G	 0.8630	 0.2940
H	 0.9270	 0.2590
I	 0.9330	 0.2080
J	 0.9690	 0.3750
M	 0.9240	 0.2230
N	 0.9300	 0.2270
P	 0.9500	 0.1920
Q	 0.9760	 0.1950
R	 0.9550	 0.1900
S	 0.9760	 0.1950

