

# wwPDB X-ray Structure Validation Summary Report (i)

May 24, 2020 – 08:39 am BST

PDB ID	:	4QB5
Title	:	$Crystal\ structure\ of\ a\ glyoxalase/bleomycin\ resistance\ protein\ from\ Albidiferax$
		ferrireducens T118
Authors	:	Kumaran, D.; Chamala, S.; Evans, B.; Foti, R.; Gizzi, A.; Hillerich, B.; Kar,
		A.; Lafleur, J.; Seidel, R.; Villigas, G.; Zencheck, W.; Al Obaidi, N.; Almo,
		S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium
		(NYSGRC)
Deposited on		
Resolution	:	2.05  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
$\operatorname{Refmac}$	:	5.8.0158
$\operatorname{CCP4}$	:	7.0.044   (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11
· ( )		

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
$R_{free}$	130704	1692(2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752(2.04-2.04)
Sidechain outliers	138945	1752(2.04-2.04)
RSRZ outliers	127900	1672(2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	148	<sup>2%</sup> 72%	7%	•	-	19%	
1	В	148	6% 70%	11%		•	16%	
1	С	148	73%	7%	•		18%	
1	D	148	<sup>2%</sup> 72%	119	6	·	16%	



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	Δ	120	Total	С	Ν	Ο	$\mathbf{S}$	$\operatorname{Se}$	0	0	0
	A	120	893	563	156	169	3	2	0	0	0
1	В	124	Total	С	Ν	Ο	$\mathbf{S}$	Se	0	0	0
	D	124	915	576	161	173	3	2	0	0	0
1	С	121	Total	С	Ν	Ο	$\mathbf{S}$	$\operatorname{Se}$	0	0	0
		121	902	569	158	170	3	2			0
1	п	195	Total	С	Ν	Ο	S	Se	0	0	0
		125	922	581	162	174	3	2	0	0	0

• Molecule 1 is a protein called Glyoxalase/bleomycin resistance protein/dioxygenase.

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MSE	-	EXPRESSION TAG	UNP Q21R90
A	-20	HIS	-	EXPRESSION TAG	UNP Q21R90
A	-19	HIS	-	EXPRESSION TAG	UNP Q21R90
A	-18	HIS	-	EXPRESSION TAG	UNP Q21R90
A	-17	HIS	-	EXPRESSION TAG	UNP Q21R90
A	-16	HIS	-	EXPRESSION TAG	UNP Q21R90
A	-15	HIS	-	EXPRESSION TAG	UNP Q21R90
A	-14	SER	-	EXPRESSION TAG	UNP Q21R90
A	-13	SER	-	EXPRESSION TAG	UNP Q21R90
А	-12	GLY	-	EXPRESSION TAG	UNP Q21R90
A	-11	VAL	-	EXPRESSION TAG	UNP Q21R90
A	-10	ASP	-	EXPRESSION TAG	UNP Q21R90
A	-9	LEU	-	EXPRESSION TAG	UNP Q21R90
A	-8	GLY	-	EXPRESSION TAG	UNP Q21R90
A	-7	THR	-	EXPRESSION TAG	UNP Q21R90
A	-6	GLU	-	EXPRESSION TAG	UNP Q21R90
А	-5	ASN	-	EXPRESSION TAG	UNP Q21R90
А	-4	LEU	-	EXPRESSION TAG	UNP Q21R90
A	-3	TYR	-	EXPRESSION TAG	UNP Q21R90
А	-2	PHE	-	EXPRESSION TAG	UNP Q21R90
А	-1	GLN	-	EXPRESSION TAG	UNP Q21R90

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP Q21R90
В	-21	MSE	-	EXPRESSION TAG	UNP Q21R90
В	-20	HIS	_	EXPRESSION TAG	UNP Q21R90
В	-19	HIS	-	EXPRESSION TAG	UNP Q21R90
В	-18	HIS	_	EXPRESSION TAG	UNP Q21R90
В	-17	HIS	-	EXPRESSION TAG	UNP Q21R90
В	-16	HIS	-	EXPRESSION TAG	UNP Q21R90
В	-15	HIS	-	EXPRESSION TAG	UNP Q21R90
В	-14	SER	-	EXPRESSION TAG	UNP Q21R90
В	-13	SER	-	EXPRESSION TAG	UNP Q21R90
В	-12	GLY	-	EXPRESSION TAG	UNP Q21R90
В	-11	VAL	-	EXPRESSION TAG	UNP Q21R90
В	-10	ASP	-	EXPRESSION TAG	UNP Q21R90
В	-9	LEU	-	EXPRESSION TAG	UNP Q21R90
В	-8	GLY	-	EXPRESSION TAG	UNP Q21R90
В	-7	THR	-	EXPRESSION TAG	UNP Q21R90
В	-6	GLU	-	EXPRESSION TAG	UNP Q21R90
В	-5	ASN	_	EXPRESSION TAG	UNP Q21R90
В	-4	LEU	-	EXPRESSION TAG	UNP Q21R90
В	-3	TYR	-	EXPRESSION TAG	UNP Q21R90
В	-2	PHE	-	EXPRESSION TAG	UNP Q21R90
В	-1	GLN	-	EXPRESSION TAG	UNP Q21R90
В	0	SER	-	EXPRESSION TAG	UNP Q21R90
С	-21	MSE	-	EXPRESSION TAG	UNP Q21R90
С	-20	HIS	-	EXPRESSION TAG	UNP Q21R90
С	-19	HIS	-	EXPRESSION TAG	UNP Q21R90
С	-18	HIS	-	EXPRESSION TAG	UNP Q21R90
С	-17	HIS	-	EXPRESSION TAG	UNP Q21R90
С	-16	HIS	-	EXPRESSION TAG	UNP Q21R90
С	-15	HIS	-	EXPRESSION TAG	
С	-14	SER	-	EXPRESSION TAG	UNP Q21R90
С	-13	SER	-	EXPRESSION TAG	UNP Q21R90
С	-12	GLY	-	EXPRESSION TAG	UNP Q21R90
С	-11	VAL	-	EXPRESSION TAG	UNP Q21R90
С	-10	ASP	-	EXPRESSION TAG	UNP Q21R90
С	-9	LEU	-	EXPRESSION TAG	UNP Q21R90
С	-8	GLY	-	EXPRESSION TAG	UNP Q21R90
С	-7	THR	-	EXPRESSION TAG	UNP Q21R90
С	-6	GLU	-	EXPRESSION TAG	UNP Q21R90
С	-5	ASN	-	EXPRESSION TAG	UNP Q21R90
С	-4	LEU	-	EXPRESSION TAG	UNP Q21R90
С	-3	TYR	-	EXPRESSION TAG	UNP Q21R90

Continued from previous page...
Chain Residue Modelled Actual


Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	PHE	-	EXPRESSION TAG	UNP Q21R90
С	-1	GLN	-	EXPRESSION TAG	UNP Q21R90
С	0	SER	-	EXPRESSION TAG	UNP Q21R90
D	-21	MSE	-	EXPRESSION TAG	UNP Q21R90
D	-20	HIS	-	EXPRESSION TAG	UNP Q21R90
D	-19	HIS	-	EXPRESSION TAG	UNP Q21R90
D	-18	HIS	-	EXPRESSION TAG	UNP Q21R90
D	-17	HIS	-	EXPRESSION TAG	UNP Q21R90
D	-16	HIS	-	EXPRESSION TAG	UNP Q21R90
D	-15	HIS	-	EXPRESSION TAG	UNP Q21R90
D	-14	SER	-	EXPRESSION TAG	UNP Q21R90
D	-13	SER	-	EXPRESSION TAG	UNP Q21R90
D	-12	GLY	-	EXPRESSION TAG	UNP Q21R90
D	-11	VAL	-	EXPRESSION TAG	UNP Q21R90
D	-10	ASP	-	EXPRESSION TAG	UNP Q21R90
D	-9	LEU	-	EXPRESSION TAG	UNP Q21R90
D	-8	GLY	-	EXPRESSION TAG	UNP Q21R90
D	-7	THR	-	EXPRESSION TAG	UNP Q21R90
D	-6	GLU	-	EXPRESSION TAG	UNP Q21R90
D	-5	ASN	-	EXPRESSION TAG	UNP Q21R90
D	-4	LEU	-	EXPRESSION TAG	UNP Q21R90
D	-3	TYR	-	EXPRESSION TAG	UNP Q21R90
D	-2	PHE	-	EXPRESSION TAG	UNP Q21R90
D	-1	GLN	-	EXPRESSION TAG	UNP Q21R90
D	0	SER	-	EXPRESSION TAG	UNP Q21R90

Continued from previous page...

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).








Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	72	Total O 72 72	0	0
4	В	45	Total         O           45         45	0	0
4	С	49	Total         O           49         49	0	0
4	D	80	Total         O           80         80	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glyoxalase/bleomycin resistance protein/dioxygenase





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	46.95Å $89.90$ Å $131.86$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	39.72 - 2.05	Depositor
Resolution (A)	39.69 - 2.05	EDS
% Data completeness	96.8 (39.72-2.05)	Depositor
(in resolution range)	96.9(39.69-2.05)	EDS
R <sub>merge</sub>	0.07	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$15.57 (at 2.05 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
D D.	0.172 , $0.211$	Depositor
$R, R_{free}$	0.187 , $0.224$	DCC
$R_{free}$ test set	1734 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	21.5	Xtriage
Anisotropy	0.538	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37, 50.3	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3897	wwPDB-VP
Average B, all atoms $(Å^2)$	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bo	ond angles
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.96	1/914~(0.1%)	0.94	4/1236~(0.3%)
1	В	0.89	2/936~(0.2%)	0.96	4/1265~(0.3%)
1	С	0.93	1/923~(0.1%)	0.92	2/1247~(0.2%)
1	D	1.01	2/944~(0.2%)	1.00	5/1277~(0.4%)
All	All	0.95	6/3717~(0.2%)	0.96	15/5025~(0.3%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	D	119	CYS	CB-SG	-11.50	1.62	1.82
1	D	29	GLY	N-CA	7.77	1.57	1.46
1	А	119	CYS	CB-SG	-7.75	1.69	1.82
1	С	29	GLY	N-CA	6.69	1.56	1.46
1	В	29	GLY	N-CA	6.60	1.55	1.46

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	6	MSE	CG-SE-CE	6.79	113.84	98.90
1	D	26	LEU	CB-CG-CD1	6.60	122.22	111.00
1	В	110	LEU	CA-CB-CG	6.10	129.34	115.30
1	С	111	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	D	29	GLY	N-CA-C	-5.97	98.18	113.10

There are no chirality outliers.

There are no planarity outliers.



#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	893	0	856	7	0
1	В	915	0	883	18	0
1	С	902	0	869	9	0
1	D	922	0	890	14	0
2	В	5	0	0	1	0
2	D	10	0	0	0	0
3	D	4	0	6	0	0
4	А	72	0	0	0	0
4	В	45	0	0	1	0
4	С	49	0	0	0	0
4	D	80	0	0	3	0
All	All	3897	0	3504	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:93:GLU:HG3	1:C:94:PRO:HD2	1.58	0.85
1:B:10:ASN:HD21	1:B:58:ALA:H	1.35	0.73
1:C:10:ASN:HD21	1:C:58:ALA:H	1.35	0.73
1:A:5:VAL:O	1:A:6:MSE:HE2	1.93	0.68
1:D:10:ASN:HD21	1:D:58:ALA:H	1.43	0.67

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	116/148~(78%)	113~(97%)	3~(3%)	0	100	100
1	В	122/148~(82%)	115~(94%)	5(4%)	2(2%)	9	2
1	С	117/148~(79%)	112 (96%)	5 (4%)	0	100	100
1	D	123/148~(83%)	117 (95%)	6~(5%)	0	100	100
All	All	478/592~(81%)	457 (96%)	19 (4%)	2(0%)	34	24

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	101	ALA
1	В	102	VAL

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentil	es
1	А	92/111~(83%)	89~(97%)	3~(3%)	38 31	
1	В	93/111~(84%)	91~(98%)	2(2%)	52 46	
1	С	93/111~(84%)	90~(97%)	3(3%)	39 32	
1	D	94/111~(85%)	91~(97%)	3(3%)	39 32	
All	All	372/444~(84%)	361~(97%)	11 (3%)	41 34	

5 of 11 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	В	110	LEU
1	С	37	ARG
1	D	26	LEU
1	В	37	ARG
1	С	93	GLU



Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	86	ASN
1	В	116	ASN
1	С	116	ASN
1	В	80	HIS
1	С	86	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type	Chain	Chain Res	Link	Bond lengths				Bond angles		
	туре	Cham	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2	
3	EDO	D	203	-	3,3,3	0.68	0	$2,\!2,\!2$	0.26	0	
2	SO4	D	201	-	4,4,4	0.50	0	$^{6,6,6}$	0.53	0	
2	SO4	D	202	-	4,4,4	0.46	0	$^{6,6,6}$	0.92	0	
2	SO4	В	201	-	4,4,4	0.35	0	$^{6,6,6}$	0.82	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	203	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	201	SO4	1	0

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$\mathbf{OWAB}(\mathrm{\AA}^2)$	$Q{<}0.9$	
1	А	118/148~(79%)	0.07	3(2%)	57	61	12, 19, 42, 64	0
1	В	122/148~(82%)	0.16	9 (7%)	14	16	16, 24, 56, 108	0
1	С	119/148~(80%)	-0.09	7(5%)	22	24	13, 19, 42, 69	0
1	D	123/148~(83%)	0.16	3(2%)	59	63	11, 19, 50, 88	0
All	All	482/592 (81%)	0.08	22 (4%)	32	35	11, 20, 49, 108	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	101	ALA	8.6
1	В	100	GLY	8.2
1	В	99	ASP	5.7
1	D	100	GLY	5.3
1	D	102	VAL	5.2

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	$\mathbf{RSR}$	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	$Q{<}0.9$
2	SO4	В	201	5/5	0.94	0.17	$55,\!58,\!62,\!68$	0
3	EDO	D	203	4/4	0.96	0.12	22,22,23,26	0
2	SO4	D	202	5/5	0.98	0.08	$30,\!31,\!34,\!35$	0
2	SO4	D	201	5/5	0.98	0.09	$30,\!35,\!35,\!42$	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

