

wwPDB X-ray Structure Validation Summary Report (i)

Jan 27, 2022 – 03:36 pm GMT

PDB ID : 7QBT

Title: B12-dependent radical SAM methyltransferase, Mmp10 with [4Fe-4S] cluster,

cobalamin, and S-methyl-5'-thioadenosine bound.

Authors: Fyfe, C.D.; Chavas, L.M.G.; Legrand, P.; Benjdia, A.; Berteau, O.

Deposited on : 2021-11-19

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.26

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0267

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned} ext{Similar resolution} \ (\# ext{Entries, resolution range}(ext{Å})) \end{aligned}$		
R_{free}	130704	6207 (1.90-1.90)		
Clashscore	141614	6847 (1.90-1.90)		
Ramachandran outliers	138981	6760 (1.90-1.90)		
Sidechain outliers	138945	6760 (1.90-1.90)		
RSRZ outliers	127900	6082 (1.90-1.90)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	436	92%	• 6%
1	В	436	91%	• 6%
1	С	436	16%	• 7%
1	D	436	91%	• 6%



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 14382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Methyl coenzyme M reductase-arginine methyltransferase Mmp10.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	٨	411	Total	С	N	О	S	0	1	0
1	A	411	3178	2027	521	609	21	U	1	
1	В	408	Total	С	N	О	S	0	1	0
1	Б	400	3161	2019	520	602	20	0		U
1	C	404	Total	С	N	О	S	0	0	0
1		404	3115	1987	513	596	19	U	0	
1	D	410	Total	С	N	О	S	0	2	0
1	ע	410	3172	2025	519	607	21	U		

There are 100 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-24	MET	-	initiating methionine	UNP A0A832SFM5
A	-23	ALA	-	expression tag	UNP A0A832SFM5
A	-22	SER	-	expression tag	UNP A0A832SFM5
A	-21	TRP	-	expression tag	UNP A0A832SFM5
A	-20	SER	-	expression tag	UNP A0A832SFM5
A	-19	HIS	-	expression tag	UNP A0A832SFM5
A	-18	PRO	-	expression tag	UNP A0A832SFM5
A	-17	GLN	-	expression tag	UNP A0A832SFM5
A	-16	PHE	-	expression tag	UNP A0A832SFM5
A	-15	GLU	-	expression tag	UNP A0A832SFM5
A	-14	LYS	-	expression tag	UNP A0A832SFM5
A	-13	SER	-	expression tag	UNP A0A832SFM5
A	-12	GLY	-	expression tag	UNP A0A832SFM5
A	-11	GLY	-	expression tag	UNP A0A832SFM5
A	-10	GLY	-	expression tag	UNP A0A832SFM5
A	-9	GLY	-	expression tag	UNP A0A832SFM5
A	-8	GLY		expression tag	UNP A0A832SFM5
A	-7	GLU	=	expression tag	UNP A0A832SFM5
A	-6	ASN	-	expression tag	UNP A0A832SFM5
A	-5	LEU	-	expression tag	UNP A0A832SFM5

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	TYR	-	expression tag	UNP A0A832SFM5
A	-3	PHE	-	expression tag	UNP A0A832SFM5
A	-2	GLN	_	expression tag	UNP A0A832SFM5
A	-1	GLY	_	expression tag	UNP A0A832SFM5
A	0	HIS	-	expression tag	UNP A0A832SFM5
В	-24	MET	-	initiating methionine	UNP A0A832SFM5
В	-23	ALA	-	expression tag	UNP A0A832SFM5
В	-22	SER	-	expression tag	UNP A0A832SFM5
В	-21	TRP	-	expression tag	UNP A0A832SFM5
В	-20	SER	-	expression tag	UNP A0A832SFM5
В	-19	HIS	-	expression tag	UNP A0A832SFM5
В	-18	PRO	-	expression tag	UNP A0A832SFM5
В	-17	GLN	-	expression tag	UNP A0A832SFM5
В	-16	PHE	-	expression tag	UNP A0A832SFM5
В	-15	GLU	-	expression tag	UNP A0A832SFM5
В	-14	LYS	-	expression tag	UNP A0A832SFM5
В	-13	SER	-	expression tag	UNP A0A832SFM5
В	-12	GLY	-	expression tag	UNP A0A832SFM5
В	-11	GLY	-	expression tag	UNP A0A832SFM5
В	-10	GLY	-	expression tag	UNP A0A832SFM5
В	-9	GLY	_	expression tag	UNP A0A832SFM5
В	-8	GLY	-	expression tag	UNP A0A832SFM5
В	-7	GLU	_	expression tag	UNP A0A832SFM5
В	-6	ASN	-	expression tag	UNP A0A832SFM5
В	-5	LEU	-	expression tag	UNP A0A832SFM5
В	-4	TYR	_	expression tag	UNP A0A832SFM5
В	-3	PHE	-	expression tag	UNP A0A832SFM5
В	-2	GLN	-	expression tag	UNP A0A832SFM5
В	-1	GLY	_	expression tag	UNP A0A832SFM5
В	0	HIS	-	expression tag	UNP A0A832SFM5
С	-24	MET	-	initiating methionine	UNP A0A832SFM5
С	-23	ALA	_	expression tag	UNP A0A832SFM5
С	-22	SER	-	expression tag	UNP A0A832SFM5
С	-21	TRP	_	expression tag	UNP A0A832SFM5
С	-20	SER	-	expression tag	UNP A0A832SFM5
С	-19	HIS	-	expression tag	UNP A0A832SFM5
С	-18	PRO	-	expression tag	UNP A0A832SFM5
С	-17	GLN	-	expression tag	UNP A0A832SFM5
С	-16	PHE		expression tag	UNP A0A832SFM5
С	-15	GLU	-	expression tag	UNP A0A832SFM5
С	-14	LYS	-	expression tag	UNP A0A832SFM5
С	-13	SER	-	expression tag	UNP A0A832SFM5

Continued on next page...




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	-12	GLY	-	expression tag	UNP A0A832SFM5
С	-11	GLY	-	expression tag	UNP A0A832SFM5
С	-10	GLY	-	expression tag	UNP A0A832SFM5
С	-9	GLY	-	expression tag	UNP A0A832SFM5
С	-8	GLY	-	expression tag	UNP A0A832SFM5
С	-7	GLU	-	expression tag	UNP A0A832SFM5
С	-6	ASN	-	expression tag	UNP A0A832SFM5
С	-5	LEU	-	expression tag	UNP A0A832SFM5
С	-4	TYR	-	expression tag	UNP A0A832SFM5
С	-3	PHE	-	expression tag	UNP A0A832SFM5
С	-2	GLN	-	expression tag	UNP A0A832SFM5
С	-1	GLY	-	expression tag	UNP A0A832SFM5
С	0	HIS	-	expression tag	UNP A0A832SFM5
D	-24	MET	_	initiating methionine	UNP A0A832SFM5
D	-23	ALA	-	expression tag	UNP A0A832SFM5
D	-22	SER	_	expression tag	UNP A0A832SFM5
D	-21	TRP	-	expression tag	UNP A0A832SFM5
D	-20	SER	-	expression tag	UNP A0A832SFM5
D	-19	HIS	-	expression tag	UNP A0A832SFM5
D	-18	PRO	-	expression tag	UNP A0A832SFM5
D	-17	GLN	-	expression tag	UNP A0A832SFM5
D	-16	PHE	-	expression tag	UNP A0A832SFM5
D	-15	GLU	-	expression tag	UNP A0A832SFM5
D	-14	LYS	-	expression tag	UNP A0A832SFM5
D	-13	SER	-	expression tag	UNP A0A832SFM5
D	-12	GLY	-	expression tag	UNP A0A832SFM5
D	-11	GLY	-	expression tag	UNP A0A832SFM5
D	-10	GLY	_	expression tag	UNP A0A832SFM5
D	-9	GLY	-	expression tag	UNP A0A832SFM5
D	-8	GLY	_	expression tag	UNP A0A832SFM5
D	-7	GLU	-	expression tag	UNP A0A832SFM5
D	-6	ASN	-	expression tag	UNP A0A832SFM5
D	-5	LEU	-	expression tag	UNP A0A832SFM5
D	-4	TYR	-	expression tag	UNP A0A832SFM5
D	-3	PHE	-	expression tag	UNP A0A832SFM5
D	-2	GLN	-	expression tag	UNP A0A832SFM5
D	-1	GLY	-	expression tag	UNP A0A832SFM5
D	0	HIS	-	expression tag	UNP A0A832SFM5

• Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe S 8 4 4	0	0
2	В	1	Total Fe S 8 4 4	0	0
2	С	1	Total Fe S 8 4 4	0	0
2	D	1	Total Fe S 8 4 4	0	0

• Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe 1 1	0	0
3	В	1	Total Fe 1 1	0	0
3	С	1	Total Fe 1 1	0	0
3	D	1	Total Fe 1 1	0	0


• Molecule 4 is 5'-DEOXY-5'-METHYLTHIOADENOSINE (three-letter code: MTA) (formula: $C_{11}H_{15}N_5O_3S$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	Λ	1	Total	С	N	О	S	0	0
4	A	T	20	11	5	3	1	0	
1	D	1	Total	С	N	О	S	0	0
4	Б	1	20	11	5	3	1	0	
4	С	1	Total	С	N	О	S	0	0
4		1	20	11	5	3	1	0	U
4	D	1	Total	С	N	О	S	0	0
4	ש	1	20	11	5	3	1	U	U

• Molecule 5 is CO-METHYLCOBALAMIN (three-letter code: COB) (formula: $C_{63}H_{91}CoN_{13}O_{14}P$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Atoms					ZeroOcc	AltConf
5	A	1	Total 92	C 63	Co 1	N 13	O 14	P	0	0
	_		Total	C	Co	13 N		P		_
5	В	1	92	63	1	13	14	1	0	0
5	С	1	Total	С	Со	N	0	Р	0	0
		-	92	63	1	13	14	1		
5	D	1	Total	С	Co	N	0	Р	0	0
		_	92	63	1	13	14	1		

• Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Na 1 1	0	0
6	В	1	Total Na 1 1	0	0
6	С	1	Total Na 1 1	0	0
6	D	1	Total Na 1 1	0	0

• Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	359	Total O 359 359	0	0
7	В	335	Total O 335 335	0	0
7	С	235	Total O 235 235	0	0
7	D	339	Total O 339 339	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Methyl coenzyme M reductase-arginine methyltransferase Mmp10



• Molecule 1: Methyl coenzyme M reductase-arginine methyltransferase Mmp10







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	156.06Å 163.91Å 77.81Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.71 - 1.90	Depositor
resolution (A)	44.76 - 1.90	EDS
% Data completeness	87.0 (44.71-1.90)	Depositor
(in resolution range)	86.9 (44.76-1.90)	EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.46 (at 1.89Å)	Xtriage
Refinement program	BUSTER 2.10.4 (16-JUL-2021)	Depositor
R, R_{free}	0.203 , 0.229	Depositor
10, 10 free	0.194 , 0.221	DCC
R_{free} test set	6765 reflections $(4.94%)$	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14382	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 14.28% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MTA, COB, SF4, NA, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.40	0/3238	0.55	0/4381	
1	В	0.40	0/3223	0.54	0/4359	
1	С	0.36	0/3172	0.55	0/4291	
1	D	0.40	0/3238	0.55	0/4381	
All	All	0.39	0/12871	0.55	0/17412	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	3178	0	3200	7	0
1	В	3161	0	3194	7	0
1	С	3115	0	3144	10	0
1	D	3172	0	3201	7	0
2	A	8	0	0	0	0
2	В	8	0	0	0	0
2	С	8	0	0	0	0
2	D	8	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	A	20	0	15	2	0
4	В	20	0	15	2	0
4	С	20	0	15	2	0
4	D	20	0	15	2	0
5	A	92	0	87	9	0
5	В	92	0	87	8	0
5	С	92	0	87	8	0
5	D	92	0	87	6	0
6	A	1	0	0	0	0
6	В	1	0	0	0	0
6	С	1	0	0	0	0
6	D	1	0	0	0	0
7	A	359	0	0	1	0
7	В	335	0	0	1	0
7	С	235	0	0	2	0
7	D	339	0	0	1	0
All	All	14382	0	13147	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ (\rm \AA) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
5:B:504:COB:H552	5:B:504:COB:H531	1.70	0.73
5:D:504:COB:H351	5:D:504:COB:H371	1.72	0.70
5:A:504:COB:H552	5:A:504:COB:H531	1.76	0.67
5:D:504:COB:H552	5:D:504:COB:H531	1.75	0.67
5:B:504:COB:H351	5:B:504:COB:H371	1.78	0.64

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	410/436 (94%)	404 (98%)	5 (1%)	1 (0%)	47	38
1	В	405/436 (93%)	400 (99%)	4 (1%)	1 (0%)	47	38
1	С	400/436 (92%)	396 (99%)	3 (1%)	1 (0%)	41	31
1	D	410/436 (94%)	405 (99%)	4 (1%)	1 (0%)	47	38
All	All	1625/1744~(93%)	1605 (99%)	16 (1%)	4 (0%)	47	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	TYR
1	В	115	TYR
1	С	115	TYR
1	D	115	TYR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Percentiles		
1	A	357/374 (96%)	355 (99%)	2 (1%)	86 87	
1	В	355/374~(95%)	354 (100%)	1 (0%)	92 93	
1	С	350/374 (94%)	350 (100%)	0	100 100	
1	D	357/374 (96%)	355 (99%)	2 (1%)	86 87	
All	All	1419/1496 (95%)	1414 (100%)	5 (0%)	91 91	



All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	275	GLU
1	A	354	LYS
1	В	282	ARG
1	D	53	LYS
1	D	326	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	С	233	HIS
1	С	402	GLN
1	D	233	HIS
1	D	215	GLN
1	В	293	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 8 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Tuna	Chain	Res	Link	Boı	Bond lengths		Во	nd angle	es
10101	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MTA	D	503	-	19,22,22	0.64	0	19,32,32	0.95	2 (10%)
5	COB	D	504	-	87,102,102	0.98	6 (6%)	127,170,170	1.37	15 (11%)
4	MTA	В	503	-	19,22,22	0.63	0	19,32,32	1.04	2 (10%)
5	COB	A	504	-	87,102,102	0.95	5 (5%)	127,170,170	1.39	12 (9%)
5	COB	В	504	-	87,102,102	0.94	4 (4%)	127,170,170	1.38	17 (13%)
4	MTA	A	503	-	19,22,22	0.61	0	19,32,32	0.89	1 (5%)
2	SF4	D	501	1	0,12,12	-	-	-		
2	SF4	В	501	1	0,12,12	-	-	-		
2	SF4	С	501	1	0,12,12	-	-	-		
5	COB	С	504	-	87,102,102	0.94	4 (4%)	127,170,170	1.40	14 (11%)
4	MTA	С	503	-	19,22,22	0.61	0	19,32,32	1.00	2 (10%)
2	SF4	A	501	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MTA	D	503	-	-	3/3/23/23	0/3/3/3
5	COB	D	504	-	-	9/52/231/231	0/3/11/11
4	MTA	В	503	-	-	3/3/23/23	0/3/3/3
5	COB	A	504	-	-	11/52/231/231	0/3/11/11
5	COB	В	504	-	-	11/52/231/231	0/3/11/11
4	MTA	A	503	-	-	3/3/23/23	0/3/3/3
2	SF4	В	501	1	-	-	0/6/5/5
2	SF4	D	501	1	-	-	0/6/5/5
2	SF4	С	501	1	-	-	0/6/5/5
5	COB	С	504	-	-	12/52/231/231	0/3/11/11
4	MTA	С	503	-	-	3/3/23/23	0/3/3/3
2	SF4	A	501	1	-	-	0/6/5/5

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\mathring{A}})$	$\operatorname{Ideal}(\text{\AA})$
5	D	504	COB	C8B-C9B	4.15	1.49	1.40
5	С	504	COB	C8B-C9B	4.13	1.48	1.40
5	A	504	COB	C8B-C9B	4.08	1.48	1.40
5	В	504	COB	C8B-C9B	4.07	1.48	1.40

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
5	С	504	COB	C6B-C5B	3.72	1.50	1.40

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	A	504	COB	C20-C1-C19	-5.79	102.26	109.42
5	В	504	COB	C20-C1-C19	-5.34	102.82	109.42
5	В	504	COB	C60-C18-C19	5.31	122.72	112.39
5	С	504	COB	C12-C11-C10	-5.25	118.74	123.54
5	A	504	COB	C12-C11-C10	-5.24	118.75	123.54

There are no chirality outliers.

5 of 55 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	MTA	O4'-C4'-C5'-S5'
4	В	503	MTA	O4'-C4'-C5'-S5'
4	В	503	MTA	C3'-C4'-C5'-S5'
4	С	503	MTA	O4'-C4'-C5'-S5'
4	D	503	MTA	O4'-C4'-C5'-S5'

There are no ring outliers.


8 monomers are involved in 39 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	503	MTA	2	0
5	D	504	COB	6	0
4	В	503	MTA	2	0
5	A	504	COB	9	0
5	В	504	COB	8	0
4	A	503	MTA	2	0
5	С	504	COB	8	0
4	С	503	MTA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the




average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	411/436 (94%)	0.71	19 (4%) 32 35	25, 37, 58, 80	0
1	В	408/436 (93%)	0.90	49 (12%) 4 4	25, 39, 71, 91	0
1	С	404/436 (92%)	1.15	69 (17%) 1 1	33, 50, 85, 112	0
1	D	410/436 (94%)	0.56	24 (5%) 22 25	25, 37, 59, 77	0
All	All	1633/1744 (93%)	0.83	161 (9%) 7 8	25, 40, 70, 112	0

The worst 5 of 161 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	330	ALA	11.2
1	A	77	SER	11.2
1	В	73	LEU	10.6
1	D	77	SER	10.5
1	С	30	ILE	9.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q<0.9
4	MTA	D	503	20/20	0.85	0.19	29,32,46,46	0
4	MTA	С	503	20/20	0.86	0.17	35,38,50,50	0
4	MTA	A	503	20/20	0.89	0.16	27,30,42,43	0
4	MTA	В	503	20/20	0.92	0.15	28,32,44,44	0
5	COB	С	504	92/92	0.93	0.14	31,33,36,37	0
6	NA	С	505	1/1	0.94	0.11	38,38,38,38	0
5	COB	A	504	92/92	0.95	0.14	24,27,32,33	0
5	COB	D	504	92/92	0.95	0.14	24,27,31,34	0
5	COB	В	504	92/92	0.95	0.14	24,27,32,34	0
2	SF4	A	501	8/8	0.96	0.12	26,27,29,29	0
2	SF4	В	501	8/8	0.96	0.11	29,29,30,30	0
2	SF4	С	501	8/8	0.96	0.09	37,37,37,37	0
2	SF4	D	501	8/8	0.97	0.10	28,29,30,30	0
6	NA	D	505	1/1	0.97	0.09	30,30,30,30	0
6	NA	В	505	1/1	0.98	0.11	30,30,30,30	0
3	FE	В	502	1/1	0.98	0.07	46,46,46,46	0
3	FE	С	502	1/1	0.98	0.07	52,52,52,52	0
6	NA	A	505	1/1	0.99	0.09	31,31,31,31	0
3	FE	A	502	1/1	1.00	0.09	32,32,32,32	0
3	FE	D	502	1/1	1.00	0.11	31,31,31,31	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MTA D 503:


 $2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)














































Electron density around SF4 A 501:









Electron density around SF4 B 501:












Electron density around SF4 C 501:








Electron density around SF4 D 501: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around NA D 505: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around FE C 502:










Electron density around NA A 505: $2 \mathrm{mF}_o\text{-DF}_c \text{ (at } 0.7 \text{ rmsd) in gray} \\ \mathrm{mF}_o\text{-DF}_c \text{ (at } 3 \text{ rmsd) in purple (negative)}$

and green (positive)











Electron density around FE A 502:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

