

# wwPDB X-ray Structure Validation Summary Report (i)

Jan 27, 2022 – 03:16 pm GMT

PDB ID : 7QBV

Title: B12-dependent radical SAM methyltransferase, Mmp10 with [4Fe-4S] cluster,

cobalamin, and S-adenosyl-L-homocysteine bound.

Authors: Fyfe, C.D.; Chavas, L.M.G.; Legrand, P.; Benjdia, A.; Berteau, O.

Deposited on : 2021-11-19

Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.26

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0267

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.26

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution

Metric	Whole archive	Similar resolution
Metric	$(\#  ext{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	436	88%	6%	6%
1	В	436	88%	6%	6%
1	С	436	87%	7%	6%
1	D	436	86%	7%	7%



# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 13237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Methyl coenzyme M reductase-arginine methyltransferase Mmp10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	411	Total	С	N	О	S	0	0	0
1	A	411	3172	2024	520	608	20	0	0	
1	В	411	Total	С	N	О	S	0	0	0
1		411	3172	2024	520	608	20	U		0
1	C	C 411	Total	С	N	О	S	0	0	0
1			3172	2024	520	608	20	U		
1	1 D	407	Total	С	N	О	S	0		0
1			3146	2010	516	600	20		U	

There are 100 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-24	MET	-	initiating methionine	UNP A0A832SFM5
A	-23	ALA	-	expression tag	UNP A0A832SFM5
A	-22	SER	-	expression tag	UNP A0A832SFM5
A	-21	TRP	-	expression tag	UNP A0A832SFM5
A	-20	SER	-	expression tag	UNP A0A832SFM5
A	-19	HIS	-	expression tag	UNP A0A832SFM5
A	-18	PRO	-	expression tag	UNP A0A832SFM5
A	-17	GLN	-	expression tag	UNP A0A832SFM5
A	-16	PHE	-	expression tag	UNP A0A832SFM5
A	-15	GLU	-	expression tag	UNP A0A832SFM5
A	-14	LYS	-	expression tag	UNP A0A832SFM5
A	-13	SER	-	expression tag	UNP A0A832SFM5
A	-12	GLY	-	expression tag	UNP A0A832SFM5
A	-11	GLY	-	expression tag	UNP A0A832SFM5
A	-10	GLY	-	expression tag	UNP A0A832SFM5
A	-9	GLY	-	expression tag	UNP A0A832SFM5
A	-8	GLY	-	expression tag	UNP A0A832SFM5
A	-7	GLU	-	expression tag	UNP A0A832SFM5
A	-6	ASN	-	expression tag	UNP A0A832SFM5
A	-5	LEU	-	expression tag	UNP A0A832SFM5

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	TYR	-	expression tag	UNP A0A832SFM5
A	-3	PHE	_	expression tag	UNP A0A832SFM5
A	-2	GLN	-	expression tag	UNP A0A832SFM5
A	-1	GLY	-	expression tag	UNP A0A832SFM5
A	0	HIS	-	expression tag	UNP A0A832SFM5
В	-24	MET	-	initiating methionine	UNP A0A832SFM5
В	-23	ALA	-	expression tag	UNP A0A832SFM5
В	-22	SER	-	expression tag	UNP A0A832SFM5
В	-21	TRP	-	expression tag	UNP A0A832SFM5
В	-20	SER	-	expression tag	UNP A0A832SFM5
В	-19	HIS	-	expression tag	UNP A0A832SFM5
В	-18	PRO	-	expression tag	UNP A0A832SFM5
В	-17	GLN	-	expression tag	UNP A0A832SFM5
В	-16	PHE	-	expression tag	UNP A0A832SFM5
В	-15	GLU	-	expression tag	UNP A0A832SFM5
В	-14	LYS	-	expression tag	UNP A0A832SFM5
В	-13	SER	-	expression tag	UNP A0A832SFM5
В	-12	GLY	-	expression tag	UNP A0A832SFM5
В	-11	GLY	_	expression tag	UNP A0A832SFM5
В	-10	GLY	-	expression tag	UNP A0A832SFM5
В	-9	GLY	-	expression tag	UNP A0A832SFM5
В	-8	GLY	-	expression tag	UNP A0A832SFM5
В	-7	GLU	-	expression tag	UNP A0A832SFM5
В	-6	ASN	-	expression tag	UNP A0A832SFM5
В	-5	LEU	-	expression tag	UNP A0A832SFM5
В	-4	TYR	_	expression tag	UNP A0A832SFM5
В	-3	PHE	-	expression tag	UNP A0A832SFM5
В	-2	GLN	-	expression tag	UNP A0A832SFM5
В	-1	GLY	-	expression tag	UNP A0A832SFM5
В	0	HIS	-	expression tag	UNP A0A832SFM5
С	-24	MET	-	initiating methionine	UNP A0A832SFM5
С	-23	ALA	-	expression tag	UNP A0A832SFM5
С	-22	SER	-	expression tag	UNP A0A832SFM5
С	-21	TRP	-	expression tag	UNP A0A832SFM5
С	-20	SER	-	expression tag	UNP A0A832SFM5
С	-19	HIS	-	expression tag	UNP A0A832SFM5
С	-18	PRO	-	expression tag	UNP A0A832SFM5
С	-17	GLN	-	expression tag	UNP A0A832SFM5
С	-16	PHE	-	expression tag	UNP A0A832SFM5
С	-15	GLU	-	expression tag	UNP A0A832SFM5
С	-14	LYS	-	expression tag	UNP A0A832SFM5
С	-13	SER	-	expression tag	UNP A0A832SFM5

Continued on next page...




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	-12	GLY	-	expression tag	UNP A0A832SFM5
С	-11	GLY	-	expression tag	UNP A0A832SFM5
С	-10	GLY	-	expression tag	UNP A0A832SFM5
С	-9	GLY	-	expression tag	UNP A0A832SFM5
С	-8	GLY	-	expression tag	UNP A0A832SFM5
С	-7	GLU	-	expression tag	UNP A0A832SFM5
С	-6	ASN	-	expression tag	UNP A0A832SFM5
С	-5	LEU	-	expression tag	UNP A0A832SFM5
С	-4	TYR	-	expression tag	UNP A0A832SFM5
С	-3	PHE	-	expression tag	UNP A0A832SFM5
С	-2	GLN	-	expression tag	UNP A0A832SFM5
С	-1	GLY	-	expression tag	UNP A0A832SFM5
С	0	HIS	-	expression tag	UNP A0A832SFM5
D	-24	MET	_	initiating methionine	UNP A0A832SFM5
D	-23	ALA	-	expression tag	UNP A0A832SFM5
D	-22	SER	_	expression tag	UNP A0A832SFM5
D	-21	TRP	-	expression tag	UNP A0A832SFM5
D	-20	SER	-	expression tag	UNP A0A832SFM5
D	-19	HIS	-	expression tag	UNP A0A832SFM5
D	-18	PRO	-	expression tag	UNP A0A832SFM5
D	-17	GLN	-	expression tag	UNP A0A832SFM5
D	-16	PHE	-	expression tag	UNP A0A832SFM5
D	-15	GLU	-	expression tag	UNP A0A832SFM5
D	-14	LYS	-	expression tag	UNP A0A832SFM5
D	-13	SER	-	expression tag	UNP A0A832SFM5
D	-12	GLY	-	expression tag	UNP A0A832SFM5
D	-11	GLY	-	expression tag	UNP A0A832SFM5
D	-10	$\operatorname{GLY}$	_	expression tag	UNP A0A832SFM5
D	-9	GLY	-	expression tag	UNP A0A832SFM5
D	-8	$\operatorname{GLY}$	_	expression tag	UNP A0A832SFM5
D	-7	GLU	-	expression tag	UNP A0A832SFM5
D	-6	ASN	-	expression tag	UNP A0A832SFM5
D	-5	LEU	-	expression tag	UNP A0A832SFM5
D	-4	TYR	-	expression tag	UNP A0A832SFM5
D	-3	PHE	-	expression tag	UNP A0A832SFM5
D	-2	GLN	-	expression tag	UNP A0A832SFM5
D	-1	GLY	-	expression tag	UNP A0A832SFM5
D	0	HIS	-	expression tag	UNP A0A832SFM5

• Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	Λ	1	Total Fe S	0	0	
	2 A	1	8 4 4	U	0	
2	В	1	Total Fe S	0	0	
	2 D	1	8 4 4	0		
2	С	1	Total Fe S	0	0	
		1	8 4 4	0	0	
2	D	1	Total Fe S	0	0	
			8 4 4	0	U	

• Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe 1 1	0	0
3	В	1	Total Fe 1 1	0	0
3	С	1	Total Fe 1 1	0	0
3	D	1	Total Fe 1 1	0	0


• Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	Λ	1	Total	С	N	О	S	0	0	
4	A		26	14	6	5	1	0		
1	D	1	Total	С	N	О	S	0	0	
4	4   B	1	26	14	6	5	1	0		
4	С	C 1	Total	С	N	О	S	0	0	
4		1	26	14	6	5	1	0	0	
4	4 D	1	Total	С	N	О	S	0	0	
4			26	14	6	5	1	U	0	

 $\bullet \ \ \mathrm{Molecule} \ 5 \ \mathrm{is} \ \mathrm{CO-METHYLCOBALAMIN} \ (\mathrm{three-letter} \ \mathrm{code} \colon \ \mathrm{COB}) \ (\mathrm{formula} \colon \ \mathrm{C}_{63}\mathrm{H}_{91}\mathrm{CoN}_{13}\mathrm{O}_{14}\mathrm{P}).$ 





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf								
5	A	1	Total 92	C 63	Co 1	N 13	O 14	P	0	0						
	_		Total	C	Co	13 N		P		_						
5	В	В	В	В	В	В	В		92	63	1	13	14	1	0	U
5	С	1	Total	С	Со	N	0	Р	0	0						
		-	92	63	1	13	14	1	Ü							
5	D	1	Total	С	Co	N	0	Р	0	0						
		_	92	63	1	13	14	1								

• Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Na 1 1	0	0
6	В	1	Total Na 1 1	0	0
6	С	1	Total Na 1 1	0	0
6	D	1	Total Na 1 1	0	0

#### • Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	14	Total O 14 14	0	0
7	В	18	Total O 18 18	0	0
7	С	16	Total O 16 16	0	0
7	D	15	Total O 15 15	0	0




# 3 Residue-property plots (i)

ALA ALA ALA ALA ALA ALA ALA ALA SER TTRP PHI S SER PHI S CLU PHI S SER GLV GLV GLV GLV GLV GLV GLV ALS HI S HI S HI S HI S SHI HI

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Methyl coenzyme M reductase-arginine methyltransferase Mmp10



• Molecule 1: Methyl coenzyme M reductase-arginine methyltransferase Mmp10

Chain D: 86% 7% 7%







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	155.50Å 163.00Å 77.33Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	49.40 - 2.70	Depositor
Resolution (A)	49.40 - 2.70	EDS
% Data completeness	76.7 (49.40-2.70)	Depositor
(in resolution range)	$76.8 \ (49.40 - 2.70)$	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.60 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.10.4 (20-APR-2021)	Depositor
D.D.	0.214 , 0.235	Depositor
$R, R_{free}$	0.210 , $0.231$	DCC
$R_{free}$ test set	2062 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.1	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning <sup>2</sup>	$< L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	13237	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 14.42% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, SF4, COB, NA, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.28	0/3232	0.49	0/4373	
1	В	0.28	0/3232	0.48	0/4373	
1	С	0.28	0/3232	0.48	0/4373	
1	D	0.26	0/3205	0.48	0/4335	
All	All	0.28	0/12901	0.48	0/17454	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3172	0	3198	19	0
1	В	3172	0	3197	17	0
1	С	3172	0	3197	16	0
1	D	3146	0	3175	19	0
2	A	8	0	0	1	0
2	В	8	0	0	1	0
2	С	8	0	0	1	0
2	D	8	0	0	1	0
3	A	1	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	A	26	0	19	1	0
4	В	26	0	19	1	0
4	С	26	0	19	1	0
4	D	26	0	19	0	0
5	A	92	0	87	10	0
5	В	92	0	87	11	0
5	С	92	0	87	9	0
5	D	92	0	87	10	0
6	A	1	0	0	0	0
6	В	1	0	0	0	0
6	С	1	0	0	0	0
6	D	1	0	0	0	0
7	A	14	0	0	0	0
7	В	18	0	0	1	0
7	С	16	0	0	0	0
7	D	15	0	0	0	0
All	All	13237	0	13191	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ (\rm \mathring{A}) \end{array}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
5:C:504:COB:H601	5:C:504:COB:H262	1.58	0.85
5:A:504:COB:H601	5:A:504:COB:H262	1.59	0.85
5:B:504:COB:H601	5:B:504:COB:H262	1.59	0.84
5:D:504:COB:H601	5:D:504:COB:H262	1.58	0.84
1:B:338:GLU:HA	1:B:364:ARG:HG2	1.64	0.77

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	A	409/436 (94%)	405 (99%)	3 (1%)	1 (0%)	47	73
1	В	409/436 (94%)	404 (99%)	4 (1%)	1 (0%)	47	73
1	С	409/436 (94%)	405 (99%)	3 (1%)	1 (0%)	47	73
1	D	403/436 (92%)	398 (99%)	4 (1%)	1 (0%)	47	73
All	All	1630/1744 (94%)	1612 (99%)	14 (1%)	4 (0%)	47	73

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	TYR
1	В	115	TYR
1	С	115	TYR
1	D	115	TYR

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles		
1	A	356/374~(95%)	353 (99%)	3 (1%)	81	93	
1	В	356/374~(95%)	354 (99%)	2 (1%)	86	95	
1	С	356/374 (95%)	354 (99%)	2 (1%)	86	95	
1	D	353/374 (94%)	352 (100%)	1 (0%)	92	98	
All	All	1421/1496 (95%)	1413 (99%)	8 (1%)	86	95	



5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	67	GLU
1	С	354	LYS
1	В	354	LYS
1	В	278	LEU
1	С	192	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	215	GLN
1	A	272	ASN
1	В	215	GLN
1	D	215	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 8 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Trino	Chain	Dag	Link	Boı	nd lengt	hs	Во	nd angle	es
Mol	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SF4	С	501	1	0,12,12	-	-	-		
4	SAH	A	503	-	21,28,28	0.61	0	20,40,40	1.00	2 (10%)
4	SAH	С	503	-	21,28,28	0.61	0	20,40,40	1.26	2 (10%)
2	SF4	A	501	1	0,12,12	-	-	-		
5	COB	A	504	-	87,102,102	0.95	5 (5%)	127,170,170	1.35	13 (10%)
2	SF4	D	501	1	0,12,12	-	-	-		
2	SF4	В	501	1	0,12,12	-	-	-		
4	SAH	D	503	-	21,28,28	0.61	0	20,40,40	1.15	2 (10%)
5	COB	С	504	-	87,102,102	0.94	5 (5%)	127,170,170	1.34	13 (10%)
4	SAH	В	503	-	21,28,28	0.61	0	20,40,40	0.96	2 (10%)
5	COB	В	504	-	87,102,102	0.97	5 (5%)	127,170,170	1.36	13 (10%)
5	COB	D	504	-	87,102,102	0.96	5 (5%)	127,170,170	1.36	11 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAH	A	503	-	-	2/7/31/31	0/3/3/3
4	SAH	С	503	-	-	0/7/31/31	0/3/3/3
2	SF4	С	501	1	-	-	0/6/5/5
2	SF4	A	501	1	-	-	0/6/5/5
5	COB	A	504	-	-	14/52/231/231	0/3/11/11
2	SF4	В	501	1	-	-	0/6/5/5
5	COB	С	504	-	-	14/52/231/231	0/3/11/11
4	SAH	D	503	-	-	4/7/31/31	0/3/3/3
2	SF4	D	501	1	-	-	0/6/5/5
4	SAH	В	503	ı	-	2/7/31/31	0/3/3/3
5	COB	В	504	-	-	14/52/231/231	0/3/11/11
5	COB	D	504	-	-	14/52/231/231	0/3/11/11

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
5	D	504	COB	C8B-C9B	4.42	1.49	1.40
5	В	504	COB	C8B-C9B	4.37	1.49	1.40
5	A	504	COB	C8B-C9B	4.20	1.49	1.40
5	С	504	COB	C8B-C9B	4.18	1.49	1.40

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	Ideal(A)	
5	D	504	COB	C6B-C5B	3.71	1.50	1.40	

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
5	В	504	COB	C12-C11-C10	-5.79	118.25	123.54
5	A	504	COB	C12-C11-C10	-5.71	118.33	123.54
5	D	504	COB	C12-C11-C10	-5.68	118.35	123.54
5	С	504	COB	C12-C11-C10	-5.51	118.51	123.54
5	В	504	COB	C20-C1-C19	-4.04	104.42	109.42

There are no chirality outliers.

5 of 64 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	SAH	O4'-C4'-C5'-SD
4	A	503	SAH	C3'-C4'-C5'-SD
4	В	503	SAH	O4'-C4'-C5'-SD
4	В	503	SAH	C3'-C4'-C5'-SD
4	D	503	SAH	O4'-C4'-C5'-SD

There are no ring outliers.


11 monomers are involved in 47 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	501	SF4	1	0
4	A	503	SAH	1	0
4	С	503	SAH	1	0
2	A	501	SF4	1	0
5	A	504	COB	10	0
2	D	501	SF4	1	0
2	В	501	SF4	1	0
5	С	504	COB	9	0
4	В	503	SAH	1	0
5	В	504	COB	11	0
5	D	504	COB	10	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is




within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









































# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSl	RZ>	>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	411/436 (94%)	-0.10	6 (1%)	73	76	38, 53, 75, 91	0
1	В	411/436 (94%)	-0.04	6 (1%)	73	76	38, 55, 82, 108	0
1	С	411/436 (94%)	0.04	17 (4%)	37	36	40, 56, 89, 112	0
1	D	407/436 (93%)	0.22	17 (4%)	36	35	45, 67, 101, 148	0
All	All	1640/1744 (94%)	0.03	46 (2%)	53	54	38, 57, 92, 148	0

The worst 5 of 46 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	411	GLU	7.5
1	A	77	SER	6.8
1	D	73	LEU	6.6
1	В	77	SER	6.1
1	С	77	SER	5.5

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q < 0.9
4	SAH	С	503	26/26	0.90	0.19	50,54,61,62	0
4	SAH	D	503	26/26	0.91	0.21	56,60,67,67	0
4	SAH	A	503	26/26	0.92	0.20	44,48,55,56	0
6	NA	С	505	1/1	0.92	0.07	42,42,42,42	0
6	NA	A	505	1/1	0.93	0.13	42,42,42,42	0
4	SAH	В	503	26/26	0.93	0.16	44,48,52,52	0
5	COB	С	504	92/92	0.96	0.19	41,42,44,45	0
5	COB	D	504	92/92	0.96	0.19	46,48,49,50	0
5	COB	A	504	92/92	0.97	0.17	37,40,43,44	0
6	NA	В	505	1/1	0.97	0.11	39,39,39,39	0
5	COB	В	504	92/92	0.97	0.18	38,40,48,48	0
3	FE	С	502	1/1	0.98	0.09	67,67,67,67	0
2	SF4	С	501	8/8	0.99	0.14	49,49,49,49	0
2	SF4	D	501	8/8	0.99	0.13	57,58,58,58	0
2	SF4	A	501	8/8	0.99	0.13	42,42,42,43	0
3	FE	D	502	1/1	0.99	0.11	74,74,74,74	0
2	SF4	В	501	8/8	0.99	0.13	40,40,40,40	0
6	NA	D	505	1/1	0.99	0.07	36,36,36,36	0
3	FE	A	502	1/1	1.00	0.11	42,42,42,42	0
3	FE	В	502	1/1	1.00	0.12	47,47,47,47	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around SAH C 503:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around SAH D 503:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)













 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)
















#### Electron density around COB A 504:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)





















# Electron density around SF4 A 501:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









# 6.5 Other polymers (i)

There are no such residues in this entry.

