



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 7, 2023 – 08:35 PM EDT

PDB ID : 1QGN
Title : CYSTATHIONINE GAMMA-SYNTHASE FROM NICOTIANA TABACUM
Authors : Steegborn, C.; Messerschmidt, A.; Laber, B.; Streber, W.; Huber, R.; Clausen, T.
Deposited on : 1999-05-02
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	51% 33% 5% 11%
1	B	445	55% 29% 5% 11%
1	C	445	51% 33% 5% 11%
1	D	445	53% 30% 6% 11%
1	E	445	56% 29% • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain				
1	F	445		52%	32%	6%	11%
1	G	445		54%	30%	5%	11%
1	H	445		55%	30%	5%	11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (CYSTATHIONINE GAMMA-SYNTHASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	398	3032	1933	514	569	16	0	0	0
1	B	398	3032	1933	514	569	16	0	0	0
1	C	398	3032	1933	514	569	16	0	0	0
1	D	398	3032	1933	514	569	16	0	0	0
1	E	398	3032	1933	514	569	16	0	0	0
1	F	398	3032	1933	514	569	16	0	0	0
1	G	398	3032	1933	514	569	16	0	0	0
1	H	398	3032	1933	514	569	16	0	0	0

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	B	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	C	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	D	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	E	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	F	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	G	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	H	1	Total 15	C 8	N 1	O 5	P 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	44	Total 44	O 44	0	0
3	B	41	Total 41	O 41	0	0
3	C	41	Total 41	O 41	0	0
3	D	37	Total 37	O 37	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	39	Total O 39 39	0	0
3	F	42	Total O 42 42	0	0
3	G	41	Total O 41 41	0	0
3	H	41	Total O 41 41	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTASE)




- Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTASE)






• Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTASE)




• Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTASE)




• Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTASE)

Chain E: 56% 29% 11%




• Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTASE)


Chain F: 52% 32% 6% 11%



• Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTASE)


Chain G: 54% 30% 5% 11%





• Molecule 1: PROTEIN (CYSTATHIONINE GAMMA-SYNTHASE)

Chain H: 55% 30% 5% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	120.00Å 129.50Å 309.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.90 24.95 – 2.79	Depositor EDS
% Data completeness (in resolution range)	88.2 (8.00-2.90) 86.2 (24.95-2.79)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.80Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.201 , 0.250 0.231 , 0.241	Depositor DCC
R_{free} test set	5276 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtrriage
Anisotropy	0.383	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 29.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	24702	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/3091	0.59	1/4190 (0.0%)
1	B	0.35	0/3091	0.58	1/4190 (0.0%)
1	C	0.33	0/3091	0.58	1/4190 (0.0%)
1	D	0.35	0/3091	0.59	1/4190 (0.0%)
1	E	0.35	0/3091	0.58	1/4190 (0.0%)
1	F	0.34	0/3091	0.57	1/4190 (0.0%)
1	G	0.34	0/3091	0.58	1/4190 (0.0%)
1	H	0.34	0/3091	0.58	1/4190 (0.0%)
All	All	0.34	0/24728	0.58	8/33520 (0.0%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	414	LYS	CD-CE-NZ	5.57	124.50	111.70
1	F	414	LYS	CD-CE-NZ	5.56	124.48	111.70
1	B	414	LYS	CD-CE-NZ	5.52	124.40	111.70
1	E	414	LYS	CD-CE-NZ	5.50	124.35	111.70
1	D	414	LYS	CD-CE-NZ	5.47	124.29	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3032	0	3054	125	0
1	B	3032	0	3055	117	0
1	C	3032	0	3055	129	0
1	D	3032	0	3055	120	0
1	E	3032	0	3055	116	0
1	F	3032	0	3055	131	0
1	G	3032	0	3055	121	0
1	H	3032	0	3055	114	0
2	A	15	0	6	0	0
2	B	15	0	6	0	0
2	C	15	0	6	0	0
2	D	15	0	6	0	0
2	E	15	0	6	0	0
2	F	15	0	6	0	0
2	G	15	0	6	0	0
2	H	15	0	6	0	0
3	A	44	0	0	2	0
3	B	41	0	0	3	0
3	C	41	0	0	3	0
3	D	37	0	0	2	0
3	E	39	0	0	1	0
3	F	42	0	0	3	0
3	G	41	0	0	1	0
3	H	41	0	0	3	0
All	All	24702	0	24487	925	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 925 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:241:THR:HG22	1:C:243:LEU:H	1.22	1.02
1:H:241:THR:HG22	1:H:243:LEU:H	1.25	1.02
1:B:241:THR:HG22	1:B:243:LEU:H	1.23	1.01
1:E:241:THR:HG22	1:E:243:LEU:H	1.20	1.00
1:G:241:THR:HG22	1:G:243:LEU:H	1.26	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/445 (89%)	360 (91%)	31 (8%)	5 (1%)	12	37
1	B	396/445 (89%)	363 (92%)	29 (7%)	4 (1%)	15	45
1	C	396/445 (89%)	361 (91%)	32 (8%)	3 (1%)	19	51
1	D	396/445 (89%)	364 (92%)	28 (7%)	4 (1%)	15	45
1	E	396/445 (89%)	363 (92%)	30 (8%)	3 (1%)	19	51
1	F	396/445 (89%)	356 (90%)	35 (9%)	5 (1%)	12	37
1	G	396/445 (89%)	363 (92%)	31 (8%)	2 (0%)	29	61
1	H	396/445 (89%)	364 (92%)	29 (7%)	3 (1%)	19	51
All	All	3168/3560 (89%)	2894 (91%)	245 (8%)	29 (1%)	17	48

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	188	ASP
1	B	188	ASP
1	C	188	ASP
1	D	320	THR
1	E	188	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/364 (90%)	275 (84%)	52 (16%)	2	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	327/364 (90%)	280 (86%)	47 (14%)	3	9
1	C	327/364 (90%)	278 (85%)	49 (15%)	3	9
1	D	327/364 (90%)	276 (84%)	51 (16%)	2	8
1	E	327/364 (90%)	282 (86%)	45 (14%)	3	10
1	F	327/364 (90%)	277 (85%)	50 (15%)	2	8
1	G	327/364 (90%)	284 (87%)	43 (13%)	4	12
1	H	327/364 (90%)	277 (85%)	50 (15%)	2	8
All	All	2616/2912 (90%)	2229 (85%)	387 (15%)	3	9

5 of 387 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	396	VAL
1	F	403	SER
1	E	420	ASN
1	F	192	LEU
1	G	117	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	420	ASN
1	G	63	HIS
1	H	420	ASN
1	F	420	ASN
1	G	343	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	A	500	1	15,15,16	3.62	6 (40%)	20,22,23	2.29	8 (40%)
2	PLP	F	500	1	15,15,16	3.43	5 (33%)	20,22,23	2.33	8 (40%)
2	PLP	B	500	1	15,15,16	3.54	4 (26%)	20,22,23	2.33	8 (40%)
2	PLP	E	500	1	15,15,16	3.43	5 (33%)	20,22,23	2.27	9 (45%)
2	PLP	D	500	1	15,15,16	3.47	5 (33%)	20,22,23	2.26	8 (40%)
2	PLP	H	500	1	15,15,16	3.60	4 (26%)	20,22,23	2.31	8 (40%)
2	PLP	G	500	1	15,15,16	3.72	4 (26%)	20,22,23	2.28	7 (35%)
2	PLP	C	500	1	15,15,16	3.83	5 (33%)	20,22,23	2.32	8 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	500	1	-	0/6/6/8	0/1/1/1
2	PLP	F	500	1	-	0/6/6/8	0/1/1/1
2	PLP	B	500	1	-	0/6/6/8	0/1/1/1
2	PLP	E	500	1	-	0/6/6/8	0/1/1/1
2	PLP	D	500	1	-	0/6/6/8	0/1/1/1
2	PLP	H	500	1	-	0/6/6/8	0/1/1/1
2	PLP	G	500	1	-	0/6/6/8	0/1/1/1
2	PLP	C	500	1	-	0/6/6/8	0/1/1/1

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	500	PLP	C5-C4	11.27	1.53	1.40
2	B	500	PLP	C5-C4	10.99	1.52	1.40
2	C	500	PLP	C5-C4	10.97	1.52	1.40
2	F	500	PLP	C5-C4	10.88	1.52	1.40
2	H	500	PLP	C5-C4	10.78	1.52	1.40

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	500	PLP	C2A-C2-C3	6.16	128.50	120.89
2	G	500	PLP	C2A-C2-C3	6.07	128.39	120.89
2	A	500	PLP	C2A-C2-C3	6.05	128.36	120.89
2	F	500	PLP	C2A-C2-C3	5.86	128.13	120.89
2	E	500	PLP	C2A-C2-C3	5.84	128.10	120.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	398/445 (89%)	-0.19	0 100 100	12, 29, 57, 95	0
1	B	398/445 (89%)	-0.36	0 100 100	9, 28, 57, 92	0
1	C	398/445 (89%)	-0.27	1 (0%) 94 94	11, 31, 61, 95	0
1	D	398/445 (89%)	-0.32	0 100 100	9, 28, 58, 91	0
1	E	398/445 (89%)	-0.11	1 (0%) 94 94	9, 28, 57, 87	0
1	F	398/445 (89%)	-0.42	0 100 100	8, 31, 60, 89	0
1	G	398/445 (89%)	-0.10	1 (0%) 94 94	7, 31, 60, 89	0
1	H	398/445 (89%)	-0.38	0 100 100	9, 29, 56, 97	0
All	All	3184/3560 (89%)	-0.27	3 (0%) 95 96	7, 30, 58, 97	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	415	TYR	3.1
1	E	415	TYR	2.8
1	G	400	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PLP	A	500	15/16	0.93	0.23	14,29,35,36	0
2	PLP	C	500	15/16	0.94	0.20	19,31,37,38	0
2	PLP	G	500	15/16	0.94	0.23	20,30,39,42	0
2	PLP	D	500	15/16	0.96	0.17	13,23,28,30	0
2	PLP	H	500	15/16	0.96	0.19	17,30,34,36	0
2	PLP	B	500	15/16	0.97	0.15	8,26,32,34	0
2	PLP	E	500	15/16	0.97	0.19	15,28,36,37	0
2	PLP	F	500	15/16	0.98	0.15	17,33,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.