



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2024 – 09:42 AM EDT

PDB ID : 2QJG
Title : M. jannaschii ADH synthase complexed with F1,6P
Authors : Ealick, S.E.; Morar, M.
Deposited on : 2007-07-07
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	 3% 74% 22% .
1	B	273	 3% 74% 19% 5% .
1	C	273	 % 73% 20% . .
1	D	273	 2% 76% 19% . .
1	E	273	 % 75% 19% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	273		71% 21%
1	G	273		75% 19%
1	H	273		76% 18%
1	I	273		77% 17%
1	J	273		73% 23%
1	K	273		70% 24%
1	L	273		71% 24%
1	M	273		74% 21%
1	N	273		73% 21%
1	O	273		75% 19%
1	P	273		68% 26%
1	Q	273		75% 21%
1	R	273		75% 20%
1	S	273		71% 22%
1	T	273		69% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	F2P	A	501	X	-	-	-
2	F2P	B	501	X	-	-	-
2	F2P	C	501	X	-	-	-
2	F2P	D	501	X	-	-	-
2	F2P	E	501	X	-	-	-
2	F2P	F	501	X	-	-	-
2	F2P	G	501	X	-	-	-
2	F2P	H	501	X	-	-	-
2	F2P	I	501	X	-	-	-
2	F2P	J	501	X	-	-	-
2	F2P	K	501	X	-	-	-
2	F2P	L	501	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	F2P	M	501	X	-	-	-
2	F2P	N	501	X	-	-	-
2	F2P	O	501	X	-	X	-
2	F2P	P	501	X	-	X	-
2	F2P	Q	501	X	-	-	-
2	F2P	R	501	X	-	-	-
2	F2P	S	501	X	-	-	-
2	F2P	T	501	X	-	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 41023 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative aldolase MJ0400.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	272	Total 2014	C 1263	N 354	O 385	S 12	0	0	0
1	B	268	Total 1998	C 1253	N 352	O 381	S 12	0	0	0
1	C	262	Total 1968	C 1237	N 347	O 373	S 11	0	0	0
1	D	267	Total 2002	C 1256	N 352	O 382	S 12	0	0	0
1	E	268	Total 2004	C 1258	N 353	O 381	S 12	0	0	0
1	K	266	Total 1997	C 1253	N 354	O 379	S 11	0	0	0
1	L	269	Total 2008	C 1260	N 354	O 382	S 12	0	0	0
1	M	265	Total 1985	C 1247	N 350	O 376	S 12	0	0	0
1	N	265	Total 1985	C 1247	N 348	O 378	S 12	0	0	0
1	O	266	Total 2000	C 1255	N 354	O 379	S 12	0	0	0
1	F	263	Total 1983	C 1245	N 351	O 376	S 11	0	0	0
1	G	268	Total 2016	C 1264	N 357	O 383	S 12	0	0	0
1	H	267	Total 1998	C 1254	N 352	O 380	S 12	0	0	0
1	I	265	Total 1996	C 1252	N 353	O 380	S 11	0	0	0
1	J	269	Total 2021	C 1267	N 358	O 384	S 12	0	0	0
1	P	268	Total 2003	C 1256	N 353	O 383	S 11	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	272	Total	C	N	O	S	0	0	0
			2035	1275	361	387	12			
1	R	269	Total	C	N	O	S	0	0	0
			2021	1267	360	382	12			
1	S	264	Total	C	N	O	S	0	0	0
			1988	1249	353	375	11			
1	T	267	Total	C	N	O	S	0	0	0
			2003	1257	353	382	11			

- Molecule 2 is 1,6-DI-O-PHOSPHONO-D-ALLITOL (three-letter code: F2P) (formula: $C_6H_{16}O_{12}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			19	6	11	2		
2	B	1	Total	C	O	P	0	0
			19	6	11	2		
2	C	1	Total	C	O	P	0	0
			19	6	11	2		
2	D	1	Total	C	O	P	0	0
			19	6	11	2		
2	E	1	Total	C	O	P	0	0
			19	6	11	2		
2	K	1	Total	C	O	P	0	0
			19	6	11	2		
2	L	1	Total	C	O	P	0	0
			19	6	11	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	M	1	Total	C	O	P	0	0
			19	6	11	2		
2	N	1	Total	C	O	P	0	0
			19	6	11	2		
2	O	1	Total	C	O	P	0	0
			19	6	11	2		
2	F	1	Total	C	O	P	0	0
			19	6	11	2		
2	G	1	Total	C	O	P	0	0
			19	6	11	2		
2	H	1	Total	C	O	P	0	0
			19	6	11	2		
2	I	1	Total	C	O	P	0	0
			19	6	11	2		
2	J	1	Total	C	O	P	0	0
			19	6	11	2		
2	P	1	Total	C	O	P	0	0
			19	6	11	2		
2	Q	1	Total	C	O	P	0	0
			19	6	11	2		
2	R	1	Total	C	O	P	0	0
			19	6	11	2		
2	S	1	Total	C	O	P	0	0
			19	6	11	2		
2	T	1	Total	C	O	P	0	0
			19	6	11	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	33	Total	O	0	0
			33	33		
3	B	34	Total	O	0	0
			34	34		
3	C	35	Total	O	0	0
			35	35		
3	D	27	Total	O	0	0
			27	27		
3	E	31	Total	O	0	0
			31	31		
3	K	19	Total	O	0	0
			19	19		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	25	Total 25	O 25	0	0
3	M	22	Total 22	O 22	0	0
3	N	23	Total 23	O 23	0	0
3	O	36	Total 36	O 36	0	0
3	F	33	Total 33	O 33	0	0
3	G	47	Total 47	O 47	0	0
3	H	38	Total 38	O 38	0	0
3	I	30	Total 30	O 30	0	0
3	J	28	Total 28	O 28	0	0
3	P	35	Total 35	O 35	0	0
3	Q	29	Total 29	O 29	0	0
3	R	39	Total 39	O 39	0	0
3	S	18	Total 18	O 18	0	0
3	T	36	Total 36	O 36	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400





- Molecule 1: Putative aldolase MJ0400








- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400



- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400









• Molecule 1: Putative aldolase MJ0400



• Molecule 1: Putative aldolase MJ0400




• Molecule 1: Putative aldolase MJ0400





• Molecule 1: Putative aldolase MJ0400







- Molecule 1: Putative aldolase MJ0400





- Molecule 1: Putative aldolase MJ0400




LYS

- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400









- Molecule 1: Putative aldolase MJ0400



- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400

Chain S:  71% 22%



- Molecule 1: Putative aldolase MJ0400

Chain T:  69% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	94.45Å 102.77Å 156.41Å 89.40° 85.83° 82.05°	Depositor
Resolution (Å)	46.31 – 2.60 46.31 – 2.60	Depositor EDS
% Data completeness (in resolution range)	89.8 (46.31-2.60) 89.5 (46.31-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.37Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.204 , 0.244 0.196 , 0.235	Depositor DCC
R_{free} test set	8568 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtrriage
Anisotropy	0.670	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	41023	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: F2P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2045	0.65	1/2770 (0.0%)
1	B	0.40	0/2028	0.63	0/2747
1	C	0.40	0/1998	0.63	0/2706
1	D	0.40	0/2032	0.63	0/2751
1	E	0.39	0/2034	0.64	0/2754
1	F	0.42	1/2013 (0.0%)	0.63	0/2725
1	G	0.42	0/2046	1.07	4/2767 (0.1%)
1	H	0.39	0/2028	0.63	0/2745
1	I	0.39	0/2026	0.62	0/2743
1	J	0.38	0/2051	0.64	0/2774
1	K	0.37	0/2027	0.62	0/2744
1	L	0.38	0/2038	0.63	0/2759
1	M	0.38	0/2015	0.62	0/2728
1	N	0.38	0/2014	0.62	0/2727
1	O	0.39	0/2030	0.65	1/2747 (0.0%)
1	P	0.40	1/2033 (0.0%)	0.66	2/2753 (0.1%)
1	Q	0.38	0/2066	0.63	0/2796
1	R	0.38	0/2051	0.62	0/2775
1	S	0.37	0/2018	0.62	0/2730
1	T	0.38	0/2033	0.64	0/2752
All	All	0.39	2/40626 (0.0%)	0.66	8/54993 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	P	183	VAL	C-N	5.92	1.47	1.34
1	F	228	GLU	CB-CG	-5.33	1.42	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	251	ARG	NE-CZ-NH1	-31.26	104.67	120.30
1	G	251	ARG	NE-CZ-NH2	28.28	134.44	120.30
1	G	251	ARG	CD-NE-CZ	14.89	144.44	123.60
1	G	251	ARG	CG-CD-NE	-10.04	90.70	111.80
1	P	183	VAL	C-N-CA	-9.70	97.45	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2014	0	1996	50	0
1	B	1998	0	1987	61	0
1	C	1968	0	1971	59	0
1	D	2002	0	1999	52	0
1	E	2004	0	1998	51	0
1	F	1983	0	1989	58	0
1	G	2016	0	2023	53	0
1	H	1998	0	1998	58	0
1	I	1996	0	1994	51	0
1	J	2021	0	2026	58	0
1	K	1997	0	1995	59	0
1	L	2008	0	2002	61	0
1	M	1985	0	1989	43	0
1	N	1985	0	1987	71	0
1	O	2000	0	2006	58	0
1	P	2003	0	1992	88	0
1	Q	2035	0	2033	54	0
1	R	2021	0	2023	56	0
1	S	1988	0	2000	76	0
1	T	2003	0	1999	71	0
2	A	19	0	10	0	0
2	B	19	0	10	2	0
2	C	19	0	10	2	0
2	D	19	0	10	5	0
2	E	19	0	10	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	19	0	10	4	0
2	G	19	0	10	2	0
2	H	19	0	10	5	0
2	I	19	0	10	0	0
2	J	19	0	10	5	0
2	K	19	0	10	4	0
2	L	19	0	10	1	0
2	M	19	0	10	2	0
2	N	19	0	10	2	0
2	O	19	0	10	13	0
2	P	19	0	10	8	0
2	Q	19	0	10	3	0
2	R	19	0	10	0	0
2	S	19	0	10	3	0
2	T	19	0	10	3	0
3	A	33	0	0	6	0
3	B	34	0	0	1	0
3	C	35	0	0	4	0
3	D	27	0	0	3	0
3	E	31	0	0	3	0
3	F	33	0	0	5	0
3	G	47	0	0	4	0
3	H	38	0	0	2	0
3	I	30	0	0	0	0
3	J	28	0	0	3	0
3	K	19	0	0	0	0
3	L	25	0	0	1	0
3	M	22	0	0	1	0
3	N	23	0	0	1	0
3	O	36	0	0	6	0
3	P	35	0	0	3	0
3	Q	29	0	0	4	0
3	R	39	0	0	2	0
3	S	18	0	0	2	0
3	T	36	0	0	2	0
All	All	41023	0	40207	1108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 1108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:N:164:ASP:HB2	1:P:214:ASN:HD21	1.02	1.14
1:D:184:LYS:HE3	2:D:501:F2P:O4	1.40	1.11
2:D:501:F2P:O13	2:D:501:F2P:H2	1.45	1.11
1:S:21:ASN:ND2	1:S:24:SER:H	1.55	1.05
1:D:140:THR:HG21	3:D:522:HOH:O	1.56	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/273 (99%)	256 (95%)	12 (4%)	2 (1%)	19	38
1	B	264/273 (97%)	249 (94%)	15 (6%)	0	100	100
1	C	258/273 (94%)	246 (95%)	11 (4%)	1 (0%)	30	52
1	D	263/273 (96%)	250 (95%)	11 (4%)	2 (1%)	16	34
1	E	264/273 (97%)	249 (94%)	14 (5%)	1 (0%)	30	52
1	F	259/273 (95%)	247 (95%)	11 (4%)	1 (0%)	30	52
1	G	264/273 (97%)	253 (96%)	11 (4%)	0	100	100
1	H	263/273 (96%)	252 (96%)	11 (4%)	0	100	100
1	I	261/273 (96%)	250 (96%)	11 (4%)	0	100	100
1	J	265/273 (97%)	249 (94%)	15 (6%)	1 (0%)	30	52
1	K	262/273 (96%)	246 (94%)	16 (6%)	0	100	100
1	L	265/273 (97%)	249 (94%)	14 (5%)	2 (1%)	16	34
1	M	261/273 (96%)	247 (95%)	14 (5%)	0	100	100
1	N	261/273 (96%)	247 (95%)	11 (4%)	3 (1%)	12	26
1	O	262/273 (96%)	248 (95%)	13 (5%)	1 (0%)	30	52
1	P	264/273 (97%)	251 (95%)	11 (4%)	2 (1%)	16	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	270/273 (99%)	256 (95%)	12 (4%)	2 (1%)	19	38
1	R	265/273 (97%)	256 (97%)	9 (3%)	0	100	100
1	S	260/273 (95%)	245 (94%)	14 (5%)	1 (0%)	30	52
1	T	263/273 (96%)	247 (94%)	16 (6%)	0	100	100
All	All	5264/5460 (96%)	4993 (95%)	252 (5%)	19 (0%)	30	52

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	74	ARG
1	A	78	LYS
1	C	70	ARG
1	Q	74	ARG
1	L	72	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/220 (94%)	192 (92%)	16 (8%)	10	22
1	B	208/220 (94%)	190 (91%)	18 (9%)	8	17
1	C	207/220 (94%)	193 (93%)	14 (7%)	13	28
1	D	210/220 (96%)	198 (94%)	12 (6%)	17	37
1	E	209/220 (95%)	194 (93%)	15 (7%)	12	26
1	F	209/220 (95%)	194 (93%)	15 (7%)	12	26
1	G	212/220 (96%)	197 (93%)	15 (7%)	12	26
1	H	209/220 (95%)	193 (92%)	16 (8%)	10	22
1	I	210/220 (96%)	197 (94%)	13 (6%)	15	33
1	J	212/220 (96%)	196 (92%)	16 (8%)	11	24
1	K	209/220 (95%)	196 (94%)	13 (6%)	15	33
1	L	209/220 (95%)	193 (92%)	16 (8%)	10	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	208/220 (94%)	193 (93%)	15 (7%)	12	26
1	N	208/220 (94%)	192 (92%)	16 (8%)	10	22
1	O	210/220 (96%)	196 (93%)	14 (7%)	13	29
1	P	209/220 (95%)	194 (93%)	15 (7%)	12	26
1	Q	212/220 (96%)	198 (93%)	14 (7%)	14	30
1	R	211/220 (96%)	197 (93%)	14 (7%)	14	30
1	S	209/220 (95%)	195 (93%)	14 (7%)	13	29
1	T	210/220 (96%)	195 (93%)	15 (7%)	12	26
All	All	4189/4400 (95%)	3893 (93%)	296 (7%)	12	26

5 of 296 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	P	95	ASN
1	T	80	VAL
1	P	236	VAL
1	R	140	THR
1	L	170	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 84 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	9	ASN
1	R	9	ASN
1	J	65	HIS
1	P	160	GLN
1	R	161	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	F2P	J	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	K	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.09	3 (13%)
2	F2P	P	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	A	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.09	3 (13%)
2	F2P	S	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	L	501	1	18,18,19	1.26	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	Q	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.09	3 (13%)
2	F2P	I	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	G	501	1	18,18,19	1.22	1 (5%)	23,26,28	1.16	3 (13%)
2	F2P	H	501	1	18,18,19	1.26	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	D	501	1	18,18,19	1.07	2 (11%)	23,26,28	1.49	5 (21%)
2	F2P	T	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.09	3 (13%)
2	F2P	C	501	1	18,18,19	1.22	1 (5%)	23,26,28	1.16	3 (13%)
2	F2P	M	501	1	18,18,19	1.26	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	O	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	R	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	B	501	1	18,18,19	1.26	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	N	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.09	3 (13%)
2	F2P	F	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.10	3 (13%)
2	F2P	E	501	1	18,18,19	1.25	1 (5%)	23,26,28	1.09	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	F2P	J	501	1	2/2/5/6	9/21/21/24	-
2	F2P	K	501	1	2/2/5/6	5/21/21/24	-
2	F2P	P	501	1	2/2/5/6	7/21/21/24	-
2	F2P	A	501	1	2/2/5/6	4/21/21/24	-
2	F2P	S	501	1	2/2/5/6	4/21/21/24	-
2	F2P	L	501	1	2/2/5/6	5/21/21/24	-
2	F2P	Q	501	1	2/2/5/6	1/21/21/24	-
2	F2P	I	501	1	2/2/5/6	4/21/21/24	-
2	F2P	G	501	1	2/2/5/6	5/21/21/24	-
2	F2P	H	501	1	2/2/5/6	2/21/21/24	-
2	F2P	D	501	1	2/2/5/6	10/21/21/24	-
2	F2P	T	501	1	2/2/5/6	5/21/21/24	-
2	F2P	C	501	1	2/2/5/6	2/21/21/24	-
2	F2P	M	501	1	2/2/5/6	3/21/21/24	-
2	F2P	O	501	1	2/2/5/6	5/21/21/24	-
2	F2P	R	501	1	2/2/5/6	6/21/21/24	-
2	F2P	B	501	1	2/2/5/6	3/21/21/24	-
2	F2P	N	501	1	2/2/5/6	2/21/21/24	-
2	F2P	F	501	1	2/2/5/6	5/21/21/24	-
2	F2P	E	501	1	2/2/5/6	6/21/21/24	-

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	501	F2P	P6-O61	3.54	1.61	1.50
2	B	501	F2P	P6-O61	3.54	1.61	1.50
2	L	501	F2P	P6-O61	3.54	1.61	1.50
2	P	501	F2P	P6-O61	3.54	1.61	1.50
2	H	501	F2P	P6-O61	3.53	1.61	1.50

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	501	F2P	O6-P6-O61	4.08	117.46	106.44
2	C	501	F2P	C5-C4-C3	-3.59	108.28	113.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	501	F2P	C5-C4-C3	-3.54	108.36	113.97
2	D	501	F2P	C5-C4-C3	-2.99	109.22	113.97
2	O	501	F2P	C5-C4-C3	-2.84	109.47	113.97

5 of 40 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	501	F2P	C4
2	A	501	F2P	C5
2	B	501	F2P	C4
2	B	501	F2P	C5
2	C	501	F2P	C4

5 of 93 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	F2P	C2-C1-O1-P1
2	C	501	F2P	C4-C5-C6-O6
2	C	501	F2P	C6-O6-P6-O62
2	D	501	F2P	C1-O1-P1-O11
2	D	501	F2P	C1-O1-P1-O12

There are no ring outliers.

17 monomers are involved in 65 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	J	501	F2P	5	0
2	K	501	F2P	4	0
2	P	501	F2P	8	0
2	S	501	F2P	3	0
2	L	501	F2P	1	0
2	Q	501	F2P	3	0
2	G	501	F2P	2	0
2	H	501	F2P	5	0
2	D	501	F2P	5	0
2	T	501	F2P	3	0
2	C	501	F2P	2	0
2	M	501	F2P	2	0
2	O	501	F2P	13	0
2	B	501	F2P	2	0
2	N	501	F2P	2	0
2	F	501	F2P	4	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	501	F2P	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/273 (99%)	-0.14	8 (2%) 54 48	19, 37, 69, 100	0
1	B	268/273 (98%)	-0.24	9 (3%) 48 42	20, 34, 69, 115	0
1	C	262/273 (95%)	-0.17	3 (1%) 77 74	16, 38, 68, 107	0
1	D	267/273 (97%)	-0.20	5 (1%) 66 61	18, 36, 64, 96	0
1	E	268/273 (98%)	-0.25	4 (1%) 71 67	19, 36, 70, 95	0
1	F	263/273 (96%)	-0.20	2 (0%) 82 79	19, 36, 64, 85	0
1	G	268/273 (98%)	-0.19	6 (2%) 62 57	18, 35, 71, 96	0
1	H	267/273 (97%)	-0.25	6 (2%) 62 57	16, 33, 65, 109	0
1	I	265/273 (97%)	-0.27	4 (1%) 71 67	16, 37, 69, 84	0
1	J	269/273 (98%)	-0.05	5 (1%) 66 61	18, 40, 76, 96	0
1	K	266/273 (97%)	0.01	5 (1%) 66 61	23, 43, 75, 117	0
1	L	269/273 (98%)	-0.24	3 (1%) 77 74	20, 36, 71, 100	0
1	M	265/273 (97%)	-0.11	5 (1%) 66 61	20, 39, 72, 96	0
1	N	265/273 (97%)	-0.20	5 (1%) 66 61	17, 39, 66, 88	0
1	O	266/273 (97%)	-0.11	5 (1%) 66 61	15, 39, 68, 98	0
1	P	268/273 (98%)	0.05	7 (2%) 57 51	21, 41, 70, 109	0
1	Q	272/273 (99%)	-0.14	4 (1%) 71 67	18, 36, 76, 118	0
1	R	269/273 (98%)	-0.10	5 (1%) 66 61	19, 39, 80, 101	0
1	S	264/273 (96%)	0.02	4 (1%) 71 67	23, 43, 72, 91	0
1	T	267/273 (97%)	-0.03	7 (2%) 57 51	19, 40, 69, 106	0
All	All	5340/5460 (97%)	-0.14	102 (1%) 66 61	15, 38, 72, 118	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	73	HIS	5.7
1	A	75	GLY	5.0
1	A	76	TYR	4.9
1	A	77	GLY	4.6
1	T	73	HIS	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	F2P	H	501	19/20	0.74	0.24	53,63,82,82	0
2	F2P	N	501	19/20	0.77	0.19	48,64,88,88	0
2	F2P	S	501	19/20	0.77	0.20	62,75,96,97	0
2	F2P	C	501	19/20	0.78	0.18	55,63,87,87	0
2	F2P	K	501	19/20	0.79	0.17	56,65,89,90	0
2	F2P	J	501	19/20	0.80	0.18	50,70,91,91	0
2	F2P	R	501	19/20	0.80	0.17	59,64,82,82	0
2	F2P	D	501	19/20	0.80	0.18	48,65,88,88	0
2	F2P	O	501	19/20	0.81	0.18	62,72,90,90	0
2	F2P	P	501	19/20	0.81	0.16	50,68,81,81	0
2	F2P	B	501	19/20	0.82	0.18	47,65,78,79	0
2	F2P	A	501	19/20	0.83	0.17	53,63,87,88	0
2	F2P	M	501	19/20	0.85	0.17	56,72,95,95	0
2	F2P	F	501	19/20	0.85	0.14	55,65,82,83	0
2	F2P	L	501	19/20	0.85	0.14	51,62,83,83	0
2	F2P	I	501	19/20	0.85	0.16	48,63,82,83	0
2	F2P	T	501	19/20	0.85	0.15	44,57,79,79	0
2	F2P	Q	501	19/20	0.86	0.17	28,50,87,88	0
2	F2P	E	501	19/20	0.87	0.16	43,57,79,80	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	F2P	G	501	19/20	0.87	0.16	44,58,88,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around F2P N 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around F2P S 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around F2P K 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P J 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P R 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P O 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P P 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around F2P M 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P F 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P L 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2P I 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around F2P T 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around F2P Q 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around F2P E 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.