

wwPDB X-ray Structure Validation Summary Report (i)

Feb 15, 2024 – 03:48 PM EST

PDB ID : 3QKJ

Title The PWWP domain of human DNA (CYTOSINE-5-)-METHYLTRANSFER

ASE 3 BETA in complex with a bis-tris molecule

Authors: Zeng, H.; Amaya, M.F.; Mackenzie, F.; Weigelt, J.; Sundstrom, M.; Arrow-

smith, C.H.; Edwards, A.M.; Botchkarev, A.; Min, J.; Wu, H.; Structural

Genomics Consortium (SGC)

Deposited on 2011-02-01

Resolution 2.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

> The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as 541 be (2020)Mogul

Xtriage (Phenix) 1.13

EDS 2.36

Percentile statistics 20191225.v01 (using entries in the PDB archive December 25th 2019)

> Refmac 5.8.0158

CCP4 7.0.044 (Gargrove) Ideal geometry (proteins) Engh & Huber (2001) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	151	82%	•	14%
1	В	151	83%	5%	13%
1	С	151	84%		13%
1	D	151	81%	5%	13%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DNA cytosine-5 methyltransferase 3 beta isoform 6 variant.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	130	Total	С	N	О	S	0	1	0
1	A	130	1027	672	178	172	5	0	1	U
1	В	132	Total	С	N	О	S	0	1	0
1	Ъ	152	1043	682	181	175	5	0	1	U
1	С	131	Total	С	N	О	S	0	0	0
1		131	1026	670	172	179	5	0	0	U
1	1 D	131	Total	С	N	О	S	0	0	0
1	D	131	1029	674	173	177	5	U	U	U

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	GLY	-	expression tag	UNP Q59H79
В	205	GLY	-	expression tag	UNP Q59H79
С	205	GLY	-	expression tag	UNP Q59H79
D	205	GLY	-	expression tag	UNP Q59H79


• Molecule 2 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	А	1	Total C N O	0	0	
	11	1	14 8 1 5	Ů	Ŭ .	
2	В	1	Total C N O	0	0	
	Ъ	1	14 8 1 5	0	U	
2	C	1	Total C N O	0	0	
	C	1	14 8 1 5	0	U	
2	D	1	Total C N O	0	0	
	D		14 8 1 5	0		

 \bullet Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$





Mol	Chain	Residues	Atoms	8	ZeroOcc	AltConf	
3	A	1	Total O	S	0	0	
	11	1	5 4	1	O	U	
3	A	1	Total O	\mathbf{S}	0	0	
	71	1	5 4	1	O	U	
3	В	1	Total O	S	0	0	
	D	1	5 4	1	O		
3	В	1	Total O	S	0	0	
	D	1	5 4	1	O	U	
3	D	1	Total O	S	0	0	
	D	1	5 4	1	O	Ŭ	
3	D	1	Total O	S	0	0	
			5 4	1		U	

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	75	Total O 75 75	0	0
4	В	76	Total O 76 76	0	0
4	С	57	Total O 57 57	0	0
4	D	55	Total O 55 55	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA cytosine-5 methyltransferase 3 beta isoform 6 variant





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32	Depositor
Cell constants	74.54Å 74.54Å 160.27Å	Domositon
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.04 - 2.04	Depositor
rtesolution (A)	28.71 - 2.04	EDS
% Data completeness	98.4 (34.04-2.04)	Depositor
(in resolution range)	98.5 (28.71-2.04)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.72 (at 2.04Å)	Xtriage
Refinement program	REFMAC 5.5.0072, BUSTER 2.8.0	Depositor
R, R_{free}	0.209 , 0.228	Depositor
	0.207 , 0.230	DCC
R_{free} test set	3163 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	48.1	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 47.6	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.35$	Xtriage
	0.012 for -h,-k,l	
Estimated twinning fraction	0.477 for h,-h-k,-l	Xtriage
	0.015 for -k,-h,-l	
F_o, F_c correlation	0.96	EDS
Total number of atoms	4474	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BTB, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	\mathbf{angles}
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.50	0/1062	0.60	0/1432
1	В	0.50	0/1075	0.60	0/1449
1	С	0.51	0/1057	0.61	0/1427
1	D	0.51	0/1061	0.60	0/1432
All	All	0.50	0/4255	0.61	0/5740

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1027	0	989	2	0
1	В	1043	0	998	2	0
1	С	1026	0	966	1	0
1	D	1029	0	980	5	0
2	A	14	0	19	3	0
2	В	14	0	19	3	0
2	С	14	0	19	2	0
2	D	14	0	19	3	0
3	A	10	0	0	0	0

Continued on next page...



I 'omtamalod	trom	mmonia	maaa
Continued	11 0116	DICUIUUS	Daue
	.,	10	1

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	10	0	0	0	0
3	D	10	0	0	0	0
4	A	75	0	0	0	0
4	В	76	0	0	0	0
4	С	57	0	0	0	0
4	D	55	0	0	0	0
All	All	4474	0	4009	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 19 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$	
2:B:356:BTB:H81	2:B:356:BTB:H62	1.55	0.85	
2:A:356:BTB:H61	2:A:356:BTB:H82	1.62	0.81	
2:A:356:BTB:H82	2:A:356:BTB:C6	2.27	0.64	
2:B:356:BTB:H62	2:B:356:BTB:C8	2.31	0.57	
1:D:239:TRP:CD2	2:D:356:BTB:H12	2.50	0.47	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles		
1	A	127/151 (84%)	125 (98%)	1 (1%)	1 (1%)	19	10	
1	В	129/151 (85%)	127 (98%)	1 (1%)	1 (1%)	19	10	
1	С	127/151 (84%)	124 (98%)	2 (2%)	1 (1%)	19	10	
1	D	127/151 (84%)	124 (98%)	2 (2%)	1 (1%)	19	10	
All	All	510/604 (84%)	500 (98%)	6 (1%)	4 (1%)	19	10	



All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	223	PHE
1	С	223	PHE
1	A	223	PHE
1	D	223	PHE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	98/121 (81%)	97 (99%)	1 (1%)	76 75		
1	В	98/121 (81%)	96 (98%)	2 (2%)	55 50		
1	С	97/121 (80%)	96 (99%)	1 (1%)	76 75		
1	D	98/121 (81%)	98 (100%)	0	100 100		
All	All	391/484 (81%)	387 (99%)	4 (1%)	76 75		

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256	SER
1	В	218	GLN
1	В	256	SER
1	С	329	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Вс	Bond lengths			Bond angles		
WIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	BTB	С	356	-	13,13,13	0.52	0	7,16,16	0.60	0	
3	SO4	В	5	-	4,4,4	0.26	0	6,6,6	0.11	0	
2	BTB	D	356	_	13,13,13	0.53	0	7,16,16	0.68	0	
3	SO4	A	3	_	4,4,4	0.33	0	6,6,6	0.15	0	
3	SO4	A	4	-	4,4,4	0.33	0	6,6,6	0.16	0	
3	SO4	D	2	_	4,4,4	0.22	0	6,6,6	0.19	0	
3	SO4	В	6	-	4,4,4	0.09	0	6,6,6	0.08	0	
2	BTB	A	356	_	13,13,13	0.47	0	7,16,16	0.25	0	
3	SO4	D	1	-	4,4,4	0.24	0	6,6,6	0.22	0	
2	BTB	В	356	_	13,13,13	0.51	0	7,16,16	0.26	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ВТВ	В	356	-	-	2/21/21/21	-
2	ВТВ	D	356	-	-	5/21/21/21	-
2	ВТВ	С	356	-	-	5/21/21/21	-
2	ВТВ	A	356	-	-	2/21/21/21	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	356	BTB	C3-C2-N-C5
2	В	356	BTB	C3-C2-N-C7
2	С	356	BTB	C1-C2-N-C7
2	С	356	BTB	C3-C2-N-C7
2	D	356	BTB	C1-C2-N-C7

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	356	BTB	2	0
2	D	356	BTB	3	0
2	A	356	BTB	3	0
2	В	356	BTB	3	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	A	130/151 (86%)	-0.53	0 100 100	38, 52, 78, 91	0
1	В	132/151 (87%)	-0.47	0 100 100	39, 52, 79, 119	0
1	С	131/151 (86%)	-0.41	0 100 100	41, 62, 92, 124	0
1	D	131/151 (86%)	-0.45	0 100 100	42, 62, 92, 114	0
All	All	524/604 (86%)	-0.46	0 100 100	38, 56, 88, 124	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	SO4	D	1	5/5	0.88	0.16	110,114,115,116	0
2	BTB	С	356	14/14	0.93	0.14	50,56,106,128	0
3	SO4	A	4	5/5	0.94	0.15	106,110,111,112	0
2	BTB	D	356	14/14	0.95	0.12	50,69,91,119	0

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	SO4	A	3	5/5	0.95	0.19	87,90,92,92	0
2	BTB	В	356	14/14	0.95	0.12	40,53,79,83	0
2	BTB	A	356	14/14	0.95	0.10	44,55,81,88	0
3	SO4	В	6	5/5	0.96	0.21	136,140,141,141	0
3	SO4	В	5	5/5	0.96	0.10	98,102,103,104	0
3	SO4	D	2	5/5	0.96	0.09	94,99,99,99	0

6.5 Other polymers (i)

There are no such residues in this entry.

