



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 06:09 PM BST

PDB ID : 6QO0  
Title : I47W mutated sulfur oxygenase reductase from *Acidianus ambivaens*  
Authors : Frazao, C.; Klezin, A.; Poell, U.  
Deposited on : 2019-02-12  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	
1	B	318	
1	C	318	
1	D	318	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfur oxygenase/reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	307	2512	1624	413	454	21	0	13	0
1	B	307	2510	1623	413	452	22	0	12	0
1	C	307	2513	1624	413	454	22	0	13	0
1	D	307	2509	1622	413	452	22	0	12	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	TRP	ILE	engineered mutation	UNP P29082
A	309	ASN	-	expression tag	UNP P29082
A	310	ALA	-	expression tag	UNP P29082
A	311	TRP	-	expression tag	UNP P29082
A	312	ARG	-	expression tag	UNP P29082
A	313	HIS	-	expression tag	UNP P29082
A	314	PRO	-	expression tag	UNP P29082
A	315	GLN	-	expression tag	UNP P29082
A	316	PHE	-	expression tag	UNP P29082
A	317	GLY	-	expression tag	UNP P29082
A	318	GLY	-	expression tag	UNP P29082
B	47	TRP	ILE	engineered mutation	UNP P29082
B	309	ASN	-	expression tag	UNP P29082
B	310	ALA	-	expression tag	UNP P29082
B	311	TRP	-	expression tag	UNP P29082
B	312	ARG	-	expression tag	UNP P29082
B	313	HIS	-	expression tag	UNP P29082
B	314	PRO	-	expression tag	UNP P29082
B	315	GLN	-	expression tag	UNP P29082
B	316	PHE	-	expression tag	UNP P29082
B	317	GLY	-	expression tag	UNP P29082

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	318	GLY	-	expression tag	UNP P29082
C	47	TRP	ILE	engineered mutation	UNP P29082
C	309	ASN	-	expression tag	UNP P29082
C	310	ALA	-	expression tag	UNP P29082
C	311	TRP	-	expression tag	UNP P29082
C	312	ARG	-	expression tag	UNP P29082
C	313	HIS	-	expression tag	UNP P29082
C	314	PRO	-	expression tag	UNP P29082
C	315	GLN	-	expression tag	UNP P29082
C	316	PHE	-	expression tag	UNP P29082
C	317	GLY	-	expression tag	UNP P29082
C	318	GLY	-	expression tag	UNP P29082
D	47	TRP	ILE	engineered mutation	UNP P29082
D	309	ASN	-	expression tag	UNP P29082
D	310	ALA	-	expression tag	UNP P29082
D	311	TRP	-	expression tag	UNP P29082
D	312	ARG	-	expression tag	UNP P29082
D	313	HIS	-	expression tag	UNP P29082
D	314	PRO	-	expression tag	UNP P29082
D	315	GLN	-	expression tag	UNP P29082
D	316	PHE	-	expression tag	UNP P29082
D	317	GLY	-	expression tag	UNP P29082
D	318	GLY	-	expression tag	UNP P29082

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Fe 1 1	0	0
2	A	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	232	Total O 232 232	0	0

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	259	Total 259	O 259	0	0
3	C	229	Total 229	O 229	0	0
3	D	271	Total 271	O 271	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sulfur oxygenase/reductase




- Molecule 1: Sulfur oxygenase/reductase



- Molecule 1: Sulfur oxygenase/reductase



- Molecule 1: Sulfur oxygenase/reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	158.16Å 158.16Å 227.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	113.80 – 1.65 136.97 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (113.80-1.65) 99.7 (136.97-1.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.15	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 1.65Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.168 , 0.191 0.167 , 0.191	Depositor DCC
$R_{free}$ test set	3200 reflections (1.60%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtrriage
Anisotropy	0.503	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11039	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2651	0.52	0/3593
1	B	0.36	0/2644	0.55	0/3583
1	C	0.33	0/2652	0.52	0/3594
1	D	0.37	0/2644	0.56	0/3583
All	All	0.35	0/10591	0.54	0/14353

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2512	0	2445	17	0
1	B	2510	0	2448	19	0
1	C	2513	0	2448	13	0
1	D	2509	0	2448	15	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	232	0	0	3	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	259	0	0	2	0
3	C	229	0	0	1	0
3	D	271	0	0	1	0
All	All	11039	0	9789	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:243[A]:ASN:ND2	3:D:501:HOH:O	2.06	0.87
1:D:45:ASN:HB2	1:D:297[B]:MET:HG2	1.57	0.86
1:A:45:ASN:HB2	1:A:297[B]:MET:HG2	1.60	0.83
1:B:45:ASN:HB2	1:B:297[B]:MET:HG2	1.60	0.81
1:C:45:ASN:HB2	1:C:297[B]:MET:HG2	1.66	0.77
1:A:132:ASP:OD1	3:A:501:HOH:O	2.03	0.76
1:C:182:ARG:NH1	3:C:503:HOH:O	2.27	0.66
1:A:99:ARG:NH2	1:A:228:GLU:OE2	2.31	0.61
1:B:308:GLU:O	3:B:501:HOH:O	2.16	0.60
1:D:307:ASN:O	1:D:308:GLU:HG2	2.04	0.57
1:A:22:MET:SD	1:A:104:CYS:HA	2.45	0.56
1:D:99:ARG:NH2	1:D:228:GLU:OE2	2.38	0.56
1:B:99:ARG:NH2	1:B:228:GLU:OE2	2.34	0.56
1:C:223:ASN:ND2	1:C:227:LEU:O	2.29	0.54
1:A:142:ALA:HB2	1:B:146:PRO:HG2	1.91	0.52
1:D:105:ALA:HA	1:D:108:MET:HG3	1.91	0.51
1:A:36[B]:ARG:NH1	3:A:504:HOH:O	2.33	0.50
1:B:223:ASN:ND2	1:B:227:LEU:O	2.28	0.50
1:B:156:TYR:HA	1:B:296:MET:O	2.13	0.49
1:B:29:LYS:HA	1:B:32:MET:HE2	1.95	0.48
1:A:262:PHE:HZ	1:C:297[A]:MET:HG2	1.79	0.48
1:C:22:MET:SD	1:C:104:CYS:HA	2.54	0.48
1:C:105:ALA:HA	1:C:108:MET:HG3	1.94	0.47
1:C:262:PHE:HZ	1:D:297[A]:MET:HG2	1.80	0.47
1:A:25:SER:HB2	1:B:56:ASN:HB3	1.96	0.46
1:B:105:ALA:HA	1:B:108:MET:HG3	1.97	0.45
1:A:105:ALA:HA	1:A:108:MET:HG3	1.98	0.45
1:A:279[B]:GLU:OE2	3:A:502:HOH:O	2.21	0.45
1:C:156:TYR:HA	1:C:296:MET:O	2.17	0.44
1:B:161:VAL:HB	1:B:293:LEU:HB2	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:MET:SD	1:B:104:CYS:HA	2.58	0.43
1:B:182:ARG:NH1	3:B:513:HOH:O	2.51	0.43
1:B:297[A]:MET:HG2	1:D:262:PHE:HZ	1.84	0.43
1:A:161:VAL:HB	1:A:293:LEU:HB2	2.01	0.43
1:A:156:TYR:HA	1:A:296:MET:O	2.19	0.43
1:C:99:ARG:NH2	1:C:228:GLU:OE2	2.48	0.43
1:A:95:SER:HA	1:A:228:GLU:HG2	2.01	0.42
1:C:51:ILE:HB	1:C:61:ALA:HB1	2.02	0.42
1:A:223:ASN:ND2	1:A:227:LEU:O	2.32	0.42
1:A:271:PRO:O	1:A:275:GLN:HG2	2.21	0.41
1:D:156:TYR:HA	1:D:296:MET:O	2.20	0.41
1:C:271:PRO:O	1:C:275:GLN:HG2	2.20	0.41
1:D:296:MET:HB3	1:D:297[B]:MET:SD	2.61	0.41
1:A:23:PHE:CD2	1:A:47:TRP:CH2	3.09	0.41
1:B:271:PRO:O	1:B:275:GLN:HG2	2.20	0.41
1:B:23:PHE:CD2	1:B:47:TRP:CH2	3.09	0.41
1:D:22:MET:HG3	1:D:23:PHE:N	2.35	0.41
1:D:161:VAL:HB	1:D:293:LEU:HB2	2.02	0.40
1:B:23:PHE:CD2	1:B:47:TRP:HH2	2.39	0.40
1:B:297[A]:MET:HG2	1:D:262:PHE:CZ	2.57	0.40
1:B:25:SER:HB2	1:D:56:ASN:HB3	2.03	0.40
1:D:51:ILE:HB	1:D:61:ALA:HB1	2.04	0.40
1:A:146:PRO:HG2	1:C:142:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/318 (100%)	312 (98%)	6 (2%)	0	100	100
1	B	317/318 (100%)	310 (98%)	7 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	318/318 (100%)	311 (98%)	7 (2%)	0	100	100
1	D	317/318 (100%)	310 (98%)	7 (2%)	0	100	100
All	All	1270/1272 (100%)	1243 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/273 (102%)	277 (100%)	0	100	100
1	B	277/273 (102%)	276 (100%)	1 (0%)	91	85
1	C	278/273 (102%)	278 (100%)	0	100	100
1	D	277/273 (102%)	275 (99%)	2 (1%)	84	73
All	All	1109/1092 (102%)	1106 (100%)	3 (0%)	92	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	22	MET
1	D	22	MET
1	D	147	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/318 (96%)	-0.60	3 (0%) 82 85	23, 30, 46, 60	0
1	B	307/318 (96%)	-0.53	3 (0%) 82 85	21, 26, 44, 63	0
1	C	307/318 (96%)	-0.57	1 (0%) 94 94	22, 29, 46, 67	0
1	D	307/318 (96%)	-0.58	2 (0%) 87 89	20, 26, 46, 65	0
All	All	1228/1272 (96%)	-0.57	9 (0%) 87 89	20, 28, 46, 67	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	23	PHE	4.4
1	A	23	PHE	3.6
1	D	23	PHE	3.3
1	B	17	PRO	2.8
1	A	102	TYR	2.5
1	D	47	TRP	2.5
1	A	47	TRP	2.3
1	B	47	TRP	2.3
1	C	47	TRP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FE	B	401	1/1	0.99	0.05	25,25,25,25	1
2	FE	A	401	1/1	0.99	0.03	28,28,28,28	1
2	FE	D	401	1/1	0.99	0.05	25,25,25,25	1
2	FE	C	401	1/1	0.99	0.05	32,32,32,32	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.