

wwPDB X-ray Structure Validation Summary Report (i)

May 16, 2020 – 12:50 pm BST

PDB ID : 2QRY

Title: Periplasmic thiamin binding protein

Authors : Ealick, S.E.; Soriano, E.V.

Deposited on : 2007-07-30

Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	330	68%	23%	• 7%				
1	В	330	70%	22%	• 6%				
1	С	330	70%	24%					
1	D	330	68%	25%	• 6%				



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 10279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Thiamine-binding periplasmic protein.

Mol	Chain	Residues		Atoms						AltConf	Trace
1	Λ	308	Total	С	N	О	S	Se	0	0	0
1	A 308	300	2395	1546	393	450	2	4	U	U	
1	В	309	Total	С	N	О	S	Se	0	0	0
1	Ъ	309	2412	1554	398	454	2	4	0		
1	С	318	Total	С	N	О	S	Se	0	0	0
1		310	2470	1590	411	462	2	5	U	U	
1	D	210	Total	С	N	О	S	Se	0	0	0
1		310	2414	1556	397	454	2	5	U	U	

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MSE	-	EXPRESSION TAG	UNP P31550
A	-1	GLY	_	EXPRESSION TAG	UNP P31550
A	0	SER	-	EXPRESSION TAG	UNP P31550
A	1	SER	-	EXPRESSION TAG	UNP P31550
A	2	HIS	-	EXPRESSION TAG	UNP P31550
A	3	HIS	-	EXPRESSION TAG	UNP P31550
A	4	HIS	-	EXPRESSION TAG	UNP P31550
A	5	HIS	-	EXPRESSION TAG	UNP P31550
A	6	HIS	=	EXPRESSION TAG	UNP P31550
A	7	HIS	-	EXPRESSION TAG	UNP P31550
A	8	SER	=	EXPRESSION TAG	UNP P31550
A	9	SER	-	EXPRESSION TAG	UNP P31550
A	10	GLY	-	EXPRESSION TAG	UNP P31550
A	11	LEU	-	EXPRESSION TAG	UNP P31550
A	12	VAL	-	EXPRESSION TAG	UNP P31550
A	13	PRO	-	EXPRESSION TAG	UNP P31550
A	14	ARG	-	EXPRESSION TAG	UNP P31550
A	15	GLY	-	EXPRESSION TAG	UNP P31550
A	16	SER	-	EXPRESSION TAG	UNP P31550
A	17	HIS	-	EXPRESSION TAG	UNP P31550
A	18	MSE	-	EXPRESSION TAG	UNP P31550

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	-2	MSE	-	EXPRESSION TAG	UNP P31550
В	-1	GLY	-	EXPRESSION TAG	UNP P31550
В	0	SER	-	EXPRESSION TAG	UNP P31550
В	1	SER	-	EXPRESSION TAG	UNP P31550
В	2	HIS	-	EXPRESSION TAG	UNP P31550
В	3	HIS	-	EXPRESSION TAG	UNP P31550
В	4	HIS	-	EXPRESSION TAG	UNP P31550
В	5	HIS	_	EXPRESSION TAG	UNP P31550
В	6	HIS	-	EXPRESSION TAG	UNP P31550
В	7	HIS	-	EXPRESSION TAG	UNP P31550
В	8	SER	-	EXPRESSION TAG	UNP P31550
В	9	SER	_	EXPRESSION TAG	UNP P31550
В	10	GLY	-	EXPRESSION TAG	UNP P31550
В	11	LEU	-	EXPRESSION TAG	UNP P31550
В	12	VAL	-	EXPRESSION TAG	UNP P31550
В	13	PRO	_	EXPRESSION TAG	UNP P31550
В	14	ARG	-	EXPRESSION TAG	UNP P31550
В	15	GLY	_	EXPRESSION TAG	UNP P31550
В	16	SER	_	EXPRESSION TAG	UNP P31550
В	17	HIS	-	EXPRESSION TAG	UNP P31550
В	18	MSE	_	EXPRESSION TAG	UNP P31550
С	-2	MSE	-	EXPRESSION TAG	UNP P31550
С	-1	GLY	-	EXPRESSION TAG	UNP P31550
С	0	SER	-	EXPRESSION TAG	UNP P31550
С	1	SER	-	EXPRESSION TAG	UNP P31550
С	2	HIS	-	EXPRESSION TAG	UNP P31550
С	3	HIS	-	EXPRESSION TAG	UNP P31550
С	4	HIS	_	EXPRESSION TAG	UNP P31550
С	5	HIS	-	EXPRESSION TAG	UNP P31550
С	6	HIS	_	EXPRESSION TAG	UNP P31550
С	7	HIS	-	EXPRESSION TAG	UNP P31550
С	8	SER	-	EXPRESSION TAG	UNP P31550
С	9	SER	-	EXPRESSION TAG	UNP P31550
С	10	GLY	-	EXPRESSION TAG	UNP P31550
С	11	LEU	-	EXPRESSION TAG	UNP P31550
С	12	VAL	-	EXPRESSION TAG	UNP P31550
С	13	PRO	-	EXPRESSION TAG	UNP P31550
С	14	ARG	-	EXPRESSION TAG	UNP P31550
С	15	GLY	-	EXPRESSION TAG	UNP P31550
С	16	SER	-	EXPRESSION TAG	UNP P31550
С	17	HIS	-	EXPRESSION TAG	UNP P31550
С	18	MSE	-	EXPRESSION TAG	UNP P31550


Continued on next page...




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	${f Comment}$	Reference
D	-2	MSE	-	EXPRESSION TAG	UNP P31550
D	-1	GLY	-	EXPRESSION TAG	UNP P31550
D	0	SER	=	EXPRESSION TAG	UNP P31550
D	1	SER	-	EXPRESSION TAG	UNP P31550
D	2	HIS	=	EXPRESSION TAG	UNP P31550
D	3	HIS	-	EXPRESSION TAG	UNP P31550
D	4	HIS	-	EXPRESSION TAG	UNP P31550
D	5	HIS	-	EXPRESSION TAG	UNP P31550
D	6	HIS	-	EXPRESSION TAG	UNP P31550
D	7	HIS	-	EXPRESSION TAG	UNP P31550
D	8	SER	-	EXPRESSION TAG	UNP P31550
D	9	SER	_	EXPRESSION TAG	UNP P31550
D	10	GLY	-	EXPRESSION TAG	UNP P31550
D	11	LEU	-	EXPRESSION TAG	UNP P31550
D	12	VAL	-	EXPRESSION TAG	UNP P31550
D	13	PRO	-	EXPRESSION TAG	UNP P31550
D	14	ARG	_	EXPRESSION TAG	UNP P31550
D	15	GLY	-	EXPRESSION TAG	UNP P31550
D	16	SER	-	EXPRESSION TAG	UNP P31550
D	17	HIS	-	EXPRESSION TAG	UNP P31550
D	18	MSE	-	EXPRESSION TAG	UNP P31550

 $\bullet \ \, \text{Molecule 2 is THIAMIN PHOSPHATE (three-letter code: TPS) (formula: $C_{12}H_{18}N_4O_4PS)$.}$



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
9	Λ.	1	Total	С	N	О	Р	S	0	0
	A	1	22	12	4	4	1	1	0	0





Continued from previous page...

Mol	Chain	Residues		${f Atoms}$					ZeroOcc	AltConf	
2	D	1	Total	С	N	Ο	Р	S	0	0	
2	Б	1	22	12	4	4	1	1	0	0	
9	С	1	Total	С	N	О	Р	S	0	0	
		1	22	12	4	4	1	1		U	
2	D	1	Total	С	N	О	Р	S	0	0	
	$\begin{array}{c c} 2 & D \end{array}$	1	22	12	4	4	1	1	0	U	

• Molecule 3 is water.


Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	A	103	Total O 103 103	0	0
3	В	138	Total O 138 138	0	0
3	С	127	Total O 127 127	0	0
3	D	132	Total O 132 132	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Thiamine-binding periplasmic protein



• Molecule 1: Thiamine-binding periplasmic protein




• Molecule 1: Thiamine-binding periplasmic protein







• Molecule 1: Thiamine-binding periplasmic protein





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	69.04Å 118.07Å 90.33Å	Depositor
a, b, c, α , β , γ	90.00° 93.74° 90.00°	Depositor
Resolution (Å)	49.39 - 2.25	Depositor
Resolution (A)	49.39 - 2.24	EDS
% Data completeness	90.1 (49.39-2.25)	Depositor
(in resolution range)	89.7 (49.39-2.24)	EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$< I/\sigma(I) > 1$	2.97 (at 2.25Å)	Xtriage
Refinement program	CNS 1.2	Depositor
D D.	0.198 , 0.241	Depositor
R, R_{free}	0.191 , 0.234	DCC
R_{free} test set	3086 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 43.2	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10279	wwPDB-VP
Average B, all atoms $(Å^2)$	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TPS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI		RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.34	0/2455	0.59	0/3341	
1	В	0.36	0/2472	0.58	0/3363	
1	С	0.35	0/2531	0.58	0/3442	
1	D	0.35	0/2474	0.59	1/3366~(0.0%)	
All	All	0.35	0/9932	0.58	$1/13512 \ (0.0\%)$	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	D	213	LEU	CA-CB-CG	-5.04	103.71	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2395	0	2345	90	0
1	В	2412	0	2364	68	0
1	С	2470	0	2418	80	0
1	D	2414	0	2364	76	0
2	A	22	0	16	1	0
2	В	22	0	16	2	0

Continued on next page...



I'amtamaiad	tmom	mmonianale	maaa
Continued	110116	DICUIUUS	Daue
0 0 10001000000	.,	F . C C C C C C C	1 00.3 0 1 1 1

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
2	С	22	0	16	2	0
2	D	22	0	16	3	0
3	A	103	0	0	5	0
3	В	138	0	0	5	0
3	С	127	0	0	3	0
3	D	132	0	0	3	0
All	All	10279	0	9555	306	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 306 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{array}{c} { m Clash} \\ { m overlap} \ ({ m \AA}) \end{array}$
1:A:135:LYS:HD3	1:A:135:LYS:H	1.14	1.11
1:A:62:SER:HA	1:A:65:ASN:ND2	1.74	1.02
1:C:135:LYS:H	1:C:135:LYS:HD3	1.26	0.97
1:A:272:GLN:NE2	1:A:284:VAL:H	1.64	0.94
1:C:93:THR:HG23	1:C:95:LEU:HD13	1.49	0.94

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles		
1	A	306/330~(93%)	293 (96%)	12 (4%)	1 (0%)	41 46		
1	В	307/330 (93%)	297 (97%)	9 (3%)	1 (0%)	41 46		
1	С	316/330 (96%)	305 (96%)	10 (3%)	1 (0%)	41 46		
1	D	308/330 (93%)	289 (94%)	17 (6%)	2 (1%)	25 25		
All	All	1237/1320 (94%)	1184 (96%)	48 (4%)	5 (0%)	34 37		



All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	60	GLY
1	С	16	SER
1	D	286	ASN
1	A	50	CYS
1	В	47	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$249/266 \ (94\%)$	239 (96%)	10 (4%)	31 37
1	В	252/266~(95%)	242 (96%)	10 (4%)	31 37
1	С	$257/266 \ (97\%)$	244 (95%)	13 (5%)	24 25
1	D	$252/266 \ (95\%)$	244 (97%)	8 (3%)	39 47
All	All	1010/1064 (95%)	969 (96%)	41 (4%)	30 36

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	287	VAL
1	С	93	THR
1	D	165	LEU
1	В	322	GLN
1	С	51	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	260	GLN
1	С	65	ASN
1	D	172	GLN
1	В	286	ASN
1	С	107	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	$\mid \# Z > 2$	Counts	RMSZ	# Z >2
2	TPS	В	400	-	20,23,23	1.96	2 (10%)	25,33,33	2.48	10 (40%)
2	TPS	A	400	-	20,23,23	1.39	1 (5%)	25,33,33	2.68	11 (44%)
2	TPS	D	400	-	20,23,23	1.71	2 (10%)	25,33,33	2.63	10 (40%)
2	TPS	С	400	-	20,23,23	1.72	3 (15%)	25,33,33	2.54	10 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	\mathbf{Type}	Chain	${ m Res}$	Link	Chirals	Torsions	Rings
2	TPS	В	400	-	-	1/10/11/11	0/2/2/2
2	TPS	A	400	-	-	1/10/11/11	0/2/2/2
2	TPS	D	400	-	-	1/10/11/11	0/2/2/2
2	TPS	С	400	_	-	1/10/11/11	0/2/2/2



The v	vorst	5	of	8	bond	length	outliers	are	listed	below:
_ I II	W OIDU	$\mathbf{\mathcal{I}}$	OI	\circ	DOM	10115011	Outilities	CULC	IDUCA	DOION.

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(ext{\AA})$
2	В	400	TPS	C4-N3	-5.93	1.34	1.39
2	D	400	TPS	C4-N3	-5.29	1.35	1.39
2	С	400	TPS	C4-N3	-5.14	1.35	1.39
2	В	400	TPS	C6-C5	-4.79	1.48	1.50
2	D	400	TPS	C6-C5	-3.68	1.49	1.50

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
2	A	400	TPS	C6-C5-C4	8.32	134.11	127.43
2	D	400	TPS	C6-C5-C4	8.21	134.02	127.43
2	С	400	TPS	C6-C5-C4	7.65	133.57	127.43
2	В	400	TPS	C6-C5-C4	6.90	132.97	127.43
2	D	400	TPS	C5-C4-N3	4.71	117.00	107.57

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	400	TPS	C4-C5-C6-C7
2	С	400	TPS	C4-C5-C6-C7
2	A	400	TPS	C4-C5-C6-C7
2	D	400	TPS	C4-C5-C6-C7

There are no ring outliers.


4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	400	TPS	2	0
2	A	400	TPS	1	0
2	D	400	TPS	3	0
2	С	400	TPS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the




average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Rings

Torsions



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	$\begin{array}{c cccc} \textbf{Analysed} & <& RSRZ> & \#RSRZ>2 \end{array}$		$OWAB(A^2)$	Q < 0.9	
1	A	304/330 (92%)	-0.12	6 (1%) 65 68	14, 28, 44, 60	0
1	В	$305/330 \; (92\%)$	-0.23	1 (0%) 94 94	12, 24, 41, 59	0
1	С	313/330 (94%)	-0.24	5 (1%) 72 74	14, 25, 41, 49	0
1	D	$305/330 \; (92\%)$	-0.23	2 (0%) 87 88	11, 25, 41, 56	0
All	All	$1227/1320 \ (92\%)$	-0.21	14 (1%) 80 82	11, 25, 43, 60	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	291	ALA	3.5
1	С	16	SER	2.9
1	D	105	ALA	2.8
1	A	132	ASN	2.4
1	С	73	ASN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f A}^2)$	Q<0.9
2	TPS	A	400	22/22	0.94	0.15	15,21,27,31	0
2	TPS	D	400	22/22	0.94	0.14	13,19,24,28	0
2	TPS	В	400	22/22	0.97	0.15	14,16,23,27	0
2	TPS	С	400	22/22	0.97	0.14	13,19,20,24	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around TPS D 400:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around TPS B 400:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

