



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2023 – 12:21 PM EDT

PDB ID : 2QUX  
Title : PP7 coat protein dimer in complex with RNA hairpin  
Authors : Chao, J.A.  
Deposited on : 2007-08-06  
Resolution : 2.44 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)
RNA backbone	3102	1062 (2.80-2.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	25	
1	F	25	
1	I	25	
1	L	25	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	O	25	
1	R	25	
2	A	125	
2	B	125	
2	D	125	
2	E	125	
2	G	125	
2	H	125	
2	J	125	
2	K	125	
2	M	125	
2	N	125	
2	P	125	
2	Q	125	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	L	26	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 15026 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (25-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	25	533	239	98	172	24	0	0	0
1	F	25	533	239	98	172	24	0	0	0
1	I	25	533	239	98	172	24	0	0	0
1	L	25	533	239	98	172	24	0	0	0
1	O	25	533	239	98	172	24	0	0	0
1	R	25	533	239	98	172	24	0	0	0

- Molecule 2 is a protein called Coat protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	123	953	595	170	187	1	0	1	0
2	B	121	934	582	168	184		0	0	0
2	D	122	942	587	169	185	1	0	0	0
2	E	122	950	592	172	185	1	0	1	0
2	G	123	948	590	170	187	1	0	0	0
2	H	119	921	575	166	180		0	0	0
2	J	123	948	590	170	187	1	0	0	0
2	K	122	940	586	169	184	1	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	M	119	Total	C	N	O	0	0	0	
			919	573	165	181				
2	N	119	Total	C	N	O	0	0	0	
			921	575	166	180				
2	P	122	Total	C	N	O	S	0	0	0
			942	587	169	185	1			
2	Q	117	Total	C	N	O		0	0	0
			908	567	164	177				

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q38062
A	-2	GLY	-	expression tag	UNP Q38062
A	-1	SER	-	expression tag	UNP Q38062
A	67	SER	-	linker	UNP Q38062
A	68	GLY	-	linker	UNP Q38062
B	-3	GLY	-	expression tag	UNP Q38062
B	-2	GLY	-	expression tag	UNP Q38062
B	-1	SER	-	expression tag	UNP Q38062
B	67	SER	-	linker	UNP Q38062
B	68	GLY	-	linker	UNP Q38062
D	-3	GLY	-	expression tag	UNP Q38062
D	-2	GLY	-	expression tag	UNP Q38062
D	-1	SER	-	expression tag	UNP Q38062
D	67	SER	-	linker	UNP Q38062
D	68	GLY	-	linker	UNP Q38062
E	-3	GLY	-	expression tag	UNP Q38062
E	-2	GLY	-	expression tag	UNP Q38062
E	-1	SER	-	expression tag	UNP Q38062
E	67	SER	-	linker	UNP Q38062
E	68	GLY	-	linker	UNP Q38062
G	-3	GLY	-	expression tag	UNP Q38062
G	-2	GLY	-	expression tag	UNP Q38062
G	-1	SER	-	expression tag	UNP Q38062
G	67	SER	-	linker	UNP Q38062
G	68	GLY	-	linker	UNP Q38062
H	-3	GLY	-	expression tag	UNP Q38062
H	-2	GLY	-	expression tag	UNP Q38062
H	-1	SER	-	expression tag	UNP Q38062
H	67	SER	-	linker	UNP Q38062
H	68	GLY	-	linker	UNP Q38062
J	-3	GLY	-	expression tag	UNP Q38062

Continued on next page...

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
J	-2	GLY	-	expression tag	UNP Q38062
J	-1	SER	-	expression tag	UNP Q38062
J	67	SER	-	linker	UNP Q38062
J	68	GLY	-	linker	UNP Q38062
K	-3	GLY	-	expression tag	UNP Q38062
K	-2	GLY	-	expression tag	UNP Q38062
K	-1	SER	-	expression tag	UNP Q38062
K	67	SER	-	linker	UNP Q38062
K	68	GLY	-	linker	UNP Q38062
M	-3	GLY	-	expression tag	UNP Q38062
M	-2	GLY	-	expression tag	UNP Q38062
M	-1	SER	-	expression tag	UNP Q38062
M	67	SER	-	linker	UNP Q38062
M	68	GLY	-	linker	UNP Q38062
N	-3	GLY	-	expression tag	UNP Q38062
N	-2	GLY	-	expression tag	UNP Q38062
N	-1	SER	-	expression tag	UNP Q38062
N	67	SER	-	linker	UNP Q38062
N	68	GLY	-	linker	UNP Q38062
P	-3	GLY	-	expression tag	UNP Q38062
P	-2	GLY	-	expression tag	UNP Q38062
P	-1	SER	-	expression tag	UNP Q38062
P	67	SER	-	linker	UNP Q38062
P	68	GLY	-	linker	UNP Q38062
Q	-3	GLY	-	expression tag	UNP Q38062
Q	-2	GLY	-	expression tag	UNP Q38062
Q	-1	SER	-	expression tag	UNP Q38062
Q	67	SER	-	linker	UNP Q38062
Q	68	GLY	-	linker	UNP Q38062

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	I	1	Total C O 6 3 3	0	0
3	L	1	Total C O 6 3 3	0	0
3	R	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	E	1	Total C O 6 3 3	0	0
3	G	1	Total C O 6 3 3	0	0
3	H	1	Total C O 6 3 3	0	0
3	J	1	Total C O 6 3 3	0	0
3	J	1	Total C O 6 3 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	K	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	20	Total	O	0	0
			20	20		
4	F	15	Total	O	0	0
			15	15		
4	I	4	Total	O	0	0
			4	4		
4	L	6	Total	O	0	0
			6	6		
4	O	4	Total	O	0	0
			4	4		
4	A	64	Total	O	0	0
			64	64		
4	B	61	Total	O	0	0
			61	61		
4	D	67	Total	O	0	0
			67	67		
4	E	59	Total	O	0	0
			59	59		
4	G	40	Total	O	0	0
			40	40		
4	H	39	Total	O	0	0
			39	39		
4	J	34	Total	O	0	0
			34	34		
4	K	30	Total	O	0	0
			30	30		
4	M	18	Total	O	0	0
			18	18		
4	N	16	Total	O	0	0
			16	16		
4	P	16	Total	O	0	0
			16	16		
4	Q	19	Total	O	0	0
			19	19		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (25-MER)




- Molecule 1: RNA (25-MER)




- Molecule 1: RNA (25-MER)




- Molecule 1: RNA (25-MER)





- Molecule 1: RNA (25-MER)





- Molecule 1: RNA (25-MER)





• Molecule 2: Coat protein





• Molecule 2: Coat protein




• Molecule 2: Coat protein





• Molecule 2: Coat protein





• Molecule 2: Coat protein





• Molecule 2: Coat protein





● Molecule 2: Coat protein





● Molecule 2: Coat protein




● Molecule 2: Coat protein





● Molecule 2: Coat protein



● Molecule 2: Coat protein



● Molecule 2: Coat protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	174.97Å 145.39Å 109.66Å 90.00° 122.94° 90.00°	Depositor
Resolution (Å)	33.80 – 2.44 34.52 – 2.44	Depositor EDS
% Data completeness (in resolution range)	99.1 (33.80-2.44) 99.0 (34.52-2.44)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.175 , 0.236 0.202 , 0.252	Depositor DCC
$R_{free}$ test set	2545 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.5	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	15026	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	C	1.01	0/596	1.82	16/928 (1.7%)
1	F	0.97	0/596	1.88	19/928 (2.0%)
1	I	1.28	8/596 (1.3%)	1.76	13/928 (1.4%)
1	L	0.91	1/596 (0.2%)	1.63	7/928 (0.8%)
1	O	0.83	0/596	1.59	8/928 (0.9%)
1	R	0.86	0/596	1.66	15/928 (1.6%)
2	A	0.60	0/965	0.91	5/1308 (0.4%)
2	B	0.59	0/943	0.86	0/1279
2	D	0.59	0/951	0.86	3/1289 (0.2%)
2	E	0.58	0/962	0.92	2/1303 (0.2%)
2	G	0.56	0/957	0.90	5/1297 (0.4%)
2	H	0.52	0/929	0.82	1/1258 (0.1%)
2	J	0.53	0/957	0.84	3/1297 (0.2%)
2	K	0.51	0/948	0.87	4/1283 (0.3%)
2	M	0.51	0/928	0.77	2/1260 (0.2%)
2	N	0.52	0/929	0.91	6/1258 (0.5%)
2	P	0.49	0/951	0.79	3/1289 (0.2%)
2	Q	0.52	0/915	0.80	0/1237
All	All	0.68	9/14911 (0.1%)	1.15	112/20926 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	P	0	1
2	Q	0	1
All	All	0	2

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	15	G	C8-N7	-8.90	1.25	1.30
1	I	15	G	N7-C5	-7.97	1.34	1.39
1	I	15	G	C5-C4	-6.26	1.33	1.38
1	I	15	G	C6-N1	-5.13	1.35	1.39
1	I	15	G	N9-C8	-5.11	1.34	1.37

The worst 5 of 112 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	99	ARG	NE-CZ-NH2	-10.67	114.97	120.30
2	E	99	ARG	NE-CZ-NH1	10.22	125.41	120.30
2	N	127	ARG	NE-CZ-NH1	10.12	125.36	120.30
2	K	99	ARG	NE-CZ-NH1	9.72	125.16	120.30
2	A	127	ARG	NE-CZ-NH2	-9.39	115.61	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	P	20	SER	Peptide
2	Q	75	LEU	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	533	0	272	0	0
1	F	533	0	272	0	0
1	I	533	0	272	0	0
1	L	533	0	272	0	0
1	O	533	0	272	0	0
1	R	533	0	272	1	0
2	A	953	0	1000	5	0
2	B	934	0	975	3	0
2	D	942	0	984	3	0
2	E	950	0	997	6	0
2	G	948	0	989	6	0
2	H	921	0	965	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	J	948	0	989	4	0
2	K	940	0	984	4	0
2	M	919	0	954	10	0
2	N	921	0	965	10	0
2	P	942	0	984	3	0
2	Q	908	0	950	6	0
3	A	12	0	16	1	0
3	B	12	0	16	0	0
3	D	12	0	16	0	0
3	E	6	0	8	0	0
3	G	6	0	8	0	0
3	H	6	0	8	0	0
3	I	6	0	8	0	0
3	J	12	0	16	1	0
3	K	6	0	8	0	0
3	L	6	0	8	0	0
3	R	6	0	8	0	0
4	A	64	0	0	1	0
4	B	61	0	0	0	0
4	C	20	0	0	0	0
4	D	67	0	0	0	0
4	E	59	0	0	0	0
4	F	15	0	0	0	0
4	G	40	0	0	0	0
4	H	39	0	0	0	0
4	I	4	0	0	0	0
4	J	34	0	0	0	0
4	K	30	0	0	0	0
4	L	6	0	0	0	0
4	M	18	0	0	0	0
4	N	16	0	0	0	0
4	O	4	0	0	0	0
4	P	16	0	0	0	0
4	Q	19	0	0	0	0
All	All	15026	0	13488	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:29:GLU:HG2	2:E:31:VAL:HG22	1.84	0.59
2:M:105:LEU:HD23	2:N:8:VAL:HG21	1.86	0.58
2:H:44:LEU:CD1	2:H:55:VAL:HG22	2.38	0.53
2:P:84:TRP:CE2	2:Q:106:THR:HG21	2.44	0.53
2:E:108:SER:O	2:E:112:THR:HG23	2.09	0.52

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	122/125 (98%)	119 (98%)	3 (2%)	0	100	100
2	B	119/125 (95%)	116 (98%)	3 (2%)	0	100	100
2	D	120/125 (96%)	118 (98%)	2 (2%)	0	100	100
2	E	121/125 (97%)	120 (99%)	1 (1%)	0	100	100
2	G	121/125 (97%)	118 (98%)	3 (2%)	0	100	100
2	H	115/125 (92%)	114 (99%)	1 (1%)	0	100	100
2	J	121/125 (97%)	119 (98%)	2 (2%)	0	100	100
2	K	118/125 (94%)	113 (96%)	5 (4%)	0	100	100
2	M	117/125 (94%)	112 (96%)	5 (4%)	0	100	100
2	N	115/125 (92%)	110 (96%)	5 (4%)	0	100	100
2	P	120/125 (96%)	112 (93%)	8 (7%)	0	100	100
2	Q	111/125 (89%)	107 (96%)	4 (4%)	0	100	100
All	All	1420/1500 (95%)	1378 (97%)	42 (3%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	109/108 (101%)	107 (98%)	2 (2%)	59	71
2	B	106/108 (98%)	104 (98%)	2 (2%)	57	69
2	D	107/108 (99%)	107 (100%)	0	100	100
2	E	108/108 (100%)	106 (98%)	2 (2%)	57	69
2	G	108/108 (100%)	107 (99%)	1 (1%)	78	87
2	H	105/108 (97%)	103 (98%)	2 (2%)	57	69
2	J	108/108 (100%)	108 (100%)	0	100	100
2	K	107/108 (99%)	104 (97%)	3 (3%)	43	56
2	M	104/108 (96%)	103 (99%)	1 (1%)	76	84
2	N	105/108 (97%)	103 (98%)	2 (2%)	57	69
2	P	107/108 (99%)	106 (99%)	1 (1%)	78	87
2	Q	103/108 (95%)	100 (97%)	3 (3%)	42	54
All	All	1277/1296 (98%)	1258 (98%)	19 (2%)	65	76

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	N	127	ARG
2	Q	63	ASP
2	Q	113	SER
2	Q	19	GLN
2	H	67	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	Q	19	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	C	24/25 (96%)	3 (12%)	0
1	F	24/25 (96%)	3 (12%)	1 (4%)
1	I	24/25 (96%)	3 (12%)	0
1	L	24/25 (96%)	5 (20%)	0
1	O	24/25 (96%)	6 (25%)	0
1	R	24/25 (96%)	5 (20%)	0
All	All	144/150 (96%)	25 (17%)	1 (0%)

5 of 25 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	C	13	A
1	C	14	U
1	C	15	G
1	F	13	A
1	F	15	G

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	F	15	G

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	J	129	-	5,5,5	0.51	0	5,5,5	0.67	0
3	GOL	K	128	-	5,5,5	0.40	0	5,5,5	0.43	0
3	GOL	B	128	-	5,5,5	0.42	0	5,5,5	0.82	0
3	GOL	I	26	-	5,5,5	0.61	0	5,5,5	1.29	1 (20%)
3	GOL	A	129	-	5,5,5	0.36	0	5,5,5	0.88	0
3	GOL	A	128	-	5,5,5	0.23	0	5,5,5	0.53	0
3	GOL	R	26	-	5,5,5	0.45	0	5,5,5	0.45	0
3	GOL	B	129	-	5,5,5	0.28	0	5,5,5	0.39	0
3	GOL	E	128	-	5,5,5	0.35	0	5,5,5	0.61	0
3	GOL	J	128	-	5,5,5	0.41	0	5,5,5	0.43	0
3	GOL	G	128	-	5,5,5	0.54	0	5,5,5	0.69	0
3	GOL	H	128	-	5,5,5	0.39	0	5,5,5	0.47	0
3	GOL	L	26	-	5,5,5	0.45	0	5,5,5	0.43	0
3	GOL	D	129	-	5,5,5	0.31	0	5,5,5	0.62	0
3	GOL	D	128	-	5,5,5	0.36	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	J	129	-	-	2/4/4/4	-
3	GOL	K	128	-	-	0/4/4/4	-
3	GOL	B	128	-	-	0/4/4/4	-
3	GOL	I	26	-	-	2/4/4/4	-
3	GOL	A	129	-	-	3/4/4/4	-
3	GOL	A	128	-	-	2/4/4/4	-
3	GOL	R	26	-	-	2/4/4/4	-
3	GOL	B	129	-	-	2/4/4/4	-
3	GOL	E	128	-	-	2/4/4/4	-
3	GOL	J	128	-	-	2/4/4/4	-
3	GOL	G	128	-	-	2/4/4/4	-
3	GOL	H	128	-	-	1/4/4/4	-
3	GOL	L	26	-	-	1/4/4/4	-
3	GOL	D	129	-	-	0/4/4/4	-
3	GOL	D	128	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	I	26	GOL	O1-C1-C2	2.02	119.90	110.20

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	I	26	GOL	O1-C1-C2-C3
3	R	26	GOL	C1-C2-C3-O3
3	R	26	GOL	O2-C2-C3-O3
3	A	128	GOL	O1-C1-C2-C3
3	A	129	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	129	GOL	1	0
3	A	128	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	25/25 (100%)	0.05	2 (8%) 12 9	27, 67, 101, 106	0
1	F	25/25 (100%)	-0.23	0 100 100	39, 66, 86, 92	0
1	I	25/25 (100%)	-0.14	1 (4%) 38 35	36, 75, 90, 96	0
1	L	25/25 (100%)	0.56	4 (16%) 1 1	45, 92, 104, 123	0
1	O	25/25 (100%)	0.47	4 (16%) 1 1	53, 100, 114, 131	0
1	R	25/25 (100%)	0.22	1 (4%) 38 35	58, 81, 110, 114	0
2	A	123/125 (98%)	-0.14	3 (2%) 59 54	21, 38, 75, 102	0
2	B	121/125 (96%)	-0.17	2 (1%) 70 66	17, 34, 69, 93	0
2	D	122/125 (97%)	-0.24	0 100 100	18, 37, 72, 99	0
2	E	122/125 (97%)	-0.13	3 (2%) 57 53	19, 39, 79, 125	0
2	G	123/125 (98%)	-0.00	3 (2%) 59 54	27, 50, 86, 101	0
2	H	119/125 (95%)	0.34	7 (5%) 22 18	30, 56, 90, 117	0
2	J	123/125 (98%)	-0.02	2 (1%) 72 69	28, 46, 83, 98	0
2	K	122/125 (97%)	0.14	7 (5%) 23 20	28, 48, 95, 141	0
2	M	119/125 (95%)	0.50	10 (8%) 11 8	40, 63, 111, 132	0
2	N	119/125 (95%)	0.34	7 (5%) 22 18	38, 62, 103, 130	0
2	P	122/125 (97%)	0.21	4 (3%) 46 43	32, 59, 101, 114	0
2	Q	117/125 (93%)	0.15	1 (0%) 84 83	30, 64, 98, 119	0
All	All	1602/1650 (97%)	0.09	61 (3%) 40 37	17, 51, 100, 141	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	P	49	ALA	7.5
2	E	75	LEU	5.6
2	G	-1	SER	5.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	K	75	LEU	5.2
2	P	48	GLY	4.5

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	129	6/6	0.67	0.24	57,74,81,83	0
3	GOL	L	26	6/6	0.77	0.42	72,88,107,115	0
3	GOL	I	26	6/6	0.79	0.30	39,73,74,82	0
3	GOL	G	128	6/6	0.79	0.21	46,54,85,88	0
3	GOL	R	26	6/6	0.83	0.27	97,104,110,110	0
3	GOL	B	129	6/6	0.86	0.35	52,64,77,80	0
3	GOL	H	128	6/6	0.86	0.18	51,73,82,83	0
3	GOL	J	129	6/6	0.87	0.34	49,86,91,105	0
3	GOL	J	128	6/6	0.91	0.24	57,85,87,90	0
3	GOL	D	129	6/6	0.91	0.20	39,51,68,77	0
3	GOL	K	128	6/6	0.91	0.26	54,72,84,90	0
3	GOL	D	128	6/6	0.92	0.13	53,69,87,89	0
3	GOL	E	128	6/6	0.93	0.15	44,48,57,62	0
3	GOL	B	128	6/6	0.94	0.14	29,41,54,62	0
3	GOL	A	128	6/6	0.95	0.12	38,42,49,62	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.