

Full wwPDB X-ray Structure Validation Report (i)

May 23, 2020 – 09:55 am BST

PDB ID : 6QXG

Title : Crystal structure of His-tag human thymidylate synthase (HT-hTS) in complex

with FdUMP

Authors : Pozzi, C.; Mangani, M.

Deposited on : 2019-03-07

Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			17%	
1	A	325	80%	6% • 12%
			13%	
1	В	325	79%	7% • 12%
			14%	
1	С	325	80%	8% 12%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7267 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Thymidylate synthase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	287	Total	С	C N O S	0	0	0		
1	A	201	2279	1461	398	407	13	U	U	
1	В	285	Total	С	N	О	S	0	0	0
1	Б	200	2248	1442	388	406	12			
1	С	287	Total	С	N	О	S	0	0	0
1		201	2259	1455	389	403	12			0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	initiating methionine	UNP P04818
A	-10	ARG	-	expression tag	UNP P04818
A	-9	GLY	-	expression tag	UNP P04818
A	-8	SER	_	expression tag	UNP P04818
A	-7	HIS	-	expression tag	UNP P04818
A	-6	HIS	_	expression tag	UNP P04818
A	-5	HIS	-	expression tag	UNP P04818
A	-4	HIS	_	expression tag	UNP P04818
A	-3	HIS	_	expression tag	UNP P04818
A	-2	HIS	_	expression tag	UNP P04818
A	-1	GLY	_	expression tag	UNP P04818
A	0	SER	-	expression tag	UNP P04818
В	-11	MET	-	initiating methionine	UNP P04818
В	-10	ARG	-	expression tag	UNP P04818
В	-9	GLY	-	expression tag	UNP P04818
В	-8	SER	_	expression tag	UNP P04818
В	-7	HIS	-	expression tag	UNP P04818
В	-6	HIS	-	expression tag	UNP P04818
В	-5	HIS	-	expression tag	UNP P04818
В	-4	HIS	-	expression tag	UNP P04818
В	-3	HIS	-	expression tag	UNP P04818
В	-2	HIS		expression tag	UNP P04818
В	-1	GLY	-	expression tag	UNP P04818



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	0	SER	-	expression tag	UNP P04818
С	-11	MET	-	initiating methionine	UNP P04818
С	-10	ARG	-	expression tag	UNP P04818
С	-9	GLY	-	expression tag	UNP P04818
С	-8	SER	_	expression tag	UNP P04818
С	-7	HIS	_	expression tag	UNP P04818
С	-6	HIS	_	expression tag	UNP P04818
С	-5	HIS	_	expression tag	UNP P04818
С	-4	HIS	-	expression tag	UNP P04818
С	-3	HIS	-	expression tag	UNP P04818
С	-2	HIS	_	expression tag	UNP P04818
С	-1	GLY	_	expression tag	UNP P04818
С	0	SER	-	expression tag	UNP P04818


• Molecule 2 is 5-FLUORO-2'-DEOXYURIDINE-5'-MONOPHOSPHATE (three-letter code: UFP) (formula: $C_9H_{12}FN_2O_8P$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C F N O P 21 9 1 2 8 1	0	0
2	В	1	Total C F N O P 21 9 1 2 8 1	0	0
2	С	1	Total C F N O P 21 9 1 2 8 1	0	0

 \bullet Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	С	1	Total 5	O 4	S 1	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	133	Total O 133 133	0	0
4	В	135	Total O 135 135	0	0
4	С	145	Total O 145 145	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thymidylate synthase









4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants	103.22Å 61.98Å 169.36Å	Depositor
a, b, c, α , β , γ	90.00° 98.96° 90.00°	Depositor
Resolution (Å)	32.36 - 2.08	Depositor
resolution (A)	32.34 - 2.08	EDS
% Data completeness	97.9 (32.36-2.08)	Depositor
(in resolution range)	97.9 (32.34-2.08)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.87 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
P. P.	0.181 , 0.225	Depositor
R, R_{free}	0.181 , 0.225	DCC
R_{free} test set	3181 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.28, 69.9	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7267	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UFP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.60	0/2339	0.99	0/3170
1	В	0.60	0/2308	0.96	0/3129
1	С	0.56	0/2319	0.93	0/3143
All	All	0.59	0/6966	0.96	0/9442

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	Α	2279	0	2218	18	0
1	В	2248	0	2153	18	0
1	С	2259	0	2192	10	0
2	A	21	0	10	1	0
2	В	21	0	10	1	0
2	С	21	0	10	0	0
3	С	5	0	0	0	0
4	A	133	0	0	1	0
4	В	135	0	0	1	0
4	С	145	0	0	1	0
All	All	7267	0	6593	44	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:ARG:HG3	1:A:163:ARG:HH11	1.32	0.91
1:A:306:THR:HG22	4:A:603:HOH:O	1.85	0.75
1:A:166:ASP:O	1:A:170:THR:HG23	1.89	0.73
1:B:165:ILE:CG2	1:B:240:ILE:HD11	2.20	0.71
1:B:32:GLN:NE2	1:B:64:ARG:O	2.25	0.70
1:A:280:ARG:HH21	1:A:280:ARG:HG2	1.59	0.65
1:B:304:HIS:HB3	1:B:305:PRO:HD2	1.80	0.63
1:B:165:ILE:HG23	1:B:240:ILE:HD11	1.81	0.61
1:B:32:GLN:HE22	1:B:64:ARG:H	1.48	0.60
1:A:280:ARG:CG	1:A:280:ARG:HH21	2.16	0.58
1:B:165:ILE:HG21	1:B:240:ILE:HD11	1.85	0.57
1:A:147:ARG:HH22	1:A:156:GLN:HE22	1.53	0.56
1:A:163:ARG:CG	1:A:163:ARG:HH11	2.13	0.56
1:A:163:ARG:NH1	1:A:163:ARG:HG3	2.12	0.54
1:C:133:PRO:HG2	1:C:186:ASP:HB3	1.89	0.54
1:B:272:GLU:HA	1:B:272:GLU:OE1	2.06	0.54
1:B:82:LYS:O	1:B:86:GLU:HB2	2.08	0.53
1:B:304:HIS:HB3	1:B:305:PRO:CD	2.38	0.53
1:B:198:LEU:HD12	1:B:198:LEU:C	2.30	0.52
1:B:195:CYS:SG	2:B:401:UFP:C6	2.99	0.51
1:A:46:ARG:HH21	1:A:46:ARG:HB2	1.76	0.51
1:A:46:ARG:NH2	1:A:46:ARG:HB2	2.26	0.50
1:B:120:SER:HB2	1:C:301:TYR:O	2.11	0.50
1:B:187:LEU:CD1	1:B:193:PRO:HB3	2.42	0.50
1:B:32:GLN:NE2	1:B:64:ARG:H	2.11	0.48
1:B:187:LEU:HD11	1:B:193:PRO:HB3	1.94	0.47
1:B:56:LEU:HD22	1:B:262:ILE:HD11	1.97	0.47
1:B:77:LYS:HE3	4:B:553:HOH:O	2.13	0.47
1:A:170:THR:HG21	4:C:637:HOH:O	2.15	0.47
1:C:193:PRO:HA	1:C:194:PRO:HD3	1.87	0.46
1:A:170:THR:HG22	1:C:267:ILE:HD13	1.98	0.45
1:A:88:LEU:HD23	1:A:232:LEU:HG	1.99	0.45
1:C:134:VAL:O	1:C:135:TYR:C	2.56	0.44
1:C:126:ARG:HG2	1:C:130:ASP:HB3	1.99	0.44
1:C:198:LEU:C	1:C:198:LEU:HD12	2.38	0.44
1:C:287:LYS:O	1:C:290:ASP:HB2	2.18	0.44
1:C:207:GLU:HA	1:C:244:LYS:O	2.18	0.43



$\alpha \cdots$	· ·	•	
Continued	trom	nromanne	naae
-	110111	picolous	payc

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({f A})$	$oxed{ overlap (\AA) }$
1:A:280:ARG:CG	1:A:280:ARG:NH2	2.79	0.43
1:C:196:HIS:HB3	1:C:212:LEU:HD11	2.01	0.43
1:A:119:ASP:OD1	1:A:124:SER:HA	2.18	0.43
1:A:195:CYS:SG	2:A:401:UFP:C6	3.07	0.43
1:A:285:VAL:CG1	1:A:290:ASP:HB3	2.49	0.42
1:A:285:VAL:HG21	1:A:291:PHE:CE1	2.54	0.42
1:B:74:LEU:HD12	1:B:224:PRO:HB3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	285/325~(88%)	276 (97%)	9 (3%)	0	100	100
1	В	283/325 (87%)	269 (95%)	13 (5%)	1 (0%)	34	31
1	С	$285/325 \ (88\%)$	271 (95%)	14 (5%)	0	100	100
All	All	853/975 (88%)	816 (96%)	36 (4%)	1 (0%)	51	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	${f Res}$	Type
1	В	134	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percer	$_{ m tiles}$
1	A	$238/281 \ (85\%)$	222 (93%)	16 (7%)	16	12
1	В	230/281 (82%)	216 (94%)	14 (6%)	18	15
1	С	232/281 (83%)	222 (96%)	10 (4%)	29	28
All	All	700/843 (83%)	660 (94%)	40 (6%)	20	17

All (40) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	A	46	ARG
1	A	69	ASP
1	A	125	THR
1	A	127	GLU
1	A	147	ARG
1	A A A A A A	163	ARG
1	A	170	THR LEU
1	A	192	LEU
1	A	221	LEU
1	A	240	ILE
1	A A A	280	ARG
1	A	288	ILE
1	A	289	ASP
1	A	294	GLU
1	A A	302	ASN
1	A	306	THR
1	В	49	ASP
1	В	56	LEU
1	В	77	LYS
1	В	86	GLU
1	В	125	THR
1	В	147	ARG
1	В	163	ARG
1	В	176	ARG LEU
1	В	187	LEU
1	В	192	LEU
1	В	198	LEU
1	В	272	GLU
1	В	289	ASP
1	В	294	GLU
1	С	46	ARG
1	С	53	THR
1	C C C	56	LEU
1	С	69	ASP



Continued from previous page...

Mol	Chain	${f Res}$	Type
1	С	121	LEU
1	С	125	THR
1	1 C 151		SER
1	С	176	ARG
1	С	229	SER
1	С	259	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	32	GLN
1	В	39	HIS
1	С	32	GLN
1	С	39	HIS
1	С	62	GLN
1	С	171	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Trino	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	$\mid \# Z > 2$	Counts	RMSZ	# Z > 2
3	SO4	С	402	-	4,4,4	0.33	0	6,6,6	0.10	0
2	UFP	В	401	-	19,22,22	2.03	2 (10%)	24,33,33	2.40	5 (20%)
2	UFP	С	401	-	19,22,22	2.08	1 (5%)	24,33,33	1.93	7 (29%)
2	UFP	A	401	-	19,22,22	1.98	2 (10%)	24,33,33	2.50	8 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UFP	В	401	_	-	1/7/22/22	0/2/2/2
2	UFP	С	401	_	-	2/7/22/22	0/2/2/2
2	UFP	A	401	-	=	1/7/22/22	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	С	401	UFP	C4-C5	8.36	1.48	1.38
2	A	401	UFP	C4-C5	7.75	1.48	1.38
2	В	401	UFP	C4-C5	7.69	1.48	1.38
2	A	401	UFP	C2-N3	-2.38	1.33	1.38
2	В	401	UFP	O3'-C3'	2.34	1.48	1.43

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$\mathbf{Ideal}(^{o})$
2	В	401	UFP	C4-N3-C2	7.82	121.74	115.14
2	A	401	UFP	C4-N3-C2	7.74	121.67	115.14
2	С	401	UFP	C4-N3-C2	5.91	120.13	115.14
2	В	401	UFP	P-O5'-C5'	4.63	131.05	118.30
2	В	401	UFP	C5-C4-N3	-4.59	117.54	122.39
2	A	401	UFP	O2P-P-O1P	4.07	126.63	110.68
2	A	401	UFP	C5-C4-N3	-3.80	118.38	122.39
2	A	401	UFP	P-O5'-C5'	3.53	128.02	118.30
2	С	401	UFP	F5-C5-C6	3.31	124.01	117.51
2	В	401	UFP	F5-C5-C6	2.83	123.07	117.51
2	С	401	UFP	P-O5'-C5'	2.78	125.95	118.30
2	A	401	UFP	F5-C5-C6	2.78	122.97	117.51
2	A	401	UFP	O2P-P-O5'	-2.74	99.45	106.73
2	С	401	UFP	O3P-P-O2P	2.57	117.47	107.64



$\alpha \cdots \tau$	e	•	
Continued	trom	meaningile	maaa
-	110116	piculuas	puyc

Mol	Chain	${f Res}$	Type	${f Atoms}$	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	401	UFP	O4'-C4'-C5'	-2.51	101.11	109.37
2	A	401	UFP	C3'-C2'-C1'	-2.40	96.53	102.54
2	A	401	UFP	O5'-C5'-C4'	-2.24	101.29	108.99
2	С	401	UFP	O5'-P-O1P	2.19	112.61	106.47
2	С	401	UFP	O3P-P-O5'	-2.15	101.02	106.73
2	В	401	UFP	O3'-C3'-C2'	-2.13	103.28	110.90

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	401	UFP	O4'-C4'-C5'-O5'
2	В	401	UFP	O4'-C4'-C5'-O5'
2	С	401	UFP	C3'-C4'-C5'-O5'
2	A	401	UFP	O4'-C4'-C5'-O5'


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	401	UFP	1	0
2	A	401	UFP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	287/325~(88%)	1.06	54 (18%) 1 1	65, 81, 119, 227	2 (0%)
1	В	$285/325 \ (87\%)$	0.96	42 (14%) 2 3	65, 81, 115, 167	3 (1%)
1	С	287/325 (88%)	0.96	44 (15%) 2 2	67, 81, 112, 140	2 (0%)
All	All	859/975 (88%)	0.99	140 (16%) 1 2	65, 81, 116, 227	7 (0%)

All (140) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	LEU	7.5
1	A	184	PRO	7.0
1	A	183	ASN	7.0
1	С	52	GLY	6.5
1	A	187	LEU	6.4
1	A	186	ASP	6.1
1	A	212	LEU	5.9
1	A	213	TYR	5.8
1	В	309	MET	5.7
1	В	212	LEU	5.2
1	С	144	ALA	4.7
1	С	153	TYR	4.6
1	В	51	THR	4.6
1	С	212	LEU	4.4
1	В	306	THR	4.4
1	В	178	ILE	4.4
1	A	120	SER	4.3
1	В	308	LYS	4.2
1	A	252	LEU	4.2
1	В	262	ILE	4.1
1	С	249	ILE	4.1
1	С	123	PHE	3.9
1	В	307	ILE	3.9



Continued from previous page...

	$\frac{nued\ from}{\Box}$			
Mol	Chain	Res	Type	RSRZ
1	C	206	SER	3.8
1	В	249	ILE	3.8
1	С	213	TYR	3.7
1	A	192	LEU	3.7
1	С	26	PRO	3.6
1	В	267	ILE	3.6
1	A	182	TRP	3.6
1	В	198	LEU	3.6
1	A	223	VAL	3.5
1	A	197	ALA	3.4
1	A	178	ILE	3.4
1	A	188	PRO	3.3
1	В	227	ILE	3.3
1	A	198	LEU	3.3
1	В	264	PRO	3.3
1	В	52	GLY	3.2
1	С	154	SER	3.2
1	С	150	GLU	3.2
1	A	253	GLY	3.2
1	С	227	ILE	3.2
1	С	128	GLU	3.2
1	A	103	SER	3.1
1	С	198	LEU	3.1
1	A	251	THR	3.0
1	A	121	LEU	3.0
1	A	227	ILE	3.0
1	С	41	LEU	3.0
1	A	180	CYS	3.0
1	В	213	TYR	3.0
1	С	124	SER	3.0
1	В	270	GLN	3.0
1	A	311	MET	2.9
1	В	180	CYS	2.9
1	С	122	GLY	2.9
1	С	180	CYS	2.9
1	A	52	GLY	2.8
1	С	156	GLN	2.8
1	A	26	PRO	2.8
1	A	285	VAL	2.8
1	A	46	ARG	2.8
1	A	283	ARG	2.8
1	A	195	CYS	2.8
		l	l	



Continued from previous page...

	$nued\ from$			
Mol	Chain	Res	Type	RSRZ
1	В	53	THR	2.8
1	A	214	GLN	2.8
1	A	215	ARG	2.8
1	В	271	ARG	2.8
1	C	84	VAL	2.8
1	С	148	ASP	2.8
1	A	185	ARG	2.7
1	A	128	GLU	2.7
1	В	230	TYR	2.7
1	С	195	CYS	2.7
1	С	251	THR	2.7
1	С	39	HIS	2.6
1	В	305	PRO	2.6
1	A	125	THR	2.6
1	С	211	GLN	2.6
1	A	216	SER	2.6
1	A	181	ALA	2.6
1	В	182	TRP	2.6
1	A	281	ILE	2.6
1	A	290	ASP	2.6
1	A	210	CYS	2.6
1	В	248	PHE	2.6
1	С	210	CYS	2.5
1	A	280	ARG	2.5
1	В	197	ALA	2.5
1	В	304	HIS	2.4
1	В	113	GLY	2.4
1	A	163	ARG	2.4
1	A	211	GLN	2.4
1	В	251	THR	2.4
1	С	50	ARG	2.4
1	A	177	ILE	2.4
1	С	121	LEU	2.3
1	В	74	LEU	2.3
1	В	43	CYS	2.3
1	A	179	MET	2.3
1	В	283	ARG	2.3
1	С	116	ASP	2.3
1	В	214	GLN	2.3
1	В	128	GLU	2.3
1	В	252	LEU	2.3
1	В	123	PHE	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	223	VAL	2.2
1	A	297	GLN	2.2
1	С	145	GLU	2.2
1	С	292	LYS	2.2
1	A	249	ILE	2.2
1	A	113	GLY	2.2
1	С	114	SER	2.2
1	С	229	SER	2.2
1	В	272	GLU	2.2
1	A	233	LEU	2.2
1	С	59	PHE	2.2
1	В	211	GLN	2.2
1	С	197	ALA	2.2
1	В	79	VAL	2.2
1	A	147	ARG	2.1
1	В	199	CYS	2.1
1	В	196	HIS	2.1
1	В	265	LEU	2.1
1	A	270	GLN	2.1
1	A	229	SER	2.1
1	С	51	THR	2.1
1	A	255	ALA	2.0
1	В	45	VAL	2.0
1	С	214	GLN	2.0
1	В	195	CYS	2.0
1	С	234	THR	2.0
1	A	59	PHE	2.0
1	A	60	GLY	2.0
1	С	217	GLY	2.0
1	С	102	SER	2.0
1	С	208	LEU	2.0
1	С	233	LEU	2.0
1	С	252	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ extbf{B-factors}(extbf{A}^2)$	Q < 0.9
3	SO4	С	402	5/5	0.78	0.32	117,122,124,127	0
2	UFP	С	401	21/21	0.94	0.13	58,68,76,100	0
2	UFP	В	401	21/21	0.95	0.11	62,68,81,103	0
2	UFP	A	401	21/21	0.95	0.14	59,66,74,109	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around UFP B 401: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around UFP A 401: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.

