

# wwPDB X-ray Structure Validation Summary Report (i)

Jan 13, 2024 – 06:07 pm GMT

PDB ID : 6QXL

Title : Crystal Structure of Pyruvate Kinase II (PykA) from Pseudomonas aeruginosa

in complex with sodium malonate, magnesium and glucose-6-phosphate

Authors: Abdelhamid, Y.; Brear, P.; Welch, M.

Deposited on : 2019-03-07

Resolution : 2.43 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	483	91%	8% •
1	В	483	94%	5% •
1	С	483	93%	6% •
1	D	483	94%	5%
1	Е	483	92%	7% •



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	483	92%	8% •
1	G	483	93%	6% •
1	Н	483	92%	7%
1	I	483	91%	8% •
1	J	483	78%	16%
1	K	483	93%	6% •
1	L	483	89%	9% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MLI	D	504	-	X	-	-



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 45923 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Pyruvate kinase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	483	Total	С	N	О	S	0	0	0
1	A	400	3662	2286	661	695	20	0	U	
1	В	483	Total	С	N	О	S	0	0	0
1	Ъ	400	3662	2286	661	695	20	U	U	U
1	С	483	Total	С	N	О	S	0	0	0
1		400	3662	2286	661	695	20	U	U	U
1	D	483	Total	С	N	Ο	S	0	0	0
1	D	400	3662	2286	661	695	20	U	U	
1	E	483	Total	С	N	О	S	0	2	0
1	Е	400	3680	2296	664	700	20	U	2	
1	F	483	Total	$\mathbf{C}$	N	O	$\mathbf{S}$	0	0	0
1	I.	400	3662	2286	661	695	20	U	U	U
1	G	483	Total	$\mathbf{C}$	N	O	S	0	0	0
1	d	400	3662	2286	661	695	20	O	O	0
1	Н	483	Total	$\mathbf{C}$	N	O	S	0	0	0
1	11	400	3662	2286	661	695	20	O	O	U
1	I	483	Total	С	N	O	S	0	0	0
1	1	400	3662	2286	661	695	20	O	U	0
1	J	404	Total	С	N	О	S	0	0	0
1	3	404	3057	1910	555	575	17	O	U	
1	K	482	Total	С	N	О	S	0	0	0
1	17	102	3654	2282	660	692	20	0	U	U
1	L	483	Total	$\mathbf{C}$	N	O	$\mathbf{S}$	0	0	0
1	L	100	3662	2286	661	695	20		U	

• Molecule 2 is 6-O-phosphono-alpha-D-glucopyranose (three-letter code: G6P) (formula:  $C_6H_{13}O_9P$ ) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
2	A	1	Total	С	О	Р	0	0
	A	1	16	6	9	1	0	0
2	В	1	Total	С	О	Р	0	0
	Б	1	16	6	9	1	U	0
2	С	1	Total	С	О	Р	0	0
2		1	16	6	9	1	U	
2	D	1	Total	С	О	Р	0	0
	D	1	16	6	9	1	U	U
2	E	1	Total	С	Ο	Р	0	0
	Ш	1	16	6	9	1	0	0
2	F	1	Total	С	Ο	Р	0	0
	1	1	16	6	9	1	0	U
2	G	1	Total	С	Ο	Р	0	0
	<u> </u>	1	16	6	9	1	0	Ŭ.
2	Н	1	Total	С	Ο	Р	0	0
	11	1	16	6	9	1	0	0
2	I	1	Total	С	Ο	Р	0	0
	1	1	16	6	9	1	0	
2	J	1	Total	С	Ο	Р	0	0
	3	1	16	6	9	1	0	0
2	K	1	Total	С	Ο	Р	0	0
	11	1	16	6	9	1		0
2	L	1	Total	С	Ο	Р	0	0
	L	1	16	6	9	1		

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	Е	1	Total Mg 1 1	0	0
3	F	1	Total Mg 1 1	0	0
3	G	1	Total Mg 1 1	0	0
3	Н	1	Total Mg 1 1	0	0
3	I	1	Total Mg 1 1	0	0
3	J	1	Total Mg 1 1	0	0
3	K	1	Total Mg 1 1	0	0
3	L	1	Total Mg 1 1	0	0


 $\bullet$  Molecule 4 is MALONATE ION (three-letter code: MLI) (formula:  $\rm C_3H_2O_4)$  (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 3 4	0	0
4	В	1	Total C O 7 3 4	0	0
4	С	1	Total C O 7 3 4	0	0
4	D	1	Total C O 7 3 4	0	0
4	Е	1	Total C O 7 3 4	0	0
4	F	1	Total C O 7 3 4	0	0
4	G	1	Total C O 7 3 4	0	0
4	Н	1	Total C O 7 3 4	0	0
4	I	1	Total C O 7 3 4	0	0
4	J	1	Total C O 7 3 4	0	0
4	K	1	Total C O 7 3 4	0	0
4	L	1	Total C O 7 3 4	0	0

 $\bullet$  Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $\mathrm{C_3H_8O_3}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0
5	E	1	Total C O 6 3 3	0	0

#### • Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	238	Total O 238 238	0	0
6	В	301	Total O 301 301	0	0
6	С	245	Total O 245 245	0	0
6	D	272	Total O 272 272	0	0
6	Е	270	Total O 270 270	0	0
6	F	164	Total O 164 164	0	0
6	G	154	Total O 154 154	0	0
6	Н	150	Total O 150 150	0	0
6	I	114	Total O 114 114	0	0
6	J	128	Total O 128 128	0	0
6	К	150	Total O 150 150	0	0
6	L	82	Total O 82 82	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyruvate kinase









• Molecule 1: Pyruvate kinase

Chain J: 78% . . 16%

#48

#48

#48

#48

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477

#477



0412 A413 R414 E422 D462 V475

• Molecule 1: Pyruvate kinase

Chain K: 93% 6% •





• Molecule 1: Pyruvate kinase

Chain L: 89% 9% •







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants	182.48Å 182.48Å 405.04Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	158.03 - 2.43	Depositor
Resolution (A)	158.03 - 2.43	EDS
% Data completeness	98.9 (158.03-2.43)	Depositor
(in resolution range)	98.7 (158.03-2.43)	EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.77 (at 2.43Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
Ρ. Р.	0.227 , $0.251$	Depositor
$R, R_{free}$	0.221 , $0.246$	DCC
$R_{free}$ test set	14353 reflections $(4.96%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.809	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33, 42.9	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.014 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	45923	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.47% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MLI, G6P, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Во	ond lengths	E	Bond angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5
1	A	0.69	3/3716 (0.1%)	0.86	19/5025 (0.4%)
1	В	0.73	1/3716 (0.0%)	0.91	15/5025 (0.3%)
1	С	0.68	$1/3716 \ (0.0\%)$	0.93	19/5025 (0.4%)
1	D	0.66	0/3716	0.85	9/5025 (0.2%)
1	Е	0.70	0/3734	0.90	16/5049 (0.3%)
1	F	0.72	3/3716 (0.1%)	0.95	$15/5025 \ (0.3\%)$
1	G	0.67	1/3716 (0.0%)	0.90	$16/5025 \ (0.3\%)$
1	Н	0.72	1/3716~(0.0%)	0.92	13/5025 (0.3%)
1	I	0.71	$1/3716 \ (0.0\%)$	0.96	21/5025 (0.4%)
1	J	0.75	3/3103 (0.1%)	0.99	15/4196 (0.4%)
1	K	0.69	0/3707	0.92	14/5011 (0.3%)
1	L	0.77	3/3716 (0.1%)	1.11	34/5025 (0.7%)
All	All	0.71	17/43988 (0.0%)	0.94	206/59481 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	В	0	3
1	С	0	1
1	D	0	1
1	Е	0	1
1	G	0	2
1	Н	0	1
1	J	0	2
1	K	0	1
1	L	0	2
All	All	0	16



The worst	5	of	17	bond	length	outliers	are	listed	below:
110 WOID	$\overline{}$	01		OILG	10115011	CULTUID	COL C	IIDCC	CIC III

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$Ideal(\AA)$
1	J	157	HIS	N-CA	-11.85	1.22	1.46
1	J	254	GLU	CD-OE1	-8.73	1.16	1.25
1	L	464	TYR	CG-CD2	-7.38	1.29	1.39
1	A	254	GLU	CG-CD	-6.30	1.42	1.51
1	F	72	PRO	C-O	-6.21	1.10	1.23

The worst 5 of 206 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	L	272	ARG	NE-CZ-NH2	22.93	131.77	120.30
1	J	76	ILE	O-C-N	-18.61	92.93	122.70
1	F	53	ARG	NE-CZ-NH2	18.36	129.48	120.30
1	J	254	GLU	OE1-CD-OE2	-15.19	105.07	123.30
1	L	198	ASP	CB-CG-OD1	14.69	131.52	118.30

There are no chirality outliers.

5 of 16 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	ASP	Mainchain
1	A	81	ASN	Mainchain
1	В	114	ASP	Mainchain
1	В	230	ASP	Sidechain
1	В	81	ASN	Mainchain

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3662	0	3726	26	0
1	В	3662	0	3726	15	0
1	С	3662	0	3726	19	0
1	D	3662	0	3726	13	0
1	Е	3680	0	3738	21	0
1	F	3662	0	3726	28	0
1	G	3662	0	3726	16	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	H(added)	Clashes	Symm-Clashes
1	Н	3662	0	3726	19	0
1	I	3662	0	3726	24	0
1	J	3057	0	3123	21	0
1	K	3654	0	3721	14	0
1	L	3662	0	3726	20	0
2	A	16	0	11	1	0
2	В	16	0	11	0	0
2	С	16	0	11	0	0
2	D	16	0	11	0	0
2	Е	16	0	11	0	0
2	F	16	0	11	2	0
2	G	16	0	11	0	0
2	Н	16	0	11	0	0
2	I	16	0	11	0	0
2	J	16	0	11	0	0
2	K	16	0	11	0	0
2	L	16	0	11	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
3	Е	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	Н	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
4	A	7	0	2	0	0
4	В	7	0	2	0	0
4	С	7	0	2	0	0
4	D	7	0	2	0	0
4	E	7	0	2	1	0
4	F	7	0	2	0	0
4	G	7	0	2	0	0
4	Н	7	0	2	0	0
4	I	7	0	2	0	0
4	J	7	0	2	0	0
4	K	7	0	2	1	0
4	L	7	0	2	0	0
5	В	6	0	8	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	6	0	8	0	0
5	Ε	6	0	8	0	0
6	A	238	0	0	10	0
6	В	301	0	0	7	0
6	С	245	0	0	7	0
6	D	272	0	0	7	0
6	Ε	270	0	0	8	0
6	F	164	0	0	3	0
6	G	154	0	0	4	0
6	Н	150	0	0	3	0
6	I	114	0	0	3	0
6	J	128	0	0	8	0
6	K	150	0	0	1	0
6	L	82	0	0	2	0
All	All	45923	0	44296	225	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:263:ILE:HD11	6:A:791:HOH:O	1.01	1.18
1:C:187:ASP:OD2	6:C:601:HOH:O	1.79	0.98
1:B:378:LYS:HE3	6:B:785:HOH:O	1.65	0.96
1:E:207:ARG:HD2	6:E:771:HOH:O	1.65	0.95
1:J:73:LYS:HG3	6:J:643:HOH:O	1.67	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	481/483 (100%)	469 (98%)	11 (2%)	1 (0%)	47	57
1	В	481/483 (100%)	469 (98%)	11 (2%)	1 (0%)	47	57
1	С	481/483 (100%)	467 (97%)	13 (3%)	1 (0%)	47	57
1	D	481/483 (100%)	465 (97%)	15 (3%)	1 (0%)	47	57
1	E	483/483 (100%)	468 (97%)	14 (3%)	1 (0%)	47	57
1	F	481/483 (100%)	464 (96%)	15 (3%)	2 (0%)	34	41
1	G	481/483 (100%)	466 (97%)	14 (3%)	1 (0%)	47	57
1	Н	481/483 (100%)	467 (97%)	13 (3%)	1 (0%)	47	57
1	I	481/483 (100%)	465 (97%)	15 (3%)	1 (0%)	47	57
1	J	400/483 (83%)	389 (97%)	10 (2%)	1 (0%)	41	49
1	K	478/483 (99%)	461 (96%)	16 (3%)	1 (0%)	47	57
1	L	481/483 (100%)	468 (97%)	12 (2%)	1 (0%)	47	57
All	All	5690/5796 (98%)	5518 (97%)	159 (3%)	13 (0%)	47	57

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	282	THR
1	В	282	THR
1	С	282	THR
1	D	282	THR
1	Е	282	THR

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$386/386 \ (100\%)$	385 (100%)	1 (0%)	92 95
1	В	386/386 (100%)	385 (100%)	1 (0%)	92 95
1	С	386/386 (100%)	384 (100%)	2 (0%)	88 93
1	D	386/386 (100%)	384 (100%)	2 (0%)	88 93



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	E	388/386 (100%)	386 (100%)	2 (0%)	88 93
1	F	386/386 (100%)	384 (100%)	2 (0%)	88 93
1	G	386/386 (100%)	384 (100%)	2 (0%)	88 93
1	Н	386/386 (100%)	385 (100%)	1 (0%)	92 95
1	I	386/386 (100%)	383 (99%)	3 (1%)	81 88
1	J	320/386 (83%)	319 (100%)	1 (0%)	92 95
1	K	385/386 (100%)	382 (99%)	3 (1%)	81 88
1	L	386/386 (100%)	385 (100%)	1 (0%)	92 95
All	All	4567/4632 (99%)	4546 (100%)	21 (0%)	88 93

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	309	TYR
1	K	83	ARG
1	L	309	TYR
1	K	168	THR
1	J	309	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	283	GLN
1	K	467	GLN
1	L	350	HIS
1	F	467	GLN
1	Е	350	HIS

## 5.3.3 RNA (i)

There are no RNA molecules in this entry.

# 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 39 ligands modelled in this entry, 12 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

TA /F - 1	<b>m</b>	C1 :	D	T ! 1	Bond lengths			Bond angles		
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MLI	F	503	3	6,6,6	1.07	0	7,7,7	1.74	2 (28%)
2	G6P	L	501	-	16,16,16	0.74	0	24,24,24	1.83	10 (41%)
4	MLI	I	503	3	6,6,6	1.32	0	7,7,7	1.32	0
4	MLI	${ m L}$	503	3	6,6,6	1.18	0	7,7,7	1.65	1 (14%)
5	GOL	D	503	-	5,5,5	0.49	0	5,5,5	0.86	0
2	G6P	J	501	-	16,16,16	0.92	0	24,24,24	1.46	4 (16%)
2	G6P	С	501	-	16,16,16	0.73	0	24,24,24	1.75	9 (37%)
5	GOL	Ε	503	-	5,5,5	0.31	0	5,5,5	0.66	0
2	G6P	Η	501	-	16,16,16	0.65	0	24,24,24	1.51	5 (20%)
2	G6P	В	501	-	16,16,16	1.09	1 (6%)	24,24,24	2.20	9 (37%)
2	G6P	G	501	-	16,16,16	0.78	0	24,24,24	1.58	5 (20%)
4	MLI	Ε	504	3	6,6,6	0.98	0	7,7,7	1.53	2 (28%)
2	G6P	Ι	501	ı	16,16,16	0.84	1 (6%)	24,24,24	1.76	7 (29%)
4	MLI	G	503	3	6,6,6	1.03	0	7,7,7	2.28	4 (57%)
4	MLI	D	504	3	6,6,6	1.14	0	7,7,7	2.53	3 (42%)
4	MLI	С	503	3	6,6,6	1.04	0	7,7,7	2.02	3 (42%)
4	MLI	J	503	3	6,6,6	1.61	1 (16%)	7,7,7	0.93	0
4	MLI	В	504	3	6,6,6	1.28	0	7,7,7	1.48	1 (14%)
2	G6P	E	501	-	16,16,16	1.02	1 (6%)	24,24,24	1.95	8 (33%)
2	G6P	F	501	-	16,16,16	0.52	0	24,24,24	1.67	6 (25%)
2	G6P	A	501	-	16,16,16	0.83	0	24,24,24	2.59	12 (50%)
2	G6P	D	501	-	16,16,16	1.06	2 (12%)	24,24,24	2.03	6 (25%)
4	MLI	K	503	3	6,6,6	1.02	0	7,7,7	1.65	2 (28%)



Mol	Tuno	Chain	Res	Link Bond lengths			Bond angles			
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	G6P	K	501	-	16,16,16	0.89	1 (6%)	24,24,24	1.69	5 (20%)
5	GOL	В	503	-	5,5,5	0.45	0	5,5,5	0.52	0
4	MLI	Н	503	3	6,6,6	1.44	0	7,7,7	0.74	0
4	MLI	A	503	3	6,6,6	1.36	1 (16%)	7,7,7	1.17	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLI	F	503	3	-	4/4/4/4	-
2	G6P	L	501	-	-	2/6/26/26	0/1/1/1
4	MLI	I	503	3	-	2/4/4/4	-
4	MLI	L	503	3	-	2/4/4/4	-
5	GOL	D	503	-	-	3/4/4/4	-
2	G6P	J	501	-	-	5/6/26/26	0/1/1/1
2	G6P	С	501	-	-	3/6/26/26	0/1/1/1
5	GOL	Е	503	-	-	2/4/4/4	-
2	G6P	Н	501	-	-	2/6/26/26	0/1/1/1
2	G6P	В	501	-	-	4/6/26/26	0/1/1/1
2	G6P	G	501	-	-	2/6/26/26	0/1/1/1
4	MLI	Е	504	3	-	2/4/4/4	-
2	G6P	I	501	-	-	5/6/26/26	0/1/1/1
4	MLI	G	503	3	-	2/4/4/4	-
4	MLI	D	504	3	-	4/4/4/4	-
4	MLI	С	503	3	-	3/4/4/4	-
4	MLI	J	503	3	-	4/4/4/4	-
4	MLI	В	504	3	-	2/4/4/4	-
2	G6P	Е	501	-	-	1/6/26/26	0/1/1/1
2	G6P	F	501	-	-	5/6/26/26	0/1/1/1
2	G6P	A	501	-	-	5/6/26/26	0/1/1/1
2	G6P	D	501	-	-	3/6/26/26	0/1/1/1
4	MLI	K	503	3	-	2/4/4/4	-
2	G6P	K	501	-	-	1/6/26/26	0/1/1/1
5	GOL	В	503	-	-	2/4/4/4	-
4	MLI	Н	503	3	-	$\frac{4/4/4/4}{2}$	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLI	A	503	3	-	2/4/4/4	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\mathring{\mathrm{A}})$	$\operatorname{Ideal}(\operatorname{\AA})$
2	Ε	501	G6P	O1-C1	2.59	1.47	1.39
2	D	501	G6P	O1-C1	2.38	1.47	1.39
2	В	501	G6P	C1-C2	-2.32	1.46	1.52
4	A	503	MLI	O7-C2	-2.26	1.23	1.30
4	J	503	MLI	C1-C2	2.20	1.54	1.51

The worst 5 of 105 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
2	A	501	G6P	O2P-P-O6	-6.97	88.18	106.73
2	D	501	G6P	O5-C1-C2	-6.14	99.33	110.28
2	В	501	G6P	O5-C1-C2	-5.67	100.17	110.28
2	Е	501	G6P	C1-O5-C5	5.06	123.21	113.66
2	F	501	G6P	C4-C3-C2	-5.00	102.09	110.82

There are no chirality outliers.

5 of 78 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	G6P	C4-C5-C6-O6
2	A	501	G6P	O5-C5-C6-O6
2	A	501	G6P	C6-O6-P-O1P
2	A	501	G6P	C6-O6-P-O2P
2	В	501	G6P	C6-O6-P-O1P

There are no ring outliers.


4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Е	504	MLI	1	0
2	F	501	G6P	2	0
2	A	501	G6P	1	0
4	K	503	MLI	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In




addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































































































### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(\AA^2)$	Q<0.9
1	A	483/483 (100%)	-0.16	2 (0%) 92 92	27, 47, 74, 105	0
1	В	483/483 (100%)	-0.20	0 100 100	22, 37, 67, 98	0
1	С	483/483 (100%)	-0.22	2 (0%) 92 92	27, 44, 72, 110	0
1	D	483/483 (100%)	-0.21	0 100 100	25, 42, 71, 102	0
1	E	483/483 (100%)	-0.20	1 (0%) 95 95	24, 39, 73, 92	0
1	F	483/483 (100%)	-0.06	3 (0%) 89 89	26, 53, 84, 111	0
1	G	483/483 (100%)	-0.10	6 (1%) 79 77	30, 56, 90, 116	0
1	Н	483/483 (100%)	-0.01	9 (1%) 66 63	23, 54, 102, 138	0
1	I	483/483 (100%)	-0.04	7 (1%) 75 73	30, 62, 91, 114	0
1	J	404/483 (83%)	-0.03	5 (1%) 79 77	28, 55, 93, 214	0
1	K	482/483 (99%)	-0.06	4 (0%) 86 85	31, 58, 92, 128	0
1	L	483/483 (100%)	0.34	31 (6%) 19 16	37, 71, 118, 153	0
All	All	5716/5796 (98%)	-0.08	70 (1%) 79 77	22, 52, 92, 214	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	76	ILE	6.3
1	L	79	PHE	4.9
1	L	104	THR	4.2
1	L	77	ALA	4.2
1	L	82	LYS	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	MG	L	502	1/1	0.68	0.09	75,75,75,75	0
3	MG	I	502	1/1	0.75	0.07	30,30,30,30	0
3	MG	В	502	1/1	0.77	0.17	44,44,44,44	0
4	MLI	L	503	7/7	0.77	0.16	64,79,81,87	0
5	GOL	D	503	6/6	0.86	0.18	45,54,60,62	0
3	MG	С	502	1/1	0.89	0.10	30,30,30,30	0
3	MG	A	502	1/1	0.90	0.05	49,49,49,49	0
4	MLI	K	503	7/7	0.90	0.14	61,67,70,84	0
3	MG	G	502	1/1	0.91	0.14	45,45,45,45	0
4	MLI	Ι	503	7/7	0.91	0.14	55,62,65,83	0
5	GOL	В	503	6/6	0.91	0.16	44,52,58,58	0
4	MLI	J	503	7/7	0.91	0.16	59,61,69,75	0
3	MG	J	502	1/1	0.92	0.04	55,55,55,55	0
3	MG	Ε	502	1/1	0.93	0.16	40,40,40,40	0
4	MLI	В	504	7/7	0.94	0.09	37,42,45,46	0
4	MLI	Н	503	7/7	0.94	0.10	47,52,54,57	0
2	G6P	F	501	16/16	0.94	0.15	53,80,96,104	0
2	G6P	G	501	16/16	0.94	0.18	49,81,94,100	0
2	G6P	K	501	16/16	0.95	0.16	51,77,92,98	0
3	MG	F	502	1/1	0.95	0.09	49,49,49,49	0
2	G6P	Н	501	16/16	0.95	0.16	50,69,82,92	0
4	MLI	С	503	7/7	0.95	0.12	38,50,64,68	0
4	MLI	D	504	7/7	0.95	0.11	33,41,49,50	0
3	MG	D	502	1/1	0.95	0.17	39,39,39,39	0
5	GOL	Ε	503	6/6	0.95	0.13	40,49,51,57	0
2	G6P	I	501	16/16	0.96	0.16	47,62,72,84	0
2	G6P	A	501	16/16	0.96	0.15	40,59,69,74	0
2	G6P	L	501	16/16	0.96	0.13	59,80,90,92	0
4	MLI	A	503	7/7	0.96	0.15	45,49,54,62	0
2	G6P	В	501	16/16	0.96	0.16	33,51,67,103	0
2	G6P	С	501	16/16	0.96	0.15	39,54,79,84	0
3	MG	Н	502	1/1	0.96	0.07	49,49,49,49	0


Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
4	MLI	Е	504	7/7	0.96	0.09	29,34,42,46	0
4	MLI	F	503	7/7	0.97	0.12	46,58,61,64	0
4	MLI	G	503	7/7	0.97	0.09	52,54,60,60	0
2	G6P	D	501	16/16	0.97	0.16	36,47,60,74	0
2	G6P	Е	501	16/16	0.97	0.17	42,60,75,78	0
2	G6P	J	501	16/16	0.97	0.17	58,85,105,109	0
3	MG	K	502	1/1	0.98	0.12	48,48,48,48	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





























### 










### Electron density around MLI I 503:


 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)
























## 










### Electron density around G6P F 501:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around G6P G 501:


 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)


























### Electron density around MLI C 503:


 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









































### Electron density around G6P B 501:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around G6P C 501:


 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)















### Electron density around MLI E 504:

 $2mF_o$ -DF<sub>c</sub> (at 0.7 rmsd) in gray  $mF_o$ -DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)










# Electron density around MLI F 503: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)



## Electron density around MLI G 503: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around G6P E 501: $2 {\rm mF}_o\text{-}{\rm DF}_c \mbox{ (at } 0.7 \mbox{ rmsd) in gray}$

 ${
m mF}_o{
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around G6P J 501:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











### 6.5 Other polymers (i)

There are no such residues in this entry.

