



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 3, 2023 – 12:06 AM EDT

PDB ID : 3QYJ  
Title : Crystal structure of ALR0039, a putative alpha/beta hydrolase from Nostoc sp PCC 7120.  
Authors : Petit, P.; Kagan, O.; Yakunin, A.F.; Savchenko, A.  
Deposited on : 2011-03-03  
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	291	 2% 86% 12%
1	B	291	 % 85% 13%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5366 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alr0039 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	291	2377	1526	415	424	12	0	4	0
1	B	290	2355	1513	408	422	12	0	2	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	325	Total 325	O 325	0	0
2	B	309	Total 309	O 309	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alr0039 protein



- Molecule 1: Alr0039 protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.54Å 88.20Å 83.59Å 90.00° 93.09° 90.00°	Depositor
Resolution (Å)	19.86 – 1.78 19.86 – 1.78	Depositor EDS
% Data completeness (in resolution range)	97.4 (19.86-1.78) 97.4 (19.86-1.78)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.17 (at 1.78Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.149 , 0.179 0.157 , 0.185	Depositor DCC
$R_{free}$ test set	2021 reflections (3.35%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtrriage
Anisotropy	0.583	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5366	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.07	3/2459 (0.1%)	0.93	5/3344 (0.1%)
1	B	1.13	6/2430 (0.2%)	1.06	14/3305 (0.4%)
All	All	1.10	9/4889 (0.2%)	1.00	19/6649 (0.3%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	257	ARG	CG-CD	-6.41	1.35	1.51
1	B	256	GLU	CG-CD	5.47	1.60	1.51
1	B	278	GLU	CG-CD	5.46	1.60	1.51
1	A	15	ALA	CA-CB	5.39	1.63	1.52
1	A	228	GLN	CG-CD	5.21	1.63	1.51
1	B	135[A]	MET	SD-CE	-5.20	1.48	1.77
1	B	135[B]	MET	SD-CE	-5.20	1.48	1.77
1	A	226	MET	SD-CE	-5.09	1.49	1.77
1	B	67	ARG	CG-CD	-5.01	1.39	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	135[A]	MET	CG-SD-CE	-11.27	82.17	100.20
1	B	135[B]	MET	CG-SD-CE	-11.27	82.17	100.20
1	B	16	ARG	NE-CZ-NH1	11.25	125.92	120.30
1	B	16	ARG	NE-CZ-NH2	-9.57	115.52	120.30
1	B	257	ARG	NE-CZ-NH2	-9.29	115.66	120.30
1	B	71	VAL	CG1-CB-CG2	7.84	123.44	110.90
1	B	196	ARG	NE-CZ-NH2	-7.55	116.53	120.30
1	A	135	MET	CG-SD-CE	-7.34	88.45	100.20
1	B	212	ARG	NE-CZ-NH1	6.89	123.74	120.30
1	A	140	ASP	CB-CG-OD1	6.81	124.42	118.30
1	B	80	ARG	NE-CZ-NH2	-6.54	117.03	120.30
1	B	249	ASP	CB-CG-OD1	6.22	123.90	118.30
1	A	255	ARG	NE-CZ-NH2	-5.97	117.32	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	196	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	B	80	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	A	137	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	B	71	VAL	N-CA-CB	-5.23	99.99	111.50
1	A	11	ASP	CB-CG-OD1	5.20	122.98	118.30
1	B	212	ARG	NE-CZ-NH2	-5.18	117.71	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2377	0	2326	30	0
1	B	2355	0	2302	24	0
2	A	325	0	0	8	0
2	B	309	0	0	6	0
All	All	5366	0	4628	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:HIS:HD2	2:A:407:HOH:O	1.69	0.73
1:B:11:ASP:OD1	1:B:16:ARG:HD2	1.89	0.72
1:B:249:ASP:O	1:B:253:THR:HG23	1.90	0.72
1:A:123:LYS:NZ	1:A:291:CYS:SG	2.59	0.69
1:B:117:HIS:HD2	2:B:320:HOH:O	1.75	0.69
1:A:119[A]:HIS:HE1	2:A:565:HOH:O	1.77	0.68
1:B:38:HIS:HE1	2:B:466:HOH:O	1.82	0.61
1:B:103:HIS:HE1	1:B:270:HIS:O	1.83	0.61
1:A:119[A]:HIS:CD2	1:A:119[A]:HIS:H	2.17	0.59
1:A:42:HIS:HD2	2:A:340:HOH:O	1.85	0.58

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:CYS:HB3	2:A:582:HOH:O	2.05	0.57
1:A:67:ARG:HH12	1:A:200:GLN:NE2	2.02	0.56
1:B:42:HIS:HD2	2:B:363:HOH:O	1.87	0.56
1:B:74:HIS:HD2	1:B:209:GLU:OE1	1.88	0.56
1:B:242:GLY:N	2:B:561:HOH:O	2.32	0.55
1:A:38:HIS:HE1	2:A:615:HOH:O	1.90	0.54
1:A:28:LEU:HD11	1:A:101:VAL:HG23	1.88	0.54
1:B:103:HIS:HD2	2:B:293:HOH:O	1.90	0.54
1:A:103:HIS:HE1	1:A:270:HIS:O	1.90	0.54
1:B:38:HIS:HD2	1:B:58:ASP:OD1	1.93	0.52
1:B:123:LYS:NZ	1:B:290:HIS:CE1	2.77	0.52
1:B:42:HIS:HE1	2:B:292:HOH:O	1.92	0.52
1:A:38:HIS:HD2	1:A:58:ASP:OD1	1.92	0.51
1:B:123:LYS:HZ1	1:B:290:HIS:CE1	2.27	0.51
1:A:74:HIS:HD2	1:A:209:GLU:OE1	1.94	0.51
1:A:257[B]:ARG:HH11	1:A:257[B]:ARG:HG2	1.76	0.50
1:A:33:GLY:HA3	1:A:104:ASP:HB3	1.94	0.50
1:A:187:HIS:HD2	1:A:189:GLN:H	1.59	0.49
1:B:231:SER:HA	1:B:259:ILE:HG12	1.95	0.48
1:A:119[A]:HIS:CE1	2:A:565:HOH:O	2.59	0.48
1:A:135:MET:HE1	1:A:143:PHE:HE1	1.78	0.48
1:A:42:HIS:HE1	2:A:309:HOH:O	1.98	0.47
1:B:187:HIS:CD2	1:B:189:GLN:H	2.33	0.47
1:B:39:VAL:HG11	1:B:190:ALA:HA	1.97	0.47
1:A:119[A]:HIS:H	1:A:119[A]:HIS:HD2	1.63	0.46
1:A:222:ASP:O	1:A:226:MET:HE2	2.16	0.45
1:A:135:MET:CE	1:A:143:PHE:HE1	2.30	0.45
1:B:187:HIS:HD2	1:B:189:GLN:H	1.64	0.45
1:A:249:ASP:O	1:A:253:THR:HG23	2.18	0.44
1:A:110:ALA:HB1	1:A:124:LEU:HD11	2.00	0.44
1:A:135:MET:HE1	1:A:143:PHE:CE1	2.53	0.44
1:A:217:ILE:HD13	1:B:92:LYS:HE2	1.99	0.44
1:B:133:HIS:CE1	1:B:137:ARG:HG3	2.53	0.44
1:A:133:HIS:CE1	1:A:137:ARG:HG3	2.53	0.43
1:B:33:GLY:HA3	1:B:104:ASP:HB3	2.00	0.42
1:B:236:VAL:O	1:B:263:GLY:HA3	2.20	0.42
1:B:135[A]:MET:CE	1:B:143:PHE:CE1	3.03	0.41
1:A:124:LEU:O	1:A:234:VAL:HA	2.20	0.41
1:A:244:ILE:HG23	1:A:245:GLY:N	2.35	0.41
1:A:103:HIS:HD2	2:A:297:HOH:O	2.03	0.41
1:B:61:GLY:HA2	1:B:67:ARG:HB3	2.03	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:LEU:HD11	1:B:101:VAL:HG23	2.03	0.41
1:A:167:ASN:C	1:A:167:ASN:HD22	2.25	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/291 (101%)	285 (97%)	8 (3%)	0	100	100
1	B	290/291 (100%)	283 (98%)	7 (2%)	0	100	100
All	All	583/582 (100%)	568 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/248 (102%)	250 (99%)	2 (1%)	81	76
1	B	249/248 (100%)	245 (98%)	4 (2%)	62	51
All	All	501/496 (101%)	495 (99%)	6 (1%)	71	62

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	ARG
1	A	167	ASN
1	B	67	ARG
1	B	71	VAL
1	B	127	LEU
1	B	191	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	HIS
1	A	42	HIS
1	A	74	HIS
1	A	103	HIS
1	A	117	HIS
1	A	155	GLN
1	A	158	ASN
1	A	167	ASN
1	A	187	HIS
1	A	200	GLN
1	B	38	HIS
1	B	42	HIS
1	B	74	HIS
1	B	103	HIS
1	B	117	HIS
1	B	155	GLN
1	B	187	HIS
1	B	290	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	291/291 (100%)	-0.21	5 (1%) 70 70	14, 20, 32, 43	0
1	B	290/291 (99%)	-0.26	4 (1%) 75 75	12, 17, 30, 54	0
All	All	581/582 (99%)	-0.24	9 (1%) 73 73	12, 19, 31, 54	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	291	CYS	7.4
1	B	290	HIS	4.8
1	A	24	HIS	4.5
1	B	243	ILE	4.1
1	B	246	ARG	3.6
1	B	24	HIS	2.8
1	A	3	THR	2.5
1	A	69	ALA	2.4
1	A	4	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.