

wwPDB X-ray Structure Validation Summary Report (i)

Jan 9, 2024 – 04:04 pm GMT

PDB ID	:	8QYF
Title	:	Crystal structure of ClpP from Staphylococcus epidermidis in complex with
		ixazomib
Authors	:	Franca, B.A.; Rohde, H.; Betzel, C.
Deposited on		
Resolution	:	2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	200	83% 8% 109	%
1	В	200	82% 6% 11%	-
1	С	200	82% 6% 10%	•
1	D	200	84% 5% 10%	-
1	Е	200	84% 6% 10%)



Mol	Chain	Length	Quality of chain	
1	F	200	80% 10%	10%
1	G	200	82% 7%	10%
1	Н	200	2% 84% 6%	10%
1	Ι	200	82% 8%	10%
1	J	200	83% 6%	10%
1	Κ	200	2% 81% 8%	11%
1	L	200	82% 7%	11%
1	М	200	82% 7%	10%
1	Ν	200	86% 5%	10%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 19892 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	181	Total	С	Ν	0	S	0	0	0
	Л	101	1371	862	232	271	6	0	0	0
1	В	178	Total	С	Ν	Ο	S	0	0	0
	D	110	1359	855	230	268	6	0	0	0
1	С	179	Total	С	Ν	Ο	S	0	0	0
1	C	115	1347	845	228	268	6	0	0	0
1	D	179	Total	С	Ν	Ο	\mathbf{S}	0	0	0
	D	115	1366	860	232	268	6	0	0	0
1	Е	179	Total	С	Ν	Ο	S	0	1	0
1	Ľ	115	1371	863	233	269	6	0	I	0
1	F	179	Total	С	Ν	Ο	S	0	1	0
L	Ľ	115	1364	856	232	270	6	0	L	
1	G	179	Total	С	Ν	0	S	0	1	0
1	G	115	1352	850	231	265	6	0	T	0
1	Н	179	Total	С	Ν	0	S	0	1	0
1	11	113	1364	857	231	270	6	0		
1	Ι	179	Total	С	Ν	0	S	0	1	0
	L	113	1372	864	232	270	6	0	1	0
1	J	179	Total	С	Ν	Ο	S	0	0	0
	0	113	1336	842	226	262	6	0	0	0
1	Κ	178	Total	С	Ν	Ο	\mathbf{S}	0	0	0
	IX	170	1332	838	229	259	6	0	0	0
1	L	178	Total	С	Ν	Ο	S	0	0	0
		110	1328	835	228	259	6	0		U
1	М	179	Total	С	Ν	0	S	0	0	0
	111	113	1358	855	231	266	6	0		U
1	Ν	181	Total	С	Ν	0	S	0	0	0
	11	101	1371	864	233	268	6	0	0	0

• Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference				
А	195	HIS	-	expression tag	UNP A0A0N1MQL5				



Chain	Residue	Modelled	Actual	Comment	Reference
А	196	HIS	-	expression tag	UNP A0A0N1MQL5
А	197	HIS	-	expression tag	UNP A0A0N1MQL5
А	198	HIS	-	expression tag	UNP A0A0N1MQL5
А	199	HIS	_	expression tag	UNP A0A0N1MQL5
А	200	HIS	-	expression tag	UNP A0A0N1MQL5
В	195	HIS	-	expression tag	UNP A0A0N1MQL5
В	196	HIS	-	expression tag	UNP A0A0N1MQL5
В	197	HIS	-	expression tag	UNP A0A0N1MQL5
В	198	HIS	-	expression tag	UNP A0A0N1MQL5
В	199	HIS	-	expression tag	UNP A0A0N1MQL5
В	200	HIS	-	expression tag	UNP A0A0N1MQL5
С	195	HIS	-	expression tag	UNP A0A0N1MQL5
С	196	HIS	-	expression tag	UNP A0A0N1MQL5
С	197	HIS	-	expression tag	UNP A0A0N1MQL5
С	198	HIS	-	expression tag	UNP A0A0N1MQL5
С	199	HIS	-	expression tag	UNP A0A0N1MQL5
С	200	HIS	-	expression tag	UNP A0A0N1MQL5
D	195	HIS	-	expression tag	UNP A0A0N1MQL5
D	196	HIS	-	expression tag	UNP A0A0N1MQL5
D	197	HIS	-	expression tag	UNP A0A0N1MQL5
D	198	HIS	-	expression tag	UNP A0A0N1MQL5
D	199	HIS	-	expression tag	UNP A0A0N1MQL5
D	200	HIS	-	expression tag	UNP A0A0N1MQL5
E	195	HIS	-	expression tag	UNP A0A0N1MQL5
E	196	HIS	-	expression tag	UNP A0A0N1MQL5
E	197	HIS	-	expression tag	UNP A0A0N1MQL5
E	198	HIS	-	expression tag	UNP A0A0N1MQL5
E	199	HIS	-	expression tag	UNP A0A0N1MQL5
E	200	HIS	-	expression tag	UNP A0A0N1MQL5
F	195	HIS	-	expression tag	UNP A0A0N1MQL5
F	196	HIS	-	expression tag	UNP A0A0N1MQL5
F	197	HIS	-	expression tag	UNP A0A0N1MQL5
F	198	HIS	-	expression tag	UNP A0A0N1MQL5
F	199	HIS	_	expression tag	UNP A0A0N1MQL5
F	200	HIS	-	expression tag	UNP A0A0N1MQL5
G	195	HIS	-	expression tag	UNP A0A0N1MQL5
G	196	HIS		expression tag	UNP A0A0N1MQL5
G	197	HIS	-	expression tag	UNP A0A0N1MQL5
G	198	HIS	-	expression tag	UNP A0A0N1MQL5
G	199	HIS	-	expression tag	UNP A0A0N1MQL5
G	200	HIS	-	expression tag	UNP A0A0N1MQL5
Н	195	HIS	-	expression tag	UNP A0A0N1MQL5




Chain	Residue	vious page Modelled	Actual	Comment	Reference
Н	196	HIS	_	expression tag	UNP A0A0N1MQL5
Н	197	HIS	-	expression tag	UNP A0A0N1MQL5
Н	198	HIS	-	expression tag	UNP A0A0N1MQL5
Н	199	HIS	-	expression tag	UNP A0A0N1MQL5
Н	200	HIS	-	expression tag	UNP A0A0N1MQL5
Ι	195	HIS	-	expression tag	UNP A0A0N1MQL5
Ι	196	HIS	-	expression tag	UNP A0A0N1MQL5
Ι	197	HIS	-	expression tag	UNP A0A0N1MQL5
Ι	198	HIS	-	expression tag	UNP A0A0N1MQL5
Ι	199	HIS	-	expression tag	UNP A0A0N1MQL5
Ι	200	HIS	-	expression tag	UNP A0A0N1MQL5
J	195	HIS	-	expression tag	UNP A0A0N1MQL5
J	196	HIS	-	expression tag	UNP A0A0N1MQL5
J	197	HIS	-	expression tag	UNP A0A0N1MQL5
J	198	HIS	-	expression tag	UNP A0A0N1MQL5
J	199	HIS	-	expression tag	UNP A0A0N1MQL5
J	200	HIS	-	expression tag	UNP A0A0N1MQL5
K	195	HIS	-	expression tag	UNP A0A0N1MQL5
K	196	HIS	-	expression tag	UNP A0A0N1MQL5
K	197	HIS	-	expression tag	UNP A0A0N1MQL5
K	198	HIS	-	expression tag	UNP A0A0N1MQL5
K	199	HIS	-	expression tag	UNP A0A0N1MQL5
K	200	HIS	-	expression tag	UNP A0A0N1MQL5
L	195	HIS	-	expression tag	UNP A0A0N1MQL5
L	196	HIS	-	expression tag	UNP A0A0N1MQL5
L	197	HIS	-	expression tag	UNP A0A0N1MQL5
L	198	HIS	-	expression tag	UNP A0A0N1MQL5
L	199	HIS	-	expression tag	UNP A0A0N1MQL5
L	200	HIS	-	expression tag	UNP A0A0N1MQL5
М	195	HIS	-	expression tag	UNP A0A0N1MQL5
М	196	HIS	-	expression tag	UNP A0A0N1MQL5
М	197	HIS	-	expression tag	UNP A0A0N1MQL5
М	198	HIS	-	expression tag	UNP A0A0N1MQL5
М	199	HIS	-	expression tag	UNP A0A0N1MQL5
М	200	HIS	-	expression tag	UNP A0A0N1MQL5
N	195	HIS	-	expression tag	UNP A0A0N1MQL5
N	196	HIS	-	expression tag	UNP A0A0N1MQL5
N	197	HIS	-	expression tag	UNP A0A0N1MQL5
N	198	HIS	-	expression tag	UNP A0A0N1MQL5
N	199	HIS	-	expression tag	UNP A0A0N1MQL5
N	200	HIS	-	expression tag	UNP A0A0N1MQL5

• Molecule 2 is [(1 {R})-1-[2-[[2,5-bis(chloranyl)phenyl]carbonylamino]ethanoylamino]-3-



methyl-butyl]boronic acid (three-letter code: 6V8) (formula: $C_{14}H_{19}BCl_2N_2O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total B C Cl N O	0	0
	Л	I	23 1 14 2 2 4	0	0
2	В	1	Total B C Cl N O	0	0
	D	1	23 1 14 2 2 4	0	0
2	С	1	Total B C Cl N O	0	0
		1	23 1 14 2 2 4		0
2	D	1	Total B C Cl N O	0	0
		-	23 1 14 2 2 4		
2	Е	1	Total B C Cl N O	0	0
		_	23 1 14 2 2 4		
2	F	1	Total B C Cl N O	0	0
2	G	1	Total B C Cl N O	0	0
			$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		
2	Н	1	Total B C Cl N O	0	0
			$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		
2	Ι	1	Total B C Cl N O	0	0
			<u>23 1 14 2 2 4</u>		
2	J	1	Total B C Cl N O	0	0
			<u>23 1 14 2 2 4</u> Tatal P C Cl N O		
2	Κ	1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0	0
2	L	1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0	0
			23 1 14 2 2 4		



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
0	2 M	1	Total	В	С	Cl	Ν	Ο	0	0
		1	23	1	14	2	2	4	0	0
0	2 N	1	Total	В	С	Cl	Ν	Ο	0	0
		1	23	1	14	2	2	4	0	0

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	40	Total O 40 40	0	0
3	В	43	Total O 43 43	0	0
3	С	44	Total O 44 44	0	0
3	D	51	Total O 51 51	0	0
3	Е	43	Total O 43 43	0	0
3	F	46	Total O 46 46	0	0
3	G	41	Total O 41 41	0	0
3	Н	52	Total O 52 52	0	0
3	Ι	36	Total O 36 36	0	0
3	J	34	$\begin{array}{cc} \text{Total} & \text{O} \\ 34 & 34 \end{array}$	0	0
3	K	22	TotalO2222	0	0
3	L	30	Total O 30 30	0	0
3	М	44	Total O 44 44	0	0
3	Ν	53	Total O 53 53	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-dependent Clp protease proteolytic subunit



• Molecule 1: ATP-dependent Clp protease proteolytic subunit





• Molecule 1: ATP-dependent Clp protease proteolytic subunit



Chain M: 82%	7% 10%
MET MET ASIN LEU LEU LEU LEU LEU THR THR THR THR ASIN ASIN ASIN ASIN ASIN ASIN ASIN ASIN	CLVS CLVS HLS HLS HLS HLS HLS HLS HLS HLS HLS
• Molecule 1: ATP-dependent Clp protease proteol	ytic subunit
Chain N: 86%	5% 10%

MET LASN LASN LASN ASN ASN ASN ASN ASN ASN ASN ASN ASN
--



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	94.96Å 123.16Å 126.24Å	Deneiten
a, b, c, α , β , γ	90.00° 91.23° 90.00°	Depositor
Resolution (Å)	48.01 - 2.33	Depositor
Resolution (A)	48.01 - 2.20	EDS
% Data completeness	99.9 (48.01-2.33)	Depositor
(in resolution range)	99.9 (48.01-2.20)	EDS
R _{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.20 (at 2.20 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.20.1-4487-000	Depositor
D D	0.211 , 0.221	Depositor
R, R_{free}	0.210 , 0.224	DCC
R_{free} test set	2101 reflections (1.43%)	wwPDB-VP
Wilson B-factor $(Å^2)$	39.9	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 33.3	EDS
L-test for twinning ²	$< L > = 0.50, < L^2 > = 0.34$	Xtriage
	0.000 for -h,l,k	
Estimated twinning fraction	0.007 for -h,-l,-k	Xtriage
	0.019 for h,-k,-l	
F_o, F_c correlation	0.95	EDS
Total number of atoms	19892	wwPDB-VP
Average B, all atoms $(Å^2)$	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $6\mathrm{V8}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bo	ond angles
WIOI	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.50	0/1388	0.66	0/1877
1	В	0.48	0/1377	0.67	1/1862~(0.1%)
1	С	0.48	0/1364	0.68	1/1849~(0.1%)
1	D	0.52	0/1384	0.66	0/1872
1	Ε	0.52	0/1392	0.68	0/1883
1	F	0.50	0/1384	0.66	0/1873
1	G	0.48	0/1372	0.66	0/1858
1	Н	0.49	0/1385	0.64	0/1875
1	Ι	0.56	1/1390~(0.1%)	0.67	1/1880~(0.1%)
1	J	0.49	1/1354~(0.1%)	0.65	0/1838
1	Κ	0.45	0/1349	0.66	1/1827~(0.1%)
1	L	0.50	0/1346	0.66	0/1826
1	М	0.57	2/1376~(0.1%)	0.70	1/1863~(0.1%)
1	Ν	0.52	0/1389	0.68	0/1881
All	All	0.51	4/19250~(0.0%)	0.67	5/26064~(0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	М	18	TYR	CB-CG	-6.99	1.41	1.51
1	Ι	18	TYR	CB-CG	-6.54	1.41	1.51
1	J	191	GLU	CB-CG	-6.06	1.40	1.52
1	М	18	TYR	CD1-CE1	-6.00	1.30	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	Κ	4	ILE	CA-CB-CG1	-6.61	98.45	111.00
1	Ι	4	ILE	CG1-CB-CG2	6.57	125.86	111.40
1	С	4	ILE	CG1-CB-CG2	-6.30	97.54	111.40



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	М	18	TYR	CB-CG-CD1	-5.33	117.80	121.00
1	В	4	ILE	CG1-CB-CG2	-5.09	100.21	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1371	0	1364	9	0
1	В	1359	0	1349	7	0
1	С	1347	0	1318	10	0
1	D	1366	0	1360	5	0
1	Е	1371	0	1366	8	0
1	F	1364	0	1352	13	0
1	G	1352	0	1338	8	0
1	Н	1364	0	1344	7	0
1	Ι	1372	0	1365	9	0
1	J	1336	0	1304	7	0
1	Κ	1332	0	1315	16	0
1	L	1328	0	1293	9	0
1	М	1358	0	1345	6	0
1	N	1371	0	1361	6	0
2	А	23	0	0	0	0
2	В	23	0	0	1	0
2	С	23	0	0	2	0
2	D	23	0	0	0	0
2	Е	23	0	0	2	0
2	F	23	0	0	0	0
2	G	23	0	0	0	0
2	Н	23	0	0	0	0
2	Ι	23	0	0	1	0
2	J	23	0	0	0	0
2	Κ	23	0	0	1	0
2	L	23	0	0	1	0
2	М	23	0	0	$\begin{bmatrix} 0 \\ C & t \end{bmatrix}$	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Ν	23	0	0	0	0
3	А	40	0	0	0	0
3	В	43	0	0	0	0
3	С	44	0	0	0	0
3	D	51	0	0	0	0
3	Ε	43	0	0	0	0
3	F	46	0	0	1	0
3	G	41	0	0	0	0
3	Η	52	0	0	0	0
3	Ι	36	0	0	0	0
3	J	34	0	0	0	0
3	Κ	22	0	0	0	0
3	L	30	0	0	0	0
3	М	44	0	0	0	0
3	Ν	53	0	0	0	0
All	All	19892	0	18774	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 101 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:152:ARG:CZ	1:K:162:ILE:HD11	2.11	0.80
1:K:152:ARG:CZ	1:K:162:ILE:CD1	2.62	0.77
1:C:4:ILE:HG12	1:C:5:PRO:HD3	1.71	0.72
1:I:42[B]:ASN:ND2	1:J:33:GLY:O	2.28	0.65
1:H:6:THR:O	1:N:22:SER:HB3	2.00	0.62

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



80	YF
પહ્ય	тт

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	177/200~(88%)	174 (98%)	3~(2%)	0	100	100
1	В	174/200~(87%)	171 (98%)	3~(2%)	0	100	100
1	С	175/200~(88%)	172 (98%)	3~(2%)	0	100	100
1	D	175/200~(88%)	172 (98%)	3~(2%)	0	100	100
1	Ε	176/200~(88%)	172 (98%)	4 (2%)	0	100	100
1	F	176/200~(88%)	173 (98%)	3(2%)	0	100	100
1	G	176/200~(88%)	173 (98%)	3 (2%)	0	100	100
1	Н	176/200~(88%)	173 (98%)	3(2%)	0	100	100
1	Ι	176/200~(88%)	173~(98%)	3~(2%)	0	100	100
1	J	175/200~(88%)	172 (98%)	3(2%)	0	100	100
1	Κ	174/200~(87%)	171 (98%)	3~(2%)	0	100	100
1	L	174/200~(87%)	171 (98%)	3(2%)	0	100	100
1	М	175/200~(88%)	172 (98%)	3 (2%)	0	100	100
1	Ν	177/200~(88%)	174 (98%)	3 (2%)	0	100	100
All	All	2456/2800~(88%)	2413 (98%)	43 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentile	s
1	А	145/167~(87%)	142 (98%)	3~(2%)	53 65	
1	В	144/167~(86%)	140 (97%)	4 (3%)	43 53	
1	С	141/167~(84%)	138 (98%)	3 (2%)	53 65	
1	D	145/167~(87%)	143 (99%)	2 (1%)	67 78	
1	Ε	146/167~(87%)	144 (99%)	2(1%)	67 78	
1	F	145/167~(87%)	143 (99%)	2(1%)	67 78	
1	G	142/167~(85%)	139~(98%)	3~(2%)	53 65	



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	Н	144/167~(86%)	142~(99%)	2(1%)	67 78
1	Ι	146/167~(87%)	142~(97%)	4(3%)	44 55
1	J	138/167~(83%)	134~(97%)	4(3%)	42 52
1	Κ	138/167~(83%)	136~(99%)	2(1%)	67 78
1	L	136/167~(81%)	134~(98%)	2(2%)	65 76
1	М	143/167~(86%)	141~(99%)	2(1%)	67 78
1	Ν	144/167~(86%)	142~(99%)	2(1%)	67 78
All	All	1997/2338~(85%)	1960~(98%)	37~(2%)	57 68

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	145	LYS
1	Ν	123	HIS
1	Κ	123	HIS
1	L	126	LEU
1	Е	123	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	Ι	54	GLN
1	М	130	GLN
1	J	54	GLN
1	N	54	GLN
1	L	54	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	6V8	Κ	301	1	$19,\!23,\!23$	0.64	0	$24,\!31,\!31$	1.32	3 (12%)
2	6V8	Ι	301	1	$19,\!23,\!23$	0.87	1 (5%)	24,31,31	1.87	4 (16%)
2	6V8	F	301	1	$19,\!23,\!23$	0.79	1 (5%)	24,31,31	1.52	5 (20%)
2	6V8	D	301	1	19,23,23	1.02	2 (10%)	24,31,31	2.15	5 (20%)
2	6V8	G	301	1	19,23,23	0.89	2 (10%)	24,31,31	1.73	5 (20%)
2	6V8	Ν	301	1	19,23,23	1.06	2 (10%)	24,31,31	2.51	9 (37%)
2	6V8	J	301	1	19,23,23	0.60	0	24,31,31	1.62	4 (16%)
2	6V8	С	301	1	19,23,23	0.68	0	24,31,31	1.73	7 (29%)
2	6V8	В	301	1	19,23,23	0.73	0	24,31,31	1.53	<u>6 (25%)</u>
2	6V8	А	301	1	19,23,23	0.74	0	24,31,31	1.96	5 (20%)
2	6V8	L	301	1	19,23,23	0.72	1 (5%)	24,31,31	1.31	1 (4%)
2	6V8	М	301	1	19,23,23	0.96	2 (10%)	24,31,31	1.88	<mark>6 (25%)</mark>
2	6V8	Е	301	1	19,23,23	0.63	0	24,31,31	1.32	2 (8%)
2	6V8	Н	301	1	19,23,23	0.67	0	24,31,31	1.48	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6V8	К	301	1	-	4/15/21/21	0/1/1/1
2	6V8	Ι	301	1	-	4/15/21/21	0/1/1/1



8QYF	٦
------	---

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6V8	F	301	1	-	4/15/21/21	0/1/1/1
2	6V8	D	301	1	-	4/15/21/21	0/1/1/1
2	6V8	G	301	1	-	4/15/21/21	0/1/1/1
2	6V8	Ν	301	1	-	8/15/21/21	0/1/1/1
2	6V8	J	301	1	-	1/15/21/21	0/1/1/1
2	6V8	С	301	1	-	3/15/21/21	0/1/1/1
2	6V8	В	301	1	-	3/15/21/21	0/1/1/1
2	6V8	А	301	1	-	4/15/21/21	0/1/1/1
2	6V8	L	301	1	-	4/15/21/21	0/1/1/1
2	6V8	М	301	1	-	3/15/21/21	0/1/1/1
2	6V8	Е	301	1	-	4/15/21/21	0/1/1/1
2	6V8	Н	301	1	-	4/15/21/21	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	301	6V8	C2-C3	2.68	1.43	1.39
2	Ν	301	6V8	C2-C3	2.53	1.43	1.39
2	Ν	301	6V8	C2-C7	2.41	1.55	1.50
2	G	301	6V8	C2-C3	2.29	1.42	1.39
2	М	301	6V8	C2-C3	2.24	1.42	1.39

The worst 5 of 67 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	N	301	6V8	C10-N9-C7	6.13	135.91	121.33
2	А	301	6V8	C21-C22-C23	5.92	122.83	115.39
2	D	301	6V8	C2-C3-CL3	5.79	129.56	121.00
2	Ι	301	6V8	C2-C3-CL3	5.64	129.34	121.00
2	N	301	6V8	C2-C3-CL3	5.10	128.55	121.00

There are no chirality outliers.

5 of 54 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	301	6V8	C21-C22-C23-C25
2	А	301	6V8	C21-C22-C23-C24
2	В	301	6V8	C21-C22-C23-C25
2	В	301	6V8	C21-C22-C23-C24



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	С	301	6V8	C21-C22-C23-C25


There are no ring outliers.

6 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Κ	301	6V8	1	0
2	Ι	301	6V8	1	0
2	С	301	6V8	2	0
2	В	301	6V8	1	0
2	L	301	6V8	1	0
2	Ε	301	6V8	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	181/200~(90%)	-0.26	2 (1%) 80 86	34, 42, 67, 78	0
1	В	178/200~(89%)	-0.37	1 (0%) 89 93	34, 42, 71, 84	0
1	С	179/200~(89%)	-0.36	0 100 100	34, 42, 70, 78	0
1	D	179/200~(89%)	-0.29	1 (0%) 89 93	34, 42, 66, 78	0
1	Е	179/200~(89%)	-0.24	0 100 100	34, 42, 68, 81	0
1	F	179/200~(89%)	-0.25	0 100 100	34, 42, 70, 87	0
1	G	179/200~(89%)	-0.38	0 100 100	34, 42, 66, 78	0
1	Н	179/200~(89%)	-0.18	3 (1%) 70 78	34, 42, 70, 103	0
1	Ι	179/200~(89%)	-0.26	1 (0%) 89 93	34, 42, 68, 83	0
1	J	179/200~(89%)	-0.34	0 100 100	34, 42, 68, 81	0
1	Κ	178/200~(89%)	-0.30	5 (2%) 53 63	34, 42, 66, 78	0
1	L	178/200~(89%)	-0.41	1 (0%) 89 93	34, 42, 66, 97	0
1	М	179/200~(89%)	-0.28	1 (0%) 89 93	34, 42, 68, 78	0
1	Ν	181/200~(90%)	-0.34	1 (0%) 89 93	34, 42, 68, 81	0
All	All	2507/2800~(89%)	-0.30	16 (0%) 89 93	34, 42, 68, 103	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	6	THR	3.2
1	В	18	TYR	3.1
1	Н	18	TYR	3.0
1	Н	7	VAL	2.9
1	D	18	TYR	2.9



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\operatorname{\AA}^2)$	Q < 0.9
2	6V8	Е	301	23/23	0.84	0.16	$26,\!39,\!72,\!99$	23
2	6V8	N	301	23/23	0.86	0.16	$37,\!50,\!114,\!128$	0
2	6V8	Н	301	23/23	0.88	0.14	21,42,78,100	23
2	6V8	А	301	23/23	0.88	0.17	$27,\!50,\!79,\!109$	0
2	6V8	С	301	23/23	0.89	0.12	$28,\!53,\!86,\!105$	0
2	6V8	L	301	23/23	0.90	0.17	37,47,73,99	23
2	6V8	D	301	23/23	0.91	0.16	37,51,98,113	0
2	6V8	Ι	301	23/23	0.91	0.11	32,53,98,105	0
2	6V8	В	301	23/23	0.91	0.17	$26,\!41,\!84,\!127$	23
2	6V8	G	301	23/23	0.91	0.13	36,46,92,127	0
2	6V8	F	301	23/23	0.92	0.11	31,50,89,116	0
2	6V8	М	301	23/23	0.92	0.12	35,54,83,111	0
2	6V8	Κ	301	23/23	0.92	0.15	34,55,98,115	23
2	6V8	J	301	23/23	0.94	0.12	35,56,118,131	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




































6.5 Other polymers (i)

There are no such residues in this entry.

