



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2024 – 02:10 am GMT

PDB ID : 6R2B  
Title : Crystal structure of the SucA domain of Mycobacterium smegmatis KGD after soaking with succinylphosphonate  
Authors : Wagner, T.; Alzari, P.M.; Bellinzoni, M.  
Deposited on : 2019-03-15  
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	868	 3% 95%
1	B	868	 2% 94%
1	C	868	 3% 94%
1	D	868	 3% 93%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 27742 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Multifunctional 2-oxoglutarate metabolism enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	844	6550	4119	1159	1248	24	0	1	0
1	B	840	6561	4127	1159	1251	24	0	2	0
1	C	844	6575	4138	1164	1249	24	0	1	0
1	D	839	6540	4116	1159	1241	24	0	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	360	GLY	-	expression tag	UNP A0R2B1
B	360	GLY	-	expression tag	UNP A0R2B1
C	360	GLY	-	expression tag	UNP A0R2B1
D	360	GLY	-	expression tag	UNP A0R2B1

- Molecule 2 is (4 {S})-4-[(2 {R})-3-[(4-azanyl-2-methyl-pyrimidin-5-yl)methyl]-4-methyl-5-[2-[oxidanyl(phosphonoxy)phosphoryl]oxyethyl]-2 {H}-1,3-thiazol-2-yl]-4-oxidanyl-4-phosphono-butanoic acid (three-letter code: QSP) (formula: C<sub>16</sub>H<sub>27</sub>N<sub>4</sub>O<sub>13</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	37	16	4	13	3	1	0	0
2	B	1	37	16	4	13	3	1	0	0
2	C	1	37	16	4	13	3	1	0	0
2	D	1	37	16	4	13	3	1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	1	1	1	0	0
3	B	1	1	1	0	0
3	C	1	1	1	0	0
3	D	1	1	1	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	A	1	1	1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ca 1 1	0	0
4	C	1	Total Ca 1 1	0	0
4	D	1	Total Ca 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	326	Total O 326 326	0	0
5	B	351	Total O 351 351	0	0
5	C	355	Total O 355 355	0	0
5	D	328	Total O 328 328	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme



- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme




- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme



- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.12Å 82.76Å 162.97Å 98.55° 97.52° 102.18°	Depositor
Resolution (Å)	158.87 – 1.96 158.87 – 1.96	Depositor EDS
% Data completeness (in resolution range)	97.1 (158.87-1.96) 97.1 (158.87-1.96)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 1.97Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.228 , 0.245 0.226 , 0.244	Depositor DCC
$R_{free}$ test set	14370 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.0	Xtrriage
Anisotropy	0.477	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.000 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	27742	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.06 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.0215e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QSP, CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/6684	0.60	0/9058
1	B	0.50	0/6700	0.60	0/9074
1	C	0.50	0/6713	0.60	0/9099
1	D	0.50	0/6678	0.60	0/9045
All	All	0.50	0/26775	0.60	0/36276

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6550	0	6354	11	0
1	B	6561	0	6412	13	0
1	C	6575	0	6399	14	0
1	D	6540	0	6373	15	0
2	A	37	0	0	0	0
2	B	37	0	0	0	0
2	C	37	0	0	0	0
2	D	37	0	0	0	0
3	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	326	0	0	0	0
5	B	351	0	0	0	0
5	C	355	0	0	1	0
5	D	328	0	0	0	0
All	All	27742	0	25538	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1112:LEU:HD21	1:D:1155:LEU:HD22	1.82	0.62
1:D:794:ARG:HA	1:D:797:GLN:HG2	1.83	0.60
1:C:492:PHE:HD2	1:C:554:TYR:HE1	1.52	0.56
1:B:492:PHE:HD1	1:B:554:TYR:HE1	1.57	0.52
1:C:673:HIS:HB3	5:C:2444:HOH:O	2.09	0.51
1:A:492:PHE:HD1	1:A:554:TYR:HE1	1.59	0.50
1:C:749:GLU:CD	1:C:749:GLU:H	2.14	0.50
1:A:1112:LEU:HD21	1:A:1155:LEU:HD22	1.94	0.50
1:A:981:VAL:HG22	1:A:988:ILE:HD11	1.94	0.48
1:B:696:THR:HG21	1:B:738:ASP:HB2	1.96	0.48
1:D:981:VAL:HG22	1:D:988:ILE:HD11	1.97	0.47
1:B:981:VAL:HG22	1:B:988:ILE:HD11	1.96	0.46
1:C:817:PRO:HB3	1:D:1217:GLU:HG2	1.98	0.46
1:B:778:LEU:HB3	1:B:784:ILE:HG12	1.97	0.46
1:C:981:VAL:HG22	1:C:988:ILE:HD11	1.98	0.46
1:D:492:PHE:HD1	1:D:554:TYR:HE2	1.63	0.46
1:B:848:LEU:HD12	1:B:868:ARG:HD3	1.99	0.45
1:B:1069:PRO:HB2	1:B:1072:MET:HB3	1.99	0.45
1:D:848:LEU:HD12	1:D:868:ARG:HD2	1.99	0.44
1:C:778:LEU:HB3	1:C:784:ILE:HG12	1.99	0.44
1:A:634:ARG:HD2	1:D:590:PHE:CD1	2.51	0.44
1:C:696:THR:HG21	1:C:738:ASP:HB2	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:696:THR:HG21	1:A:738:ASP:HB2	1.98	0.44
1:B:1112:LEU:HD21	1:B:1155:LEU:HD22	2.00	0.44
1:B:492:PHE:CD1	1:B:554:TYR:HE1	2.34	0.44
1:A:749:GLU:H	1:A:749:GLU:CD	2.21	0.43
1:C:1069:PRO:HB2	1:C:1072:MET:HB3	1.99	0.43
1:C:784:ILE:HD12	1:C:789:ALA:HB2	2.00	0.43
1:C:749:GLU:CD	1:C:749:GLU:N	2.72	0.43
1:C:848:LEU:HD12	1:C:868:ARG:HD2	2.00	0.43
1:B:628:GLU:HG2	1:B:664:ARG:O	2.19	0.43
1:A:1069:PRO:HB2	1:A:1072:MET:HB3	2.00	0.42
1:A:1069:PRO:CB	1:A:1072:MET:HB3	2.50	0.42
1:B:908:PHE:CZ	1:B:1070:LYS:HG2	2.54	0.42
1:D:1016:GLN:HB3	1:D:1020:HIS:HB2	2.01	0.42
1:D:1069:PRO:CB	1:D:1072:MET:HB3	2.50	0.41
1:A:908:PHE:CZ	1:A:1070:LYS:HG2	2.55	0.41
1:C:1112:LEU:HD23	1:C:1114:LEU:HD11	2.02	0.41
1:D:696:THR:HG21	1:D:738:ASP:HB2	2.03	0.41
1:D:908:PHE:CZ	1:D:1070:LYS:HG2	2.56	0.41
1:D:1069:PRO:HB2	1:D:1072:MET:HB3	2.02	0.41
1:D:628:GLU:HG2	1:D:664:ARG:O	2.21	0.41
1:B:1069:PRO:CB	1:B:1072:MET:HB3	2.51	0.41
1:A:634:ARG:HD2	1:D:590:PHE:HD1	1.85	0.41
1:B:1177:TRP:CD1	1:B:1197:ARG:HD3	2.56	0.41
1:C:1177:TRP:CD1	1:C:1197:ARG:HD3	2.56	0.40
1:A:1217:GLU:HG2	1:B:817:PRO:HB3	2.02	0.40
1:D:1112:LEU:HD12	1:D:1136:ALA:HB3	2.03	0.40
1:C:1069:PRO:CB	1:C:1072:MET:HB3	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	837/868 (96%)	817 (98%)	18 (2%)	2 (0%)	47	38
1	B	836/868 (96%)	814 (97%)	20 (2%)	2 (0%)	47	38
1	C	839/868 (97%)	816 (97%)	21 (2%)	2 (0%)	47	38
1	D	833/868 (96%)	810 (97%)	21 (2%)	2 (0%)	47	38
All	All	3345/3472 (96%)	3257 (97%)	80 (2%)	8 (0%)	47	38

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	538	PRO
1	B	538	PRO
1	B	605	HIS
1	C	538	PRO
1	C	605	HIS
1	D	538	PRO
1	A	605	HIS
1	D	605	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	680/726 (94%)	674 (99%)	6 (1%)	78	77
1	B	689/726 (95%)	685 (99%)	4 (1%)	86	85
1	C	685/726 (94%)	681 (99%)	4 (1%)	86	85
1	D	682/726 (94%)	679 (100%)	3 (0%)	91	90
All	All	2736/2904 (94%)	2719 (99%)	17 (1%)	86	85

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	514	VAL
1	A	682	PHE
1	A	791	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	812	LYS
1	A	953	PHE
1	A	1131	ASN
1	B	373	LEU
1	B	464	GLN
1	B	682	PHE
1	B	953	PHE
1	C	682	PHE
1	C	767	LYS
1	C	812	LYS
1	C	953	PHE
1	D	682	PHE
1	D	953	PHE
1	D	1131	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	976	GLN
1	C	976	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	QSP	B	2001	3	32,38,38	1.98	11 (34%)	44,59,59	1.34	6 (13%)
2	QSP	A	2001	3	32,38,38	1.78	12 (37%)	44,59,59	1.16	3 (6%)
2	QSP	C	2001	3	32,38,38	1.81	11 (34%)	44,59,59	1.26	5 (11%)
2	QSP	D	2001	3	32,38,38	1.71	9 (28%)	44,59,59	1.38	5 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QSP	B	2001	3	-	2/29/54/54	0/2/2/2
2	QSP	A	2001	3	-	2/29/54/54	0/2/2/2
2	QSP	C	2001	3	-	1/29/54/54	0/2/2/2
2	QSP	D	2001	3	-	1/29/54/54	0/2/2/2

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2001	QSP	C4'-N3'	3.75	1.40	1.35
2	B	2001	QSP	C4'-N3'	3.70	1.40	1.35
2	B	2001	QSP	O11-C11	-3.59	1.40	1.43
2	B	2001	QSP	C2-N3	3.58	1.50	1.46
2	B	2001	QSP	C4-N3	3.48	1.45	1.36
2	D	2001	QSP	C4'-N3'	3.47	1.40	1.35
2	D	2001	QSP	C2-S1	3.29	1.90	1.83
2	C	2001	QSP	C4-N3	3.20	1.45	1.36
2	A	2001	QSP	PA-O7	3.07	1.71	1.59
2	B	2001	QSP	C2-S1	3.06	1.89	1.83
2	D	2001	QSP	PA-O7	2.85	1.70	1.59
2	A	2001	QSP	C2-S1	2.78	1.89	1.83
2	B	2001	QSP	C7'-N3	2.77	1.51	1.46
2	C	2001	QSP	C5'-C4'	2.74	1.47	1.42
2	C	2001	QSP	C2-S1	2.72	1.89	1.83
2	D	2001	QSP	C4-N3	2.71	1.43	1.36

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	QSP	C4'-N3'	2.69	1.38	1.35
2	A	2001	QSP	O7-C7	-2.58	1.34	1.44
2	D	2001	QSP	C2'-N1'	2.56	1.38	1.34
2	D	2001	QSP	C6-C5	-2.55	1.45	1.50
2	C	2001	QSP	C7'-N3	2.53	1.50	1.46
2	A	2001	QSP	CM4-C4	2.52	1.53	1.49
2	B	2001	QSP	C13-C14	2.51	1.56	1.50
2	C	2001	QSP	C6-C7	2.50	1.58	1.51
2	C	2001	QSP	PA-O7	2.49	1.69	1.59
2	B	2001	QSP	O7-C7	-2.48	1.34	1.44
2	C	2001	QSP	O7-C7	-2.43	1.34	1.44
2	D	2001	QSP	O7-C7	-2.41	1.34	1.44
2	B	2001	QSP	C5'-C4'	2.41	1.47	1.42
2	A	2001	QSP	C4-N3	2.40	1.42	1.36
2	C	2001	QSP	C13-C14	2.40	1.56	1.50
2	A	2001	QSP	C5-C4	2.40	1.40	1.35
2	D	2001	QSP	C6-C7	2.38	1.57	1.51
2	B	2001	QSP	P4-O8	-2.24	1.50	1.54
2	A	2001	QSP	C6'-N1'	2.23	1.39	1.34
2	A	2001	QSP	O11-C11	-2.22	1.41	1.43
2	D	2001	QSP	P4-O8	-2.19	1.50	1.54
2	A	2001	QSP	C2-N3	2.14	1.48	1.46
2	C	2001	QSP	C2'-N1'	2.13	1.37	1.34
2	B	2001	QSP	P4-O5	2.09	1.53	1.50
2	A	2001	QSP	C6-C5	-2.08	1.46	1.50
2	A	2001	QSP	C6-C7	2.08	1.57	1.51
2	C	2001	QSP	P4-O8	-2.04	1.51	1.54

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2001	QSP	S1-C2-N3	-4.47	103.22	106.91
2	B	2001	QSP	S1-C2-N3	-3.72	103.84	106.91
2	B	2001	QSP	C13-C12-C11	2.99	121.53	114.33
2	C	2001	QSP	C2'-N3'-C4'	-2.82	113.68	118.08
2	D	2001	QSP	C2'-N3'-C4'	-2.74	113.82	118.08
2	D	2001	QSP	CM4-C4-N3	2.60	122.37	119.89
2	B	2001	QSP	C2'-N3'-C4'	-2.50	114.19	118.08
2	D	2001	QSP	O6-P4-O5	2.48	118.63	113.06
2	A	2001	QSP	C13-C12-C11	2.48	120.29	114.33
2	B	2001	QSP	O3B-PB-O2B	2.38	116.75	107.64
2	D	2001	QSP	O3B-PB-O2B	2.38	116.72	107.64

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	QSP	O3B-PB-O2B	2.35	116.61	107.64
2	C	2001	QSP	C13-C12-C11	2.30	119.86	114.33
2	B	2001	QSP	C7'-N3-C2	-2.29	116.94	119.47
2	C	2001	QSP	C5'-C7'-N3	2.28	116.60	113.34
2	C	2001	QSP	S1-C2-N3	-2.25	105.05	106.91
2	C	2001	QSP	O8-P4-O6	2.18	114.14	107.99
2	B	2001	QSP	O6-P4-O5	2.16	117.91	113.06
2	A	2001	QSP	C2'-N3'-C4'	-2.07	114.85	118.08

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2001	QSP	PA-O3A-PB-O2B
2	B	2001	QSP	PA-O3A-PB-O2B
2	B	2001	QSP	PA-O3A-PB-O3B
2	D	2001	QSP	PA-O3A-PB-O2B
2	A	2001	QSP	PA-O3A-PB-O3B
2	C	2001	QSP	PA-O3A-PB-O3B


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	844/868 (97%)	0.11	24 (2%) 53 62	18, 31, 60, 98	0
1	B	840/868 (96%)	0.09	17 (2%) 65 73	18, 30, 58, 89	0
1	C	844/868 (97%)	0.10	24 (2%) 53 62	17, 29, 57, 92	0
1	D	839/868 (96%)	0.16	25 (2%) 50 59	18, 30, 58, 85	0
All	All	3367/3472 (96%)	0.11	90 (2%) 54 63	17, 30, 58, 98	0

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	394	ASN	6.7
1	C	394	ASN	6.4
1	D	412	LEU	6.4
1	C	828	PRO	6.2
1	B	828	PRO	5.7
1	D	410	LEU	5.3
1	D	570	ALA	5.3
1	D	371	ILE	4.9
1	B	502	GLY	4.4
1	D	394	ASN	4.4
1	B	829	SER	4.3
1	A	570	ALA	4.3
1	C	413	TRP	4.2
1	C	790	GLU	4.2
1	A	502	GLY	3.9
1	B	426	GLY	3.8
1	B	570	ALA	3.7
1	C	502	GLY	3.7
1	C	829	SER	3.6
1	D	419	PHE	3.4
1	B	827	ILE	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	1149	ARG	3.3
1	D	796	TYR	3.3
1	D	411	THR	3.2
1	A	796	TYR	3.2
1	A	411	THR	3.2
1	C	501	VAL	3.2
1	B	796	TYR	3.1
1	C	827	ILE	3.1
1	A	368	ALA	2.9
1	A	1102	GLY	2.9
1	B	779	ILE	2.9
1	D	397	PHE	2.8
1	B	790	GLU	2.8
1	C	796	TYR	2.8
1	A	412	LEU	2.8
1	D	502	GLY	2.8
1	C	368	ALA	2.7
1	C	779	ILE	2.7
1	B	501	VAL	2.7
1	D	786	MET	2.7
1	C	417	ARG	2.7
1	A	779	ILE	2.6
1	C	410	LEU	2.6
1	A	746	GLY	2.6
1	B	786	MET	2.6
1	A	563	GLY	2.6
1	A	790	GLU	2.6
1	A	371	ILE	2.5
1	C	626	THR	2.5
1	D	472	ASP	2.4
1	C	472	ASP	2.4
1	A	413	TRP	2.4
1	A	783	ASP	2.4
1	D	392	LEU	2.3
1	D	368	ALA	2.3
1	A	568	SER	2.3
1	C	554	TYR	2.3
1	D	627	GLY	2.3
1	A	501	VAL	2.3
1	D	827	ILE	2.3
1	D	568	SER	2.2
1	C	784	ILE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	797	GLN	2.2
1	B	392	LEU	2.2
1	A	786	MET	2.2
1	D	420	LYS	2.2
1	A	794	ARG	2.2
1	A	807	VAL	2.2
1	C	492	PHE	2.2
1	C	786	MET	2.1
1	B	390	LEU	2.1
1	B	626	THR	2.1
1	B	793	LEU	2.1
1	D	415	LEU	2.1
1	D	779	ILE	2.1
1	C	392	LEU	2.1
1	B	554	TYR	2.1
1	B	804	PHE	2.1
1	C	627	GLY	2.1
1	D	491	ALA	2.1
1	A	830	LYS	2.1
1	D	555	SER	2.1
1	C	371	ILE	2.1
1	A	373	LEU	2.1
1	A	365	ASP	2.0
1	D	492	PHE	2.0
1	C	570	ALA	2.0
1	D	413	TRP	2.0
1	C	415	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	B	2002	1/1	0.95	0.06	19,19,19,19	0
2	QSP	C	2001	37/37	0.96	0.13	14,20,29,31	0
3	MG	A	2002	1/1	0.96	0.09	20,20,20,20	0
2	QSP	B	2001	37/37	0.96	0.12	17,21,30,31	0
3	MG	D	2002	1/1	0.96	0.08	19,19,19,19	0
2	QSP	D	2001	37/37	0.97	0.12	15,21,30,31	0
2	QSP	A	2001	37/37	0.97	0.13	14,22,30,31	0
4	CA	C	2003	1/1	0.97	0.06	26,26,26,26	0
3	MG	C	2002	1/1	0.98	0.07	20,20,20,20	0
4	CA	B	2003	1/1	0.99	0.05	26,26,26,26	0
4	CA	A	2003	1/1	0.99	0.06	28,28,28,28	0
4	CA	D	2003	1/1	0.99	0.05	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around QSP C 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around QSP B 2001:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around QSP D 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.