

Full wwPDB X-ray Structure Validation Report (i)

Nov 6, 2024 - 03:52 am GMT

PDB ID : 8R40

Title: Crystal structure of diabody CR57 in complex with rabies virus protein G

domain III

Authors : Kedari, A.; Rissanen, I.

Deposited on : 2023-11-10

Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	262	84%		8% 8%			
1	R	262	80%		11% 8%			
2	С	122	61%	11%	27%			
2	G	122	57%	11% •	30%			



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Homospecific Diabody CR57.

	\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
Ī	1	D	240	Total	С	N	О	S	0	0	0
	1	π	240	1805	1131	304	362	8	0	U	U
Ī	1	Λ	242	Total	С	N	О	S	0	0	0
	1	Α	242	1816	1137	305	366	8	0	U	U

• Molecule 2 is a protein called Glycoprotein.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	С	85	Total	С	N	О	S	0	0	0
	G	00	647	404	114	120	9	U	U	0
9	C	89	Total	С	N	О	S	0	0	0
		09	681	426	118	128	9	U	U	U

There are 82 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	28	GLU	-	expression tag	UNP O92284
G	29	THR	-	expression tag	UNP O92284
G	30	GLY	-	expression tag	UNP O92284
G	31	GLU	-	expression tag	UNP O92284
G	32	ASP	-	expression tag	UNP O92284
G	33	GLU	-	expression tag	UNP O92284
G	34	GLY	-	expression tag	UNP O92284
G	35	CYS	-	expression tag	UNP O92284
G	36	THR	-	expression tag	UNP O92284
G	37	ASN	-	expression tag	UNP O92284
G	38	LEU	-	expression tag	UNP O92284
G	39	SER	-	expression tag	UNP O92284
G	40	GLU	-	expression tag	UNP O92284
G	41	PHE	-	expression tag	UNP O92284
G	42	SER	-	expression tag	UNP O92284
G	43	TYR	-	expression tag	UNP O92284



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
G	44	MET	-	expression tag	UNP O92284
G	45	GLU	-	expression tag	UNP O92284
G	46	LEU	-	expression tag	UNP O92284
G	47	LYS	-	expression tag	UNP O92284
G	48	VAL	-	expression tag	UNP O92284
G	49	GLY	_	expression tag	UNP O92284
G	50	TYR	-	expression tag	UNP O92284
G	51	ILE	-	expression tag	UNP O92284
G	52	SER	_	expression tag	UNP O92284
G	53	ALA	-	expression tag	UNP O92284
G	176	ILE	_	expression tag	UNP O92284
G	177	LYS	_	expression tag	UNP O92284
G	178	VAL	_	expression tag	UNP O92284
G	179	GLY	_	expression tag	UNP O92284
G	180	GLY	_	expression tag	UNP O92284
G	181	GLY	_	expression tag	UNP O92284
G	263	GLY	_	expression tag	UNP O92284
G	264	THR	_	expression tag	UNP O92284
G	265	LYS	_	expression tag	UNP O92284
G	266	HIS	-	expression tag	UNP O92284
G	267	HIS	-	expression tag	UNP O92284
G	268	HIS	-	expression tag	UNP O92284
G	269	HIS	-	expression tag	UNP O92284
G	270	HIS	-	expression tag	UNP O92284
G	271	HIS	-	expression tag	UNP O92284
С	28	GLU	-	expression tag	UNP O92284
С	29	THR	-	expression tag	UNP O92284
С	30	GLY	-	expression tag	UNP O92284
С	31	GLU	-	expression tag	UNP O92284
С	32	ASP	-	expression tag	UNP O92284
С	33	GLU	-	expression tag	UNP O92284
С	34	GLY	-	expression tag	UNP O92284
С	35	CYS	-	expression tag	UNP O92284
С	36	THR	-	expression tag	UNP O92284
С	37	ASN	-	expression tag	UNP O92284
С	38	LEU		expression tag	UNP O92284
С	39	SER	_	expression tag	UNP O92284
С	40	GLU	-	expression tag	UNP O92284
С	41	PHE	-	expression tag	UNP O92284
С	42	SER	-	expression tag	UNP O92284
С	43	TYR	-	expression tag	UNP O92284
С	44	MET	-	expression tag	UNP O92284



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	45	GLU	-	expression tag	UNP O92284
С	46	LEU	-	expression tag	UNP O92284
С	47	LYS	-	expression tag	UNP O92284
С	48	VAL	-	expression tag	UNP O92284
С	49	GLY	-	expression tag	UNP O92284
С	50	TYR	-	expression tag	UNP O92284
С	51	ILE	-	expression tag	UNP O92284
С	52	SER	-	expression tag	UNP O92284
С	53	ALA	-	expression tag	UNP O92284
С	176	ILE	-	expression tag	UNP O92284
С	177	LYS	ı	expression tag	UNP O92284
С	178	VAL	-	expression tag	UNP O92284
С	179	GLY	-	expression tag	UNP O92284
С	180	GLY	-	expression tag	UNP O92284
С	181	GLY	-	expression tag	UNP O92284
С	263	GLY	-	expression tag	UNP O92284
С	264	THR	-	expression tag	UNP O92284
С	265	LYS	-	expression tag	UNP O92284
С	266	HIS	ı	expression tag	UNP O92284
С	267	HIS	-	expression tag	UNP O92284
С	268	HIS	=	expression tag	UNP O92284
С	269	HIS	-	expression tag	UNP O92284
С	270	HIS	=	expression tag	UNP O92284
С	271	HIS	-	expression tag	UNP O92284

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	R	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	С	1	Total C O 6 3 3	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Homospecific Diabody CR57 Chain R: • Molecule 1: Homospecific Diabody CR57 Chain A: 84% 8% • Molecule 2: Glycoprotein Chain G: 57% 11% 30% GGLN
VAL
AASN
LEU
LEU
HIS
GLY
GLY
HIS
HIS
HIS
HIS
HIS • Molecule 2: Glycoprotein Chain C: 61% 11% 27%







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants	57.20Å 69.72Å 190.73Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.28 - 2.70	Depositor
Resolution (A)	56.28 - 2.70	EDS
% Data completeness	100.0 (56.28-2.70)	Depositor
(in resolution range)	100.0 (56.28-2.70)	EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.76 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487, PHENIX 1.20.1_4487	Depositor
R R.	0.217 , 0.259	Depositor
R, R_{free}	0.219 , 0.257	DCC
R_{free} test set	1111 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	66.4	Xtriage
Anisotropy	0.143	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 58.9	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4973	wwPDB-VP
Average B, all atoms $(Å^2)$	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 22.36 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.7921e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.25	0/1860	0.49	0/2528
1	R	0.26	0/1849	0.49	0/2513
2	С	0.25	0/692	0.53	0/927
2	G	0.26	0/655	0.55	0/873
All	All	0.26	0/5056	0.50	0/6841

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1816	0	1724	11	0
1	R	1805	0	1713	18	0
2	С	681	0	674	7	0
2	G	647	0	643	6	0
3	A	12	0	16	1	0
3	С	6	0	8	0	0
3	R	6	0	8	0	0
All	All	4973	0	4786	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A	A. 0	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:C:44:MET:HB3	2:C:242:ALA:HB3	1.72	0.70
1:R:6:GLN:H	1:R:119:GLN:HE22	1.38	0.69
1:R:51:ILE:HD12	1:R:58:ALA:HB2	1.83	0.60
1:R:168:HIS:HB2	1:R:171:LYS:HE2	1.83	0.60
1:A:174:LYS:HE2	1:A:176:MET:HE1	1.86	0.57
2:C:215:LEU:HD22	2:C:217:LYS:HD3	1.89	0.55
1:R:29:PHE:HB3	1:R:53:PRO:HG2	1.90	0.54
2:G:211:ASP:OD1	2:G:213:ARG:HG2	2.08	0.53
2:C:40:GLU:OE2	2:C:196:ARG:NH1	2.42	0.53
1:R:40:ALA:HB3	1:R:43:GLN:HG3	1.92	0.52
1:A:225:THR:O	2:G:226:LYS:NZ	2.24	0.52
2:C:223:CYS:HB3	2:C:250:LYS:HE3	1.92	0.52
1:A:87:ARG:HG3	1:A:89:ASP:H	1.77	0.50
1:A:164:TRP:HB2	1:A:177:ILE:HB	1.94	0.49
1:R:36:TRP:CD1	1:R:70:ILE:HD12	2.47	0.49
1:R:60:TYR:HE1	1:R:70:ILE:HG12	1.79	0.48
1:A:163:SER:HB3	1:A:218:CYS:HB3	1.96	0.47
1:R:60:TYR:CE1	1:R:70:ILE:HG12	2.49	0.47
1:A:40:ALA:HB3	1:A:43:GLN:HB2	1.95	0.47
2:C:199:ARG:NH2	2:C:206:THR:OG1	2.49	0.47
1:A:-1:THR:HB	1:A:27:GLY:HA2	1.97	0.46
1:R:221:ALA:HB1	1:R:225:THR:HB	1.97	0.46
1:R:88:SER:O	1:R:91:THR:HG22	2.15	0.46
1:R:7:SER:HB3	1:R:21:SER:H	1.81	0.45
2:C:205:LYS:N	2:C:205:LYS:HD3	2.32	0.45
1:A:73:ASP:OD1	1:A:75:SER:OG	2.22	0.43
1:R:54:ILE:HD12	1:R:54:ILE:HA	1.86	0.43
1:A:166:GLN:HB3	1:A:176:MET:HE2	2.01	0.43
2:C:227:LEU:HD23	2:C:227:LEU:HA	1.88	0.43
2:G:221:GLY:HA3	2:G:236:MET:SD	2.59	0.42
1:R:6:GLN:H	1:R:119:GLN:NE2	2.12	0.42
1:R:68:LEU:HD11	1:R:70:ILE:HD11	2.03	0.41
1:A:207:LEU:HD12	1:A:207:LEU:HA	1.86	0.41
2:G:48:VAL:HG21	2:G:53:ALA:HA	2.02	0.41
2:G:198:LYS:HB2	2:G:210:VAL:HG22	2.03	0.41
1:R:46:GLU:HG3	3:A:301:GOL:H31	2.02	0.41
1:R:165:TYR:OH	1:A:112:GLY:O	2.39	0.41
2:G:196:ARG:HD2	2:G:196:ARG:N	2.36	0.41



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ (\rm \AA) \end{array}$	Clash overlap (Å)
1:R:38:ARG:HB3	1:R:48:MET:SD	2.62	0.40
1:R:190:ARG:H	1:R:190:ARG:HG2	1.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$240/262 \ (92\%)$	233 (97%)	7 (3%)	0	100	100
1	R	238/262 (91%)	235 (99%)	3 (1%)	0	100	100
2	C	85/122 (70%)	79 (93%)	6 (7%)	0	100	100
2	G	77/122 (63%)	73 (95%)	4 (5%)	0	100	100
All	All	640/768~(83%)	620 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	195/212 (92%)	192 (98%)	3 (2%)	60 83		
1	R	194/212 (92%)	190 (98%)	4 (2%)	48 76		
2	С	75/103 (73%)	73 (97%)	2 (3%)	40 69		



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
2	G	71/103 (69%)	64 (90%)	7 (10%)	6 16		
All	All	535/630~(85%)	519 (97%)	16 (3%)	36 65		

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	33	THR
1	R	65	GLN
1	R	87	ARG
1	R	135	ARG
1	A	1	GLU
1	A	185	SER
1	A	194	SER
2	G	41	PHE
2	G	42	SER
2	G	189	CYS
2	G	191	ILE
2	G	195	SER
2	G	196	ARG
2	G	226	LYS
2	С	41	PHE
2	С	213	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res Link		Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	GOL	С	301	-	5,5,5	0.92	0	5,5,5	0.97	0
3	GOL	R	301	-	5,5,5	0.93	0	5,5,5	0.97	0
3	GOL	A	301	-	5,5,5	0.92	0	5,5,5	0.99	0
3	GOL	A	302	-	5,5,5	0.94	0	5,5,5	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	С	301	-	=	0/4/4/4	-
3	GOL	R	301	-	-	0/4/4/4	-
3	GOL	A	301	-	=	0/4/4/4	-
3	GOL	A	302	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will




also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9	
1	A	242/262 (92%)	0.03	3 (1%)	76	76	43, 65, 99, 140	0
1	R	240/262 (91%)	0.12	4 (1%)	69	68	48, 70, 124, 161	0
2	С	89/122 (72%)	0.51	2 (2%)	62	61	61, 99, 132, 150	0
2	G	85/122 (69%)	0.66	7 (8%)	19	17	70, 105, 132, 140	0
All	All	656/768~(85%)	0.21	16 (2%)	59	58	43, 74, 127, 161	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	53	ALA	4.7
1	A	240	GLY	3.4
2	G	48	VAL	2.8
2	G	227	LEU	2.7
2	С	53	ALA	2.7
2	G	244	GLN	2.7
2	G	230	VAL	2.5
1	R	240	GLY	2.5
1	R	29	PHE	2.5
1	A	55	PHE	2.5
2	G	191	ILE	2.5
2	G	245	THR	2.4
1	A	155	ASP	2.2
1	R	88	SER	2.2
1	R	64	PHE	2.1
2	С	213	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)


There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	GOL	С	301	6/6	0.73	0.19	74,90,98,100	0
3	GOL	A	302	6/6	0.77	0.15	65,74,94,101	0
3	GOL	R	301	6/6	0.81	0.14	70,75,82,84	0
3	GOL	A	301	6/6	0.90	0.14	61,74,76,80	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around GOL A 302:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GOL R 301:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

