



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 19, 2024 – 03:11 pm GMT

PDB ID : 8R7C  
Title : MutSbeta bound to compound CHDI-00915542 in the canonical DNA-mismatch bound form  
Authors : Thomsen, M.; Thieulin-Pardo, G.; Steinbacher, S.; Brace, G.; Tillack, K.; Johnson, P.; Ritzefeld, M.; Schaertl, S.; Frush, E.; Warfield, B.; Ballantyne, G.; Lee, J.-H.; Witte, D.; Finley, M.; Prasad, B.; Monteagudo, E.; Plotnikov, N.; Pacifici, R.; Maillard, M.; Wilkinson, H.; Iyer, R.; Dominguez, C.; Vogt, T.; Felsenfeld, D.; Haque, T.  
Deposited on : 2023-11-24  
Resolution : 2.82 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4293 (2.84-2.80)
Clashscore	180529	4801 (2.84-2.80)
Ramachandran outliers	177936	4739 (2.84-2.80)
Sidechain outliers	177891	4741 (2.84-2.80)
RSRZ outliers	164620	4295 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	930	 94%
1	E	930	 91% 7%
2	B	918	 89% 7%

*Continued on next page...*

Ideal geometry (proteins) : Engh & Huber (2001)  
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.39

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	918	<p>% 89% 5% 5%</p>
3	C	24	<p>79% 8% 12%</p>
3	G	24	<p>83% 12%</p>
4	D	24	<p>96%</p>
5	H	24	<p>96%</p>

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 29762 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA mismatch repair protein Msh2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	894	Total	C	N	O	S	248	3	0
			7089	4498	1202	1351	38			
1	E	869	Total	C	N	O	S	330	0	0
			6879	4370	1166	1308	35			

- Molecule 2 is a protein called DNA mismatch repair protein Msh3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	852	Total	C	N	O	S	256	1	0
			6826	4354	1166	1274	32			
2	F	868	Total	C	N	O	S	151	1	0
			6958	4437	1193	1297	31			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	217	GLY	-	expression tag	UNP P20585
B	218	PRO	-	expression tag	UNP P20585
B	949	ARG	GLN	conflict	UNP P20585
F	217	GLY	-	expression tag	UNP P20585
F	218	PRO	-	expression tag	UNP P20585
F	949	ARG	GLN	conflict	UNP P20585

- Molecule 3 is a DNA chain called DNA (5'-D(P\*CP\*TP\*AP\*TP\*CP\*TP\*GP\*AP\*AP\*GP\*CP\*CP\*GP\*AP\*TP\*CP\*GP\*AP\*TP\*GP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	21	Total	C	N	O	P	0	0	0
			432	205	80	126	21			
3	G	21	Total	C	N	O	P	0	0	0
			432	205	80	126	21			


- Molecule 4 is a DNA chain called DNA (5'-D(\*CP\*CP\*AP\*TP\*CP\*GP\*AP\*TP\*CP\*GP\*CP\*AP\*GP\*CP\*TP\*TP\*CP\*AP\*GP\*AP\*TP\*AP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	23	466	223	86	135	22	0	0	0

- Molecule 5 is a DNA chain called DNA (5'-D(P\*CP\*AP\*TP\*CP\*GP\*AP\*TP\*CP\*GP\*CP\*AP\*GP\*CP\*TP\*TP\*CP\*AP\*GP\*AP\*TP\*AP\*GP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	H	23	472	224	88	137	23	0	0	0

- Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
6	A	1	27	10	5	10	2	0	0
6	E	1	27	10	5	10	2	0	0

- Molecule 7 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 8 is {N}-(4-chlorophenyl)-2-(4-ethylpiperazin-1-yl)pteridin-4-amine (three-letter code: YKW) (formula: C<sub>18</sub>H<sub>20</sub>ClN<sub>7</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	Cl	N	1	0
			26	18	1	7		
8	F	1	Total	C	Cl	N	0	0
			26	18	1	7		

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	1	Total Cl 1 1	0	0
9	F	2	Total Cl 2 2	0	0

- Molecule 10 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	B	1	Total O S 5 4 1	0	0
10	B	1	Total O S 5 4 1	0	0
10	F	1	Total O S 5 4 1	0	0
10	F	1	Total O S 5 4 1	0	0

- Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	18	Total O 18 18	0	0
11	B	16	Total O 16 16	0	0
11	E	6	Total O 6 6	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
11	F	28	Total 28	O 28	0	0
11	H	1	Total 1	O 1	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA mismatch repair protein Msh2




- Molecule 1: DNA mismatch repair protein Msh2







- Molecule 2: DNA mismatch repair protein Msh3







- Molecule 2: DNA mismatch repair protein Msh3



- Molecule 3: DNA (5'-D(P\*CP\*TP\*AP\*TP\*CP\*TP\*GP\*AP\*AP\*GP\*CP\*CP\*GP\*AP\*TP\*C P\*GP\*AP\*TP\*GP\*G)-3')



- Molecule 3: DNA (5'-D(P\*CP\*TP\*AP\*TP\*CP\*TP\*GP\*AP\*AP\*GP\*CP\*CP\*GP\*AP\*TP\*C P\*GP\*AP\*TP\*GP\*G)-3')



- Molecule 4: DNA (5'-D(\*CP\*CP\*AP\*TP\*CP\*GP\*AP\*TP\*CP\*GP\*CP\*AP\*GP\*CP\*TP\*TP \*CP\*AP\*GP\*AP\*TP\*AP\*G)-3')



- Molecule 5: DNA (5'-D(P\*CP\*AP\*TP\*CP\*GP\*AP\*TP\*CP\*GP\*CP\*AP\*GP\*CP\*TP\*TP\*C P\*AP\*GP\*AP\*TP\*AP\*G)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.22Å 104.79Å 121.26Å 110.03° 91.20° 110.50°	Depositor
Resolution (Å)	112.46 – 2.82 112.46 – 2.82	Depositor EDS
% Data completeness (in resolution range)	70.1 (112.46-2.82) 70.2 (112.46-2.82)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 2.82Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.201 , 0.231 0.205 , 0.234	Depositor DCC
$R_{free}$ test set	5160 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	68.5	Xtrriage
Anisotropy	0.026	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 66.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	29762	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, SO4, YKW, CL, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	0/7200	0.78	0/9698
1	E	0.69	0/6985	0.77	0/9410
2	B	0.69	0/6949	0.78	0/9377
2	F	0.68	0/7084	0.79	0/9560
3	C	0.29	0/484	0.77	0/745
3	G	0.28	0/484	0.74	0/745
4	D	0.35	0/522	0.77	0/803
5	H	0.37	0/529	0.79	0/814
All	All	0.67	0/30237	0.78	0/41152

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7089	0	7136	11	0
1	E	6879	0	6935	8	0
2	B	6826	0	6938	13	0
2	F	6958	0	7071	20	0
3	C	432	0	237	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	432	0	237	1	0
4	D	466	0	260	0	0
5	H	472	0	259	0	0
6	A	27	0	12	0	0
6	E	27	0	12	0	0
7	A	10	0	14	0	0
8	B	26	0	0	0	0
8	F	26	0	0	0	0
9	B	1	0	0	0	0
9	F	2	0	0	0	0
10	B	10	0	0	0	0
10	F	10	0	0	0	0
11	A	18	0	0	0	0
11	B	16	0	0	0	0
11	E	6	0	0	0	0
11	F	28	0	0	0	0
11	H	1	0	0	0	0
All	All	29762	0	29111	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:741[B]:LYS:CD	1:A:741[B]:LYS:H	2.07	0.67
1:A:741[B]:LYS:H	1:A:741[B]:LYS:HD2	1.67	0.58
2:F:378:VAL:HG13	2:F:379:ARG:HD2	1.87	0.56
2:B:884:LEU:HB3	2:B:1004:LEU:HD22	1.87	0.55
2:B:846:TYR:HB3	2:B:921:TYR:HB3	1.89	0.55

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	885/930 (95%)	841 (95%)	44 (5%)	0	100	100
1	E	853/930 (92%)	816 (96%)	37 (4%)	0	100	100
2	B	841/918 (92%)	806 (96%)	35 (4%)	0	100	100
2	F	859/918 (94%)	824 (96%)	34 (4%)	1 (0%)	48	76
All	All	3438/3696 (93%)	3287 (96%)	150 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	385	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	779/804 (97%)	771 (99%)	8 (1%)	73	91
1	E	757/804 (94%)	748 (99%)	9 (1%)	67	89
2	B	763/818 (93%)	754 (99%)	9 (1%)	67	89
2	F	776/818 (95%)	759 (98%)	17 (2%)	47	78
All	All	3075/3244 (95%)	3032 (99%)	43 (1%)	62	86

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	377	ASN
2	F	855	ARG
2	F	379	ARG
2	F	516	GLU
2	F	938	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	462	GLN
2	B	422	GLN
1	E	824	GLN
1	E	835	ASN
2	F	377	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	ADP	A	1001	-	24,29,29	0.64	0	29,45,45	0.84	1 (3%)
6	ADP	E	1001	-	24,29,29	0.67	0	29,45,45	0.76	0
10	SO4	B	1203	-	4,4,4	0.40	0	6,6,6	0.07	0
10	SO4	F	1204	-	4,4,4	0.37	0	6,6,6	0.06	0
8	YKW	F	1201	-	28,29,29	0.33	0	34,40,40	0.48	0
8	YKW	B	1201	-	28,29,29	0.47	0	34,40,40	0.53	0
10	SO4	F	1202	-	4,4,4	0.39	0	6,6,6	0.09	0
10	SO4	B	1204	-	4,4,4	0.39	0	6,6,6	0.05	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	PGE	A	1002	-	9,9,9	0.30	0	8,8,8	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADP	A	1001	-	-	2/12/32/32	0/3/3/3
6	ADP	E	1001	-	-	0/12/32/32	0/3/3/3
8	YKW	F	1201	-	-	0/10/20/20	0/4/4/4
8	YKW	B	1201	-	-	2/10/20/20	0/4/4/4
7	PGE	A	1002	-	-	1/7/7/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1001	ADP	PA-O3A-PB	-2.16	125.42	132.83

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1001	ADP	C5'-O5'-PA-O1A
8	B	1201	YKW	C12-C5-N3-C6
8	B	1201	YKW	N2-C5-N3-C6
7	A	1002	PGE	O2-C3-C4-O3
6	A	1001	ADP	C5'-O5'-PA-O3A


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the



average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	894/930 (96%)	-0.12	6 (0%) 84 79	28, 67, 112, 144	77 (8%)
1	E	869/930 (93%)	-0.04	9 (1%) 79 73	28, 73, 117, 154	92 (10%)
2	B	852/918 (92%)	0.02	10 (1%) 76 70	24, 75, 118, 167	80 (9%)
2	F	868/918 (94%)	-0.29	11 (1%) 74 68	26, 56, 102, 160	48 (5%)
3	C	21/24 (87%)	-0.36	0 100 100	56, 96, 183, 192	0
3	G	21/24 (87%)	-0.31	0 100 100	40, 81, 153, 163	0
4	D	23/24 (95%)	-0.46	0 100 100	47, 87, 157, 195	0
5	H	23/24 (95%)	-0.49	0 100 100	41, 77, 128, 161	0
All	All	3571/3792 (94%)	-0.11	36 (1%) 79 73	24, 68, 117, 195	297 (8%)

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	547	ASN	4.4
1	A	547	ASN	3.6
2	B	771	VAL	3.4
1	E	296	PHE	3.2
1	A	751	GLY	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
10	SO4	B	1203	5/5	0.81	0.09	120,122,124,126	0
7	PGE	A	1002	10/10	0.89	0.12	55,62,64,65	0
10	SO4	B	1204	5/5	0.90	0.09	121,121,123,128	0
8	YKW	B	1201	26/26	0.93	0.10	77,85,95,97	1
10	SO4	F	1204	5/5	0.93	0.11	92,94,97,98	0
10	SO4	F	1202	5/5	0.94	0.11	64,66,68,72	0
8	YKW	F	1201	26/26	0.95	0.07	47,49,61,68	0
6	ADP	E	1001	27/27	0.96	0.06	50,59,64,65	0
6	ADP	A	1001	27/27	0.96	0.06	48,59,64,68	0
9	CL	F	1203	1/1	0.97	0.07	58,58,58,58	0
9	CL	B	1202	1/1	0.99	0.03	41,41,41,41	0
9	CL	F	1205	1/1	0.99	0.02	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around YKW B 1201:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around YKW F 1201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.