



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 06:58 AM BST

PDB ID : 4RNC  
Title : Crystal structure of an esterase RhEst1 from Rhodococcus sp. ECU1013  
Authors : Dou, S.; Kong, X.D.; Xu, J.H.; Zhou, J.  
Deposited on : 2014-10-23  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	
1	B	310	
1	C	310	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Esterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	281	Total 2045	C 1280	N 363	O 396	S 6	0	0	0
1	B	275	Total 2009	C 1257	N 356	O 390	S 6	0	0	0
1	C	275	Total 2009	C 1257	N 356	O 390	S 6	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 4	O 3	P 1	0	0
2	B	1	Total 4	O 3	P 1	0	0
2	C	1	Total 4	O 3	P 1	0	0

- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	245	Total 245	O 245	0	0
3	B	201	Total 201	O 201	0	0
3	C	151	Total 151	O 151	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	212.93Å 45.38Å 77.44Å 90.00° 105.82° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 44.31 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-1.95) 99.9 (44.31-1.95)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.161 , 0.212 0.171 , 0.223	Depositor DCC
$R_{free}$ test set	2674 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.0	Xtrriage
Anisotropy	0.348	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 39.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6672	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.87	0/2086	0.90	3/2839 (0.1%)
1	B	0.83	0/2050	0.94	7/2791 (0.3%)
1	C	0.80	0/2050	0.84	2/2791 (0.1%)
All	All	0.84	0/6186	0.89	12/8421 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	ARG	NE-CZ-NH2	-8.59	116.00	120.30
1	B	196	ARG	NE-CZ-NH1	8.45	124.52	120.30
1	B	196	ARG	NE-CZ-NH2	-8.14	116.23	120.30
1	B	211	ARG	NE-CZ-NH2	-7.26	116.67	120.30
1	B	74	ASP	CB-CG-OD2	-7.05	111.95	118.30
1	A	196	ARG	NE-CZ-NH1	7.05	123.82	120.30
1	B	192	ARG	NE-CZ-NH2	-7.02	116.79	120.30
1	B	211	ARG	NE-CZ-NH1	6.70	123.65	120.30
1	B	192	ARG	NE-CZ-NH1	6.67	123.64	120.30
1	C	192	ARG	NE-CZ-NH1	6.19	123.39	120.30
1	A	211	ARG	NE-CZ-NH2	-6.08	117.26	120.30
1	C	211	ARG	NE-CZ-NH1	5.86	123.23	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	143	ALA	Peptide
1	C	145	ARG	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2045	0	2003	10	0
1	B	2009	0	1961	15	0
1	C	2009	0	1961	10	0
2	A	4	0	0	0	0
2	B	4	0	0	1	0
2	C	4	0	0	0	0
3	A	245	0	0	3	0
3	B	201	0	0	3	0
3	C	151	0	0	2	0
All	All	6672	0	5925	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:140:VAL:HG13	1:B:144:MET:HB3	1.62	0.81
1:B:38:GLN:HG3	3:B:599:HOH:O	1.80	0.80
1:A:211:ARG:NH1	3:A:507:HOH:O	2.10	0.66
1:B:89:GLU:O	3:B:519:HOH:O	2.14	0.65
1:C:140:VAL:CG1	1:C:144:MET:HB3	2.26	0.65
1:A:21:ASN:OD1	3:A:461:HOH:O	2.15	0.63
1:B:273:SER:O	1:B:275:GLY:N	2.27	0.63
1:C:234:LYS:HD2	1:C:247:TYR:CZ	2.40	0.56
1:B:140:VAL:CG1	1:B:144:MET:HB3	2.35	0.55
1:C:175:GLU:OE1	1:C:175:GLU:N	2.40	0.55

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:268:ARG:NH1	3:C:541:HOH:O	2.41	0.52
1:B:140:VAL:HG13	1:B:144:MET:CB	2.38	0.49
1:A:21:ASN:CG	3:A:461:HOH:O	2.49	0.48
1:B:273:SER:C	1:B:275:GLY:H	2.15	0.48
1:A:1:MET:CG	1:B:5:GLU:HG2	2.44	0.48
1:C:101:SER:HA	1:C:127:ALA:HA	1.96	0.47
1:A:21:ASN:OD1	1:A:24:GLY:N	2.48	0.45
1:B:173:PRO:HG2	1:B:177:LYS:CE	2.47	0.44
1:B:173:PRO:HB2	1:B:174:PRO:HA	1.99	0.44
1:A:1:MET:HG3	1:B:5:GLU:HG2	2.00	0.44
1:B:144:MET:CE	2:B:301:PO4:O1	2.66	0.44
1:C:29:LEU:HD11	1:C:54:LEU:HD11	2.00	0.43
1:B:187:TYR:OH	3:B:482:HOH:O	2.15	0.43
1:A:12:THR:HG23	1:A:66:ASP:HB2	1.99	0.43
1:C:261:THR:O	1:C:265:SER:HB2	2.19	0.43
1:A:126:GLY:HA2	1:A:221:HIS:CE1	2.54	0.43
1:B:93:GLU:HA	1:B:118:ALA:HA	1.99	0.42
1:C:144:MET:O	1:C:148:VAL:HG23	2.19	0.42
1:A:140:VAL:CG1	1:A:144:MET:HB3	2.49	0.42
1:A:22:SER:HA	1:A:53:ARG:HD3	2.02	0.42
1:C:268:ARG:HD2	3:C:541:HOH:O	2.19	0.41
1:B:171:THR:O	1:B:173:PRO:HD2	2.21	0.41
1:C:12:THR:OG1	1:C:68:PRO:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	279/310 (90%)	272 (98%)	7 (2%)	0	100 100
1	B	273/310 (88%)	265 (97%)	6 (2%)	2 (1%)	22 11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	273/310 (88%)	263 (96%)	9 (3%)	1 (0%)	34	22
All	All	825/930 (89%)	800 (97%)	22 (3%)	3 (0%)	34	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	274	LEU
1	C	149	PRO
1	B	173	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/231 (89%)	198 (97%)	7 (3%)	37	25
1	B	203/231 (88%)	201 (99%)	2 (1%)	76	74
1	C	203/231 (88%)	202 (100%)	1 (0%)	88	88
All	All	611/693 (88%)	601 (98%)	10 (2%)	62	58

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	SER
1	A	144	MET
1	A	158	GLU
1	A	161	ARG
1	A	170	LEU
1	A	234	LYS
1	A	277	LEU
1	B	152	MET
1	B	173	PRO
1	C	175	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	301	-	0,3,4	0.00	-	0,3,6	0.00	-
2	PO4	C	301	-	0,3,4	0.00	-	0,3,6	0.00	-
2	PO4	B	301	-	0,3,4	0.00	-	0,3,6	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	281/310 (90%)	-0.62	1 (0%) 92 95	22, 28, 44, 60	0
1	B	275/310 (88%)	-0.58	3 (1%) 80 85	23, 31, 49, 85	0
1	C	275/310 (88%)	-0.45	6 (2%) 62 70	26, 34, 62, 102	0
All	All	831/930 (89%)	-0.55	10 (1%) 79 84	22, 32, 54, 102	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	MET	6.3
1	B	275	GLY	3.6
1	C	149	PRO	3.3
1	C	145	ARG	3.3
1	C	142	SER	2.9
1	C	148	VAL	2.7
1	B	173	PRO	2.6
1	A	278	ALA	2.5
1	C	147	ALA	2.2
1	B	274	LEU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	PO4	B	301	4/5	0.96	0.09	41,42,43,47	0
2	PO4	C	301	4/5	0.98	0.07	35,41,42,45	0
2	PO4	A	301	4/5	0.99	0.05	37,40,43,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.