



# Full wwPDB X-ray Structure Validation Report

Jan 31, 2024 – 02:12 PM EST

PDB ID : 7RNC  
Title : Crystal structure of caspase-3 with inhibitor Ac-VDVVD-CHO  
Authors : McCue, W.; Finzel, B.C.  
Deposited on : 2021-07-29  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	 6% 88% 12%
1	C	141	 % 94% 6%
2	B	95	 3% 83% 12% . .
2	D	95	 4% 91% 7% .
3	F	6	 50% 50%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	G	6	 83% 17%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-3 subunit p17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1110	686	201	215	8	1	0	0
1	C	141	1109	687	202	211	9	0	0	0

- Molecule 2 is a protein called Caspase-3 subunit p12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	91	757	499	119	134	5	0	0	0
2	D	93	772	508	123	136	5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	278	HIS	-	expression tag	UNP P42574
D	278	HIS	-	expression tag	UNP P42574

- Molecule 3 is a protein called Ac-VDVVD-CHO.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	6	40	25	5	10	0	0	0
3	G	6	40	25	5	10	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	B	36	Total O 36 36	0	0
4	C	67	Total O 67 67	0	0
4	D	33	Total O 33 33	0	0
4	F	3	Total O 3 3	0	0
4	G	6	Total O 6 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Caspase-3 subunit p17




- Molecule 1: Caspase-3 subunit p17



- Molecule 2: Caspase-3 subunit p12




- Molecule 2: Caspase-3 subunit p12



- Molecule 3: Ac-VDVVD-CHO



- Molecule 3: Ac-VDVVD-CHO



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.25Å 85.22Å 97.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.61 – 1.93 42.61 – 1.93	Depositor EDS
% Data completeness (in resolution range)	96.8 (42.61-1.93) 96.8 (42.61-1.93)	Depositor EDS
$R_{merge}$	0.27	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.30 (at 1.94Å)	Xtrriage
Refinement program	PHENIX 1.19.2	Depositor
R, $R_{free}$	0.234 , 0.284 0.234 , 0.284	Depositor DCC
$R_{free}$ test set	2060 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.0	Xtrriage
Anisotropy	0.818	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4038	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.65 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.6331e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASA, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.38	0/1125	0.58	0/1509
1	C	0.41	0/1124	0.61	0/1506
2	B	0.47	0/780	0.62	1/1053 (0.1%)
2	D	0.43	0/796	0.57	0/1075
3	F	0.35	0/29	0.51	0/40
3	G	0.47	0/29	0.52	0/40
All	All	0.42	0/3883	0.59	1/5223 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	195	TYR	CD1-CE1-CZ	5.12	124.41	119.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1110	0	1093	10	0
1	C	1109	0	1103	7	0
2	B	757	0	740	13	0
2	D	772	0	749	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	40	0	37	3	0
3	G	40	0	37	0	0
4	A	65	0	0	0	0
4	B	36	0	0	2	1
4	C	67	0	0	1	1
4	D	33	0	0	0	0
4	F	3	0	0	0	0
4	G	6	0	0	0	0
All	All	4038	0	3759	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:195:TYR:HE1	2:B:268:MET:CE	1.87	0.88
2:B:271:LYS:NZ	4:B:301:HOH:O	2.14	0.75
2:B:195:TYR:HE1	2:B:268:MET:HE2	1.51	0.74
2:B:195:TYR:CE1	2:B:268:MET:HE1	2.23	0.73
2:D:195:TYR:HE2	2:D:268:MET:HE2	1.55	0.71
2:B:195:TYR:HE1	2:B:268:MET:HE1	1.56	0.70
1:A:59:THR:HG23	1:A:61:MET:H	1.58	0.69
2:B:195:TYR:CE1	2:B:268:MET:CE	2.75	0.69
2:B:186:LYS:N	4:B:302:HOH:O	2.26	0.69
1:C:58:SER:O	4:C:201:HOH:O	2.10	0.68
2:D:195:TYR:HE2	2:D:268:MET:CE	2.10	0.63
1:A:100:MET:HG3	1:A:139:ILE:HG23	1.80	0.62
2:B:207:ARG:HA	2:B:213:SER:HA	1.82	0.60
1:A:56:HIS:O	1:A:59:THR:HG22	2.04	0.57
1:C:93:ARG:HG3	1:C:134:VAL:HG22	1.85	0.57
2:B:207:ARG:HH11	3:F:6:ASA:HB2	1.70	0.56
2:B:210:LYS:HE3	2:B:211:ASP:OD1	2.07	0.55
2:D:207:ARG:HA	2:D:213:SER:HA	1.90	0.53
1:C:168:LEU:HD13	2:D:259:LYS:HE3	1.92	0.52
1:C:117:VAL:HG22	1:C:159:ILE:HB	1.93	0.51
1:C:74:LEU:HD13	1:C:117:VAL:HG11	1.93	0.51
2:D:195:TYR:CE2	2:D:268:MET:CE	2.94	0.49
1:A:120:SER:HB3	1:A:127:ILE:HD11	1.95	0.49
2:B:231:GLU:OE1	2:B:233:MET:HB2	2.13	0.47
1:C:93:ARG:CG	1:C:134:VAL:HG22	2.45	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:LEU:HD13	1:A:117:VAL:HG11	1.96	0.46
3:F:4:VAL:HG22	3:F:5:VAL:H	1.80	0.46
1:A:168:LEU:HD13	2:B:259:LYS:HE3	1.99	0.45
1:C:88:LYS:HA	1:C:88:LYS:HD3	1.79	0.45
1:A:51:ASN:HD22	1:A:89:ASN:HD21	1.64	0.45
1:A:167:GLU:HB2	2:B:203:TYR:HA	1.99	0.45
1:A:151:LEU:HA	1:A:154:LYS:HD2	2.02	0.41
1:A:38:LYS:HD3	1:A:41:TYR:CE2	2.56	0.41
2:D:218:SER:HB3	2:D:242:LYS:HD3	2.03	0.40
3:F:4:VAL:HG22	3:F:5:VAL:N	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:333:HOH:O	4:C:219:HOH:O[3_555]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	136 (98%)	3 (2%)	0	100	100
1	C	139/141 (99%)	137 (99%)	2 (1%)	0	100	100
2	B	89/95 (94%)	88 (99%)	1 (1%)	0	100	100
2	D	91/95 (96%)	91 (100%)	0	0	100	100
3	F	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
3	G	4/6 (67%)	4 (100%)	0	0	100	100
All	All	466/484 (96%)	459 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/127 (98%)	124 (100%)	0	100	100
1	C	124/127 (98%)	124 (100%)	0	100	100
2	B	81/85 (95%)	81 (100%)	0	100	100
2	D	82/85 (96%)	82 (100%)	0	100	100
3	F	4/4 (100%)	4 (100%)	0	100	100
3	G	4/4 (100%)	4 (100%)	0	100	100
All	All	419/432 (97%)	419 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ASA	F	6	1,3	6,7,7	0.94	0	5,8,8	1.34	0
3	ASA	G	6	1,3	6,7,7	0.79	0	5,8,8	1.43	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASA	F	6	1,3	-	2/5/6/6	-
3	ASA	G	6	1,3	-	2/5/6/6	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	6	ASA	OD1-CG-CB	-2.01	116.35	122.80

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	6	ASA	C-CA-CB-CG
3	F	6	ASA	CA-CB-CG-OD1
3	F	6	ASA	CA-CB-CG-OD2
3	G	6	ASA	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	6	ASA	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	141/141 (100%)	0.69	8 (5%) 23 30	20, 28, 45, 54	2 (1%)
1	C	141/141 (100%)	0.51	2 (1%) 75 80	19, 26, 34, 49	0
2	B	91/95 (95%)	0.76	3 (3%) 46 54	17, 23, 33, 40	1 (1%)
2	D	93/95 (97%)	0.71	4 (4%) 35 42	17, 22, 35, 48	0
3	F	4/6 (66%)	0.92	0 100 100	25, 27, 27, 28	0
3	G	4/6 (66%)	0.57	0 100 100	24, 24, 26, 29	0
All	All	474/484 (97%)	0.65	17 (3%) 42 50	17, 25, 39, 54	3 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	174	THR	5.0
1	A	60	GLY	3.8
1	A	174	THR	3.7
1	A	62	THR	3.0
1	A	109	SER	2.9
2	B	264	CYS	2.8
2	B	210	LYS	2.8
2	D	268	MET	2.6
2	D	264	CYS	2.5
1	A	41	TYR	2.5
1	A	121	HIS	2.5
1	A	110	LYS	2.4
2	B	266	VAL	2.3
2	D	266	VAL	2.2
1	C	101	ARG	2.2
2	D	185	HIS	2.2
1	A	101	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ASA	F	6	8/8	0.66	0.27	26,32,38,44	0
3	ASA	G	6	8/8	0.80	0.17	22,24,26,28	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.