



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2023 – 12:35 AM EDT

PDB ID : 7S00  
Title : X-ray structure of the phage AR9 non-virion RNA polymerase core  
Authors : Leiman, P.G.; Sokolova, M.L.; Fraser, A.  
Deposited on : 2021-08-28  
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1149 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	c	496	 9% 96%
1	e	496	 11% 97%
2	C	665	 5% 95%
2	E	665	 8% 96%
3	D	631	 4% 76% 24%
3	F	631	 14% 94% 6%
4	d	448	 17% 89% 8%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	f	448	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '14%', a large green segment in the middle labeled '91%', and a small grey segment on the right labeled '8%'.</p>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 33894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	c	484	Total 4003	C 2580	N 658	O 754	S 11	0	0	0
1	e	484	Total 4003	C 2580	N 658	O 754	S 11	0	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	640	Total 5208	C 3296	N 857	O 1031	S 24	0	1	0
2	E	640	Total 5208	C 3296	N 857	O 1031	S 24	0	1	0

- Molecule 3 is a protein called DNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	481	Total 3893	C 2481	N 640	O 757	S 15	0	0	0
3	F	596	Total 4853	C 3092	N 791	O 952	S 18	0	0	0

- Molecule 4 is a protein called DNA-directed RNA polymerase beta' subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	d	410	Total 3362	C 2182	N 541	O 631	S 8	0	0	0
4	f	410	Total 3362	C 2182	N 541	O 631	S 8	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
d	-21	MET	-	expression tag	UNP A0A172JIH0
d	-20	GLY	-	expression tag	UNP A0A172JIH0
d	-19	SER	-	expression tag	UNP A0A172JIH0
d	-18	SER	-	expression tag	UNP A0A172JIH0
d	-17	HIS	-	expression tag	UNP A0A172JIH0
d	-16	HIS	-	expression tag	UNP A0A172JIH0
d	-15	HIS	-	expression tag	UNP A0A172JIH0
d	-14	HIS	-	expression tag	UNP A0A172JIH0
d	-13	HIS	-	expression tag	UNP A0A172JIH0
d	-12	HIS	-	expression tag	UNP A0A172JIH0
d	-11	SER	-	expression tag	UNP A0A172JIH0
d	-10	SER	-	expression tag	UNP A0A172JIH0
d	-9	GLY	-	expression tag	UNP A0A172JIH0
d	-8	GLU	-	expression tag	UNP A0A172JIH0
d	-7	ASN	-	expression tag	UNP A0A172JIH0
d	-6	LEU	-	expression tag	UNP A0A172JIH0
d	-5	TYR	-	expression tag	UNP A0A172JIH0
d	-4	PHE	-	expression tag	UNP A0A172JIH0
d	-3	GLN	-	expression tag	UNP A0A172JIH0
d	-2	GLY	-	expression tag	UNP A0A172JIH0
d	-1	HIS	-	expression tag	UNP A0A172JIH0
d	0	HIS	-	expression tag	UNP A0A172JIH0
f	-21	MET	-	expression tag	UNP A0A172JIH0
f	-20	GLY	-	expression tag	UNP A0A172JIH0
f	-19	SER	-	expression tag	UNP A0A172JIH0
f	-18	SER	-	expression tag	UNP A0A172JIH0
f	-17	HIS	-	expression tag	UNP A0A172JIH0
f	-16	HIS	-	expression tag	UNP A0A172JIH0
f	-15	HIS	-	expression tag	UNP A0A172JIH0
f	-14	HIS	-	expression tag	UNP A0A172JIH0
f	-13	HIS	-	expression tag	UNP A0A172JIH0
f	-12	HIS	-	expression tag	UNP A0A172JIH0
f	-11	SER	-	expression tag	UNP A0A172JIH0
f	-10	SER	-	expression tag	UNP A0A172JIH0
f	-9	GLY	-	expression tag	UNP A0A172JIH0
f	-8	GLU	-	expression tag	UNP A0A172JIH0
f	-7	ASN	-	expression tag	UNP A0A172JIH0
f	-6	LEU	-	expression tag	UNP A0A172JIH0
f	-5	TYR	-	expression tag	UNP A0A172JIH0
f	-4	PHE	-	expression tag	UNP A0A172JIH0
f	-3	GLN	-	expression tag	UNP A0A172JIH0
f	-2	GLY	-	expression tag	UNP A0A172JIH0
f	-1	HIS	-	expression tag	UNP A0A172JIH0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
f	0	HIS	-	expression tag	UNP A0A172JIH0


- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Zn 1 1	0	0
5	F	1	Total Zn 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase beta subunit



- Molecule 1: DNA-directed RNA polymerase beta subunit




- Molecule 2: DNA-directed RNA polymerase





- Molecule 2: DNA-directed RNA polymerase









• Molecule 3: DNA-directed RNA polymerase





• Molecule 3: DNA-directed RNA polymerase






• Molecule 4: DNA-directed RNA polymerase beta' subunit







● Molecule 4: DNA-directed RNA polymerase beta' subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.86Å 166.27Å 307.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.42 – 3.30 49.42 – 3.30	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.42-3.30) 98.6 (49.42-3.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.59 (at 3.33Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.219 , 0.258 0.220 , 0.256	Depositor DCC
$R_{free}$ test set	4308 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	107.3	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 96.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	33894	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	155.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	c	0.26	0/4082	0.48	0/5502
1	e	0.26	0/4082	0.47	0/5502
2	C	0.27	0/5308	0.48	0/7156
2	E	0.26	0/5308	0.47	0/7156
3	D	0.25	0/3952	0.44	0/5328
3	F	0.25	0/4932	0.44	0/6650
4	d	0.25	0/3431	0.44	0/4630
4	f	0.25	0/3431	0.44	0/4630
All	All	0.26	0/34526	0.46	0/46554

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	c	482/496 (97%)	473 (98%)	9 (2%)	0	100	100
1	e	482/496 (97%)	474 (98%)	8 (2%)	0	100	100
2	C	637/665 (96%)	625 (98%)	12 (2%)	0	100	100
2	E	637/665 (96%)	627 (98%)	10 (2%)	0	100	100
3	D	477/631 (76%)	471 (99%)	6 (1%)	0	100	100
3	F	594/631 (94%)	584 (98%)	10 (2%)	0	100	100
4	d	406/448 (91%)	398 (98%)	8 (2%)	0	100	100
4	f	406/448 (91%)	399 (98%)	6 (2%)	1 (0%)	47	77
All	All	4121/4480 (92%)	4051 (98%)	69 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	f	201	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	c	457/469 (97%)	450 (98%)	7 (2%)	65	81
1	e	457/469 (97%)	454 (99%)	3 (1%)	84	90
2	C	587/608 (96%)	581 (99%)	6 (1%)	76	86
2	E	587/608 (96%)	583 (99%)	4 (1%)	84	90
3	D	445/590 (75%)	441 (99%)	4 (1%)	78	87
3	F	555/590 (94%)	549 (99%)	6 (1%)	73	85
4	d	375/409 (92%)	365 (97%)	10 (3%)	44	71
4	f	375/409 (92%)	375 (100%)	0	100	100
All	All	3838/4152 (92%)	3798 (99%)	40 (1%)	76	86

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	e	373	PHE
3	F	293	ASP
1	e	425	PHE
2	E	516	GLU
3	F	541	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
4	d	15	ASN
4	d	180	ASN
4	d	377	ASN
4	f	247	ASN
4	f	377	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	c	484/496 (97%)	0.57	43 (8%) 9 10	57, 129, 227, 365	0
1	e	484/496 (97%)	0.56	56 (11%) 4 4	109, 173, 251, 319	0
2	C	640/665 (96%)	0.37	31 (4%) 30 28	49, 92, 202, 317	0
2	E	640/665 (96%)	0.51	51 (7%) 12 11	98, 151, 214, 279	0
3	D	481/631 (76%)	0.33	24 (4%) 28 27	54, 112, 194, 261	0
3	F	596/631 (94%)	0.80	87 (14%) 2 2	121, 175, 272, 329	0
4	d	410/448 (91%)	0.94	78 (19%) 1 1	71, 179, 259, 367	0
4	f	410/448 (91%)	0.83	63 (15%) 2 2	123, 196, 278, 427	0
All	All	4145/4480 (92%)	0.60	433 (10%) 6 6	49, 152, 250, 427	0

The worst 5 of 433 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	450	PHE	9.6
4	f	49	ASP	9.1
2	E	557	SER	8.7
3	F	429	VAL	8.5
3	F	491	ILE	7.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	ZN	F	701	1/1	0.71	0.15	188,188,188,188	0
5	ZN	D	701	1/1	0.91	0.17	135,135,135,135	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.